

STABLE POLARIZED DEL PEZZO SURFACES

IVAN CHELTSOV AND JESUS MARTINEZ-GARCIA

ABSTRACT. We give a simple sufficient condition for K -stability of polarized del Pezzo surfaces.

1. INTRODUCTION

In recent years the notion of K -stability has been of great importance in the study of the existence of canonical metrics on complex varieties. This is mainly because of the following

Conjecture (Yau–Tian–Donaldson). *Let X be a smooth variety, and let L be an ample line bundle on X . Then X admits a constant scalar curvature Kähler (cscK) metric in $c_1(L)$ if and only if the pair (X, L) is K -stable.*

In the case of toric surfaces, the conjecture follows from the classical results of Donaldson [9] and recent results of Codogni and Stoppa [7]. It is known in different degrees of generality that K -stability is a necessary condition for the existence of a cscK metric, with the most general result due to Berman, Darvas and Lu [2]. To show that it is also a sufficient condition is currently one of the main open questions in the field. For smooth Fano varieties polarized by anticanonical line bundles, this was recently proved by Chen, Donaldson and Sun in [6].

The goal of this paper is to study K -stability of polarized smooth del Pezzo surfaces. This problem is explicitly solved in the toric case by Donaldson [9]. Surprisingly, we do not know many results about the non-toric case. By the famous theorem of Tian [13], all non-toric smooth del Pezzo surfaces are Kähler-Einstein, so that they are K -stable for the anticanonical polarization. Arezzo and Pacard’s results [1] imply the K -stability for many other polarizations. On the other hand, one can use Ross and Thomas’s [11, Example 5.30] to produce many K -unstable polarizations on every smooth non-toric del Pezzo surface.

The main result of this paper is

Theorem 1.1. *Let S be a smooth del Pezzo surface such that $K_S^2 \leq 2$, and let L be an ample \mathbb{Q} -divisor on S . If $-K_S - \frac{2}{3} \frac{-K_S \cdot L}{L^2} L$ is nef, then (S, L) is K -stable.*

For each smooth del Pezzo surface of degree one or two, this result provides a closed subset in its ample cone that consists of K -stable polarizations. This subset contains an anticanonical divisor, so that Theorem 1.1 also implies the existence of Kähler-Einstein metrics on smooth del Pezzo surfaces of degree one and two. The nefness condition in Theorem 1.1 is easy to check, because the Mori cones of these del Pezzo surfaces are generated by finitely many (-1) -curves.

For smooth del Pezzo surfaces of degree one, the assertion of Theorem 1.1 was recently proved by Hong and Won in [10] using a different approach.

The α -invariant of Tian, originally defined in terms of complex differential geometry, has an algebraic formulation which we use to study the K -stability of the anticanonically polarized del Pezzo surfaces. For a variety X and an ample \mathbb{Q} -divisor L on it, we let

$$\alpha(X, L) = \sup \left\{ \lambda \in \mathbb{Q} \mid \begin{array}{l} \text{the log pair } (X, \lambda D) \text{ is log canonical} \\ \text{for every effective } \mathbb{Q}\text{-divisor } D \sim_{\mathbb{Q}} L \end{array} \right\} \in \mathbb{R}_{>0}.$$

Key words and phrases. K -stability, cscK metric, α -invariant, del Pezzo surface.

While the α -invariant is very difficult to estimate for an arbitrary polarization, it is the only effective invariant to provide sufficient condition for K -stability. In this article we will prove Theorem 1.1 using the following criterion of Dervan:

Theorem 1.2 ([8, Theorem 1.1]). *Let X be a Fano variety with log canonical singularities, and let L be an ample \mathbb{Q} -divisor on it. Let $\nu(L) = \frac{-K_X \cdot L^{n-1}}{L^n}$. Then (X, L) is K -stable provided that*

- (A) *the \mathbb{Q} -divisor $-K_X - \frac{n}{n+1}\nu(L)L$ is nef,*
- (B) *and $\alpha(X, L) > \frac{n}{n+1}\nu(L)$.*

Namely, we will show that for smooth del Pezzo surfaces of degree one and two, the condition (B) in this theorem follows from the condition (A). The reader may wonder whether the condition (B) in Theorem 1.2 is redundant. In general, this is not the case:

Example 1.3. Let S be a smooth cubic surface in \mathbb{P}^3 that does not contain Eckardt points, and let E be a line in S . Let $L = -K_S + xE$, where x is a non-negative rational number such that $x < 1$. Then L is ample and

$$\frac{-K_S \cdot L}{L^2} = \frac{3 + x}{3 + 2x - x^2},$$

so that $-K_S - \frac{2}{3}\frac{-K_S \cdot L}{L^2}L$ is nef if and only if $x \leq \frac{3}{5}$. On the other hand, we have $(1+x)E + C \sim_{\mathbb{Q}} L$, where C is a conic in $| -K_S - E |$ that is tangent to E . This immediately gives $\alpha(S, L) \leq \frac{3}{4+2x}$. Thus, if $\frac{3}{5} \geq x \geq \frac{-1+2\sqrt{10}}{13}$, then $-K_S - \frac{2}{3}\frac{-K_S \cdot L}{L^2}L$ is nef, while $\alpha(S, L) \leq \frac{2}{3}\frac{-K_S \cdot L}{L^2}$.

In this paper, we also prove the following negative result.

Theorem 1.4. *Let S be a smooth del Pezzo surface such that $K_S^2 \geq 4$, and let L be an ample \mathbb{Q} -divisor on S . Then $\alpha(S, L) \leq \frac{2}{3}\frac{-K_S \cdot L}{L^2}$. Moreover, the equality holds if and only if $K_S^2 = 4$ and $L \in \mathbb{Q}_{>0}[-K_S]$.*

Hence Theorem 1.1 exhausts the application of Theorem 1.2 to all smooth del Pezzo surfaces other than cubic surfaces. Fortunately, we can still obtain partial results in the latter case.

Theorem 1.5. *Let S be a smooth cubic surface in \mathbb{P}^3 , and let $E_1, E_2, E_3, E_4, E_5, E_6$ be disjoint lines on the surface S . Let $L = -K_S + x \sum_{i=1}^6 E_i$, where x is a non-negative rational number such that $x < 1$. Then L is ample. Furthermore, if $0 < x \leq \frac{1}{10}$, then (S, L) is K -stable.*

Since all smooth cubic surfaces are Kähler-Einstein by Tian's theorem [13], the K -stability of the pair (S, L) in this theorem holds also for $x = 0$. Moreover, Arezzo and Pacard's results [1] imply the K -stability of the pair (S, L) in the case when x is sufficiently close to 1. Thus, one may expect that the pair (S, L) is K -stable for all positive rational numbers $x < 1$. Unfortunately, we were unable to prove this.

Let us describe the structure of this paper. In Section 2 we present some auxiliary results. In Section 3, we give a very short proof of our Theorem 1.1 for smooth del Pezzo surfaces of degree one. In Section 4, we prove Theorem 1.1 for smooth del Pezzo surfaces of degree two. This section is the main part of the paper. Then we prove Theorem 1.5 in Section 5. Finally, we tackle Theorem 1.4 in Section 6. Its proof involves two very technical inequalities which are left to the Appendix A.

Acknowledgements. We would like to thank Ruadhaí Dervan, Sir Simon Donaldson, Julius Ross, Yanir Rubinshtein and Richard Thomas for helpful discussions and valuable comments.

We proved Theorem 1.1 during our visit to the Simons Center for Geometry and Physics in November 2015. We would like to thank the center for perfect working conditions.

2. PRELIMINARIES

Let S be a smooth surface, let D be an effective \mathbb{Q} -divisor on the surface S , and let P be a point in the surface S . Let $D = \sum_{i=1}^r a_i C_i$, where each C_i is an irreducible curve on S , and each a_i is a non-negative rational number. We assume here that all curves C_1, \dots, C_r are different.

Let $\pi: \tilde{S} \rightarrow S$ be a birational morphism such that \tilde{S} is also smooth. Then π is a composition of n blow ups of smooth points. For each C_i , denote by \tilde{C}_i its proper transform on \tilde{S} . Let F_1, \dots, F_n be π -exceptional curves. Then

$$K_{\tilde{S}} + \sum_{i=1}^r a_i \tilde{C}_i + \sum_{j=1}^n b_j F_j \sim_{\mathbb{Q}} \pi^*(K_S + D)$$

for some rational numbers b_1, \dots, b_n . Suppose, in addition, that $\sum_{i=1}^r \tilde{C}_i + \sum_{j=1}^n F_j$ is a divisor with simple normal crossings (in particular each \tilde{C}_i is smooth).

Definition 2.1. The log pair (S, D) is *log canonical* (respectively *Kawamata log terminal*) at a point $P \in S$ if the following two conditions are satisfied:

- $a_i \leq 1$ (respectively $a_i < 1$) for every C_i such that $P \in C_i$,
- $b_j \leq 1$ (respectively $b_j < 1$) for every F_j such that $\pi(F_j) = P$.

This definition is independent on the choice of birational morphism $\pi: \tilde{S} \rightarrow S$. The log pair (S, D) is said to be *log canonical* (respectively *Kawamata log terminal*) if it is log canonical (respectively, *Kawamata log terminal*) at every point of S .

Remark 2.2. Let R be any effective \mathbb{Q} -divisor on S such that $R \sim_{\mathbb{Q}} D$ and $R \neq D$. Let

$$D_\epsilon = (1 + \epsilon)D - \epsilon R,$$

where $\epsilon \in \mathbb{Q}_{\geq 0}$. Then $D_\epsilon \sim_{\mathbb{Q}} D$. Let ϵ_0 be the greatest rational number such that D_{ϵ_0} is effective. Then the support of D_{ϵ_0} does not contain at least one irreducible component of $\text{Supp}(R)$. Moreover, if the log pair (S, D) is not log canonical at P , and (S, R) is log canonical at P , then the log pair (S, D_{ϵ_0}) is not log canonical at P , because

$$D = \frac{1}{1 + \epsilon_0} D_{\epsilon_0} + \frac{\epsilon_0}{1 + \epsilon_0} R.$$

The following result is well known and is very easy to prove.

Lemma 2.3. *Suppose that (S, D) is not log canonical at P . Then $\text{mult}_P(D) > 1$.*

The following result is also well known (see [12, Corollary 3.12] or [4, Theorem 7]).

Lemma 2.4. *Suppose that (S, D) is not log canonical at P , the curve C_1 is smooth at P , and $a_1 \leq 1$. Let $\Delta = \sum_{i=2}^r a_i C_i$. Then $\text{mult}_P(C_1 \cdot \Delta) > 1$.*

The following result plays an important role in the proof of Theorem 1.1.

Theorem 2.5 ([4, Theorem 13]). *Suppose that (S, D) is not log canonical at P , the curves C_1 and C_2 are smooth at P and intersect each other transversally at P , $a_1 \leq 1$ and $a_2 \leq 1$. Let $\Delta = \sum_{i=3}^r a_i C_i$. If $\text{mult}_P(\Delta) \leq 1$, then $\text{mult}_P(C_1 \cdot \Delta) > 2(1 - a_2)$ or $\text{mult}_P(C_2 \cdot \Delta) > 2(1 - a_1)$.*

Let $\rho: \hat{S} \rightarrow S$ be the blow up of a point $P \in S$, and let F be the ρ -exceptional curve. Denote by \hat{D} the proper transform of the divisor D on the surface \hat{S} via ρ . Then

$$K_{\hat{S}} + \hat{D} + \left(\text{mult}_P(D) - 1 \right) F \sim_{\mathbb{Q}} \rho^*(K_S + D).$$

By Definition 2.1, if (S, D) is not log canonical at P , then $(\hat{S}, \hat{D} + (\text{mult}_P(D) - 1)F)$ is not log canonical at some point of the curve F .

3. DEL PEZZO SURFACES OF DEGREE ONE

Let S be a smooth del Pezzo surface such that $K_S^2 = 1$ and L be an ample \mathbb{Q} -divisor on the surface S . The goal of this section is to prove the following result:

Theorem 3.1. *If $-K_S - \frac{2}{3}\frac{-K_S \cdot L}{L^2}L$ is nef, then $\alpha(S, L) > \frac{2}{3}\frac{-K_S \cdot L}{L^2}$.*

Let us prove this result. Suppose that $-K_S - \frac{2}{3}\frac{-K_S \cdot L}{L^2}L$ is nef. We can swap L with $\frac{2}{3}\frac{-K_S \cdot L}{L^2}L$ and assume that $\frac{-K_S \cdot L}{L^2} = \frac{3}{2}$. Then

$$(3.1) \quad \left(L - \frac{2}{3}(-K_S)\right) \cdot L = 0,$$

and the class $-K_S - L$ is nef. We have to show that $\alpha(S, L) > 1$.

Let $R = -K_S - L$ and $\epsilon = -K_S \cdot R$. Then $\epsilon > 0$, because $R \not\sim_{\mathbb{Q}} 0$ by (3.1). Let

$$(3.2) \quad \gamma = \begin{cases} \frac{6}{5} & \text{if } \epsilon \geq \frac{1}{2}, \\ \frac{3}{3-\epsilon} & \text{if } \epsilon < \frac{1}{2}. \end{cases}$$

Then $\gamma > 1$. We will prove that $\alpha(S, L) \geq \gamma$ by *reductio ad absurdum*.

Suppose that $\alpha(S, L) < \gamma$. Then there exist a positive rational number $\lambda < \gamma$ and an effective \mathbb{Q} -divisor D on S such that $D \sim_{\mathbb{Q}} L$, and $(S, \lambda D)$ is not log canonical at some point $P \in S$.

Lemma 3.2. *One has $L \not\sim_{\mathbb{Q}} -\frac{2}{3}K_S$.*

Proof. By [3, Theorem 1.7], $\alpha(S, -K_S) \geq \frac{5}{6}$. Thus, if $L \sim_{\mathbb{Q}} -\frac{2}{3}K_S$, then $(S, \frac{5}{4}D)$ is log canonical, which is impossible, because $\lambda < \gamma \leq \frac{6}{5} < \frac{5}{4}$ by (3.2). \square

Let C be a curve in the pencil $|-K_S|$ that passes through the point P . Then C is irreducible. Write $D = aC + \Delta$, where a is a non-negative rational number, and Δ is an effective \mathbb{Q} -divisor whose support does not contain the curve C .

Lemma 3.3. *One has $a < \frac{2}{3}$.*

Proof. Suppose that $a \geq \frac{2}{3}$. Then it follows from (3.1) that

$$0 = L \cdot \left(L - \frac{2}{3}(-K_S)\right) = L \cdot \left(D - \frac{2}{3}C\right) = L \cdot \left(\left(a - \frac{2}{3}\right)C + \Delta\right) \geq \left(a - \frac{2}{3}\right)L \cdot C,$$

so that $a = \frac{2}{3}$ and $L \cdot \Delta = 0$, because $L \cdot C > 0$ and $L \cdot \Delta \geq 0$. Then $\Delta = 0$, because L is ample. Thus, we have $L \sim_{\mathbb{Q}} -\frac{2}{3}K_S$, which is impossible by Lemma 3.2. \square

If C is smooth at P , then it follows from Lemma 2.4 that

$$\frac{1}{\lambda} < C \cdot \Delta = C \cdot (D - aC) = 1 - R \cdot C - a \leq 1 - R \cdot C = 1 - \epsilon,$$

which gives $\epsilon < 1$ and $\gamma > \lambda > \frac{1}{1-\epsilon} > \frac{3}{3-\epsilon}$, which contradicts (3.2). Thus, $\text{mult}_P(C) = 2$.

Let $m = \text{mult}_P(\Delta)$. Then $2m \leq C \cdot \Delta = 1 - \epsilon - a < 1 - a$.

Corollary 3.4. *One has $a + 2m < 1$.*

Let $f: \tilde{S} \rightarrow S$ be the blow-up at P , and let F be its exceptional curve. Denote by \tilde{C} and $\tilde{\Delta}$ the proper transforms on the surface \tilde{S} of the curve C and the divisor Δ , respectively. Then the log pair $(\tilde{S}, \lambda a \tilde{C} + \lambda \tilde{\Delta} + (\lambda(2a+m) - 1)F)$ is not log canonical at some point $Q \in F$. Since $a < \frac{2}{3}$ by Lemma 3.3 and $a + 2m < 1$ by Corollary 3.4, we have $2a + m < \frac{3}{2}$. Then $\lambda(2a+m) - 1 < 1$, because $\lambda < \frac{6}{5}$ by (3.2). We also have $\lambda(2a+m) - 1 > 0$ by Lemma 2.3.

Lemma 3.5. *The point Q is contained in the curve \tilde{C} .*

Proof. If $Q \notin \tilde{C}$, then $\frac{1}{2} \geq \frac{1-a}{2} \geq m = F \cdot \tilde{\Delta} > \frac{1}{\lambda} > \frac{5}{6}$ by Lemma 2.4, Corollary 3.4 and (3.2). \square

By Lemma 3.3 and (3.2), we have $\lambda a < 1$. Observe that $\tilde{C} \cdot F = 2$ and the curve \tilde{C} is smooth. Thus, applying Lemma 2.4, we get

$$1 < \tilde{C} \cdot \left(\lambda \tilde{\Delta} + (\lambda(2a + m) - 1)F \right) = \lambda(C \cdot \Delta - 2m) + 2\lambda(2a + m) - 2 = \lambda(1 - \epsilon + 3a) - 2.$$

Now using $\epsilon > 0$ and $a < \frac{2}{3}$, we obtain $3 < \lambda(1 + 3a - \epsilon) < \lambda(3 - \epsilon) < 3$ by (3.2).

The obtained contradiction completes the proof of Theorem 3.1.

4. DEL PEZZO SURFACES OF DEGREE TWO

Let S be a smooth del Pezzo surface such that $K_S^2 = 2$ and L be an ample \mathbb{Q} -divisor on the surface S . In this section we prove the following result:

Theorem 4.1. *If $-K_S - \frac{2}{3} \frac{-K_S \cdot L}{L^2} L$ is nef, then $\alpha(S, L) > \frac{2}{3} \frac{-K_S \cdot L}{L^2}$.*

As in the proof of Theorem 3.1, we may assume that $\frac{-K_S \cdot L}{L^2} = \frac{3}{2}$. Then

$$(4.1) \quad \left(L - \frac{2}{3}(-K_S) \right) \cdot L = 0.$$

Suppose that $-K_S - L$ is nef. To prove Theorem 4.1, we have to show that $\alpha(S, L) > 1$.

Let $R = -K_S - L$ and $\epsilon = -K_S \cdot R$. Then $\epsilon > 0$, because $R \not\sim_{\mathbb{Q}} 0$ by (4.1). Let

$$(4.2) \quad \gamma = \begin{cases} \frac{12}{11} & \text{if } \epsilon \geq 1, \\ \frac{12}{12 - \epsilon} & \text{if } \epsilon < 1. \end{cases}$$

Then $\gamma > 1$. We will prove that $\alpha(S, L) \geq \gamma$ by *reductio ad absurdum*.

Suppose that $\alpha(S, L) < \gamma$. Then there exist a positive rational number $\lambda < \gamma$ and an effective \mathbb{Q} -divisor D on S such that $D \sim_{\mathbb{Q}} L$, and $(S, \lambda D)$ is not log canonical at some point $P \in S$.

Lemma 4.2. *One has $L \not\sim_{\mathbb{Q}} -\frac{2}{3}K_S$.*

Proof. By [3, Theorem 1.7], $\alpha(S, -K_S) \geq \frac{3}{4}$. Thus, if $L \sim_{\mathbb{Q}} -\frac{2}{3}K_S$, then $(S, \frac{9}{8}D)$ is log canonical, which is impossible, because $\lambda < \gamma \leq \frac{12}{11} < \frac{9}{8}$ by (4.2). \square

Lemma 4.3. *Suppose that $D = \sum_{i=1}^k a_i C_i + \Delta$, where a_1, \dots, a_k are non-negative rational numbers, C_1, \dots, C_k are irreducible curves on the surface S such that*

$$\sum_{i=1}^k C_i \sim_{\mathbb{Q}} -K_S,$$

and Δ is an effective \mathbb{Q} -divisor on the surface S whose support does not contain any curve among the curves C_1, \dots, C_k . Then $a_r < \frac{2}{3}$ for some r .

Proof. Suppose that $a_1 \geq \frac{2}{3}, \dots, a_k \geq \frac{2}{3}$. Then it follows from (4.1) that

$$0 = L \cdot \left(L - \frac{2}{3}(-K_S) \right) = \sum_{i=1}^k \left(a_i - \frac{2}{3} \right) L \cdot C_i + L \cdot \Delta \geq \sum_{i=1}^k \left(a_i - \frac{2}{3} \right) L \cdot C_i \geq 0,$$

which implies that $a_1 = \dots = a_k = \frac{2}{3}$ and $L \cdot \Delta = 0$. Then $\Delta = 0$, since the divisor L is ample and $L \sim_{\mathbb{Q}} -\frac{2}{3}K_S$, which contradicts Lemma 4.2. \square

Let $f: \tilde{S} \rightarrow S$ be a blow-up of the point P , let F be its exceptional curve, and let \tilde{D} be the proper transform on \tilde{S} of the divisor D . Then $(\tilde{S}, \lambda\tilde{D} + (\lambda\text{mult}_P(D) - 1)F)$ is not log canonical at some point $Q \in F$. Note that $\lambda\text{mult}_P(D) > 1$ by Lemma 2.3.

Lemma 4.4. *One has $\text{mult}_P(D) \leq 2 - \epsilon$ and $\lambda\text{mult}_P(D) < 2$.*

Proof. Let C be a general curve in $|-K_S|$ that passes through P . Then $\text{mult}_P(D) \leq D \cdot C = 2 - \epsilon$, which implies that $\lambda\text{mult}_P(D) < 2$ by (4.2). \square

Let $g: \hat{S} \rightarrow \tilde{S}$ be the blow-up at the point Q , and let G be its exceptional curve. Denote by \hat{F} and \hat{D} the proper transforms on \hat{S} of the curve F and the divisor \tilde{D} , respectively. Then the log pair $(\hat{S}, \lambda\hat{D} + (\lambda\text{mult}_P(D) - 1)\hat{F} + (\lambda(\text{mult}_P(D) + \text{mult}_Q(\tilde{D})) - 2)G)$ is not log canonical at some point $O \in G$. Note that it follows from Lemma 2.3 that $\lambda\text{mult}_P(D) + \lambda\text{mult}_Q(\tilde{D}) > 2$.

The linear system $|-K_S|$ gives a double cover $\pi: S \rightarrow \mathbb{P}^2$ branched over a smooth quartic curve $C_4 \subset \mathbb{P}^2$. Thus, if $\pi(P) \in C_4$, then $|-K_S|$ contains a unique curve that is singular at P .

Lemma 4.5. *Suppose that $\pi(P)$ is contained in the curve C_4 . Let T be the unique curve in the linear system $|-K_S|$ that is singular at the point P . Then T is reducible.*

Proof. Suppose that T is irreducible. Write $D = aT + \Delta$, where a is a non-negative rational number, and Δ is an effective \mathbb{Q} -divisor whose support does not contain T . By Lemma 4.3, we have $a < \frac{2}{3}$. Let $m = \text{mult}_P(\Delta)$. Then

$$(4.3) \quad 2m \leq T \cdot \Delta = 2 - \epsilon - 2a \leq 2 - \epsilon.$$

Denote by \tilde{T} and $\tilde{\Delta}$ the proper transforms on \tilde{S} of the curve T and the divisor Δ , respectively. If $Q \notin \tilde{T}$, then Lemma 2.4 gives $\frac{1}{\lambda} < F \cdot \tilde{\Delta} = m \leq 1 - \frac{\epsilon}{2}$, which contradicts (4.2). Then $Q \in \tilde{T}$.

Since T is irreducible, it has either a nodal point at P or a cuspidal point. In particular, the curve \tilde{T} is smooth. If T has a nodal point at P , then Lemma 2.4 gives

$$2 - \epsilon - 2a - 2m = \tilde{T} \cdot \tilde{\Delta} \geq \text{mult}_Q(\tilde{T} \cdot \tilde{\Delta}) > \frac{2}{\lambda} - 2a - m,$$

so that $2 - \epsilon \geq 2 - \epsilon - m > \frac{2}{\lambda}$, which contradicts (4.2). Thus, T has a cuspidal singularity at P .

Denote by \hat{T} and $\hat{\Delta}$ the proper transforms on \hat{S} of the curve \tilde{T} and the divisor $\tilde{\Delta}$, respectively. Let $\tilde{m} = \text{mult}_Q(\tilde{\Delta})$. Then $(\hat{S}, \lambda a \hat{T} + \lambda \hat{\Delta} + (\lambda(2a + m) - 1)\hat{F} + (\lambda(3a + m + \tilde{m}) - 2)G)$ is not log canonical at O . Using Lemma 4.4 and (4.2), we get $\lambda(3a + m + \tilde{m}) \leq (\frac{8}{3} - \epsilon) < 3$, since $a < \frac{2}{3}$.

If $O \neq \hat{T} \cap \hat{F} \cap G$, then $1 - \frac{\epsilon}{2} < \frac{1}{\lambda} < G \cdot \hat{\Delta} = \tilde{m} \leq m$ by Lemma 2.4 and (4.2), which is absurd, because $m \leq 1 - \frac{\epsilon}{2}$ by (4.3). Thus, we have $O = \hat{T} \cap \hat{F} \cap G$. Then

$$1 < \hat{T} \cdot (\lambda \hat{\Delta} + (\lambda(2a + m) - 1)F + (\lambda(3a + m + \tilde{m}) - 2)G) = \lambda(2 + 3a - \epsilon) - 3$$

by Lemma 2.4. Since $a < \frac{2}{3}$, this gives $1 < \lambda(4 - \epsilon) - 3$, which is impossible by (4.2). \square

Lemma 4.6. *The point $\pi(P)$ is not contained in the curve C_4 .*

Proof. Suppose that $\pi(P) \in C_4$. Let T be the curve in $|-K_S|$ that is singular at the point P . By Lemma 4.5, it is reducible, so that $T = L_1 + L_2$, where L_1 and L_2 are irreducible smooth curves such that $L_1^2 = L_2^2 = -1$, $L_1 \cdot L_2 = 2$ and $P \in L_1 \cap L_2$. Let $\epsilon_1 = L_1 \cdot R$ and $\epsilon_2 = L_2 \cdot R$. Then $\epsilon = \epsilon_1 + \epsilon_2$, where $\epsilon_1 \geq 0$ and $\epsilon_2 \geq 0$, since R is nef.

Write $D = a_1 L_1 + a_2 L_2 + \Delta$, where a_1 and a_2 are non-negative rational numbers, and Δ is an effective \mathbb{Q} -divisor whose support contains neither L_1 nor L_2 . Then $a_1 < \frac{2}{3}$ or $a_2 < \frac{2}{3}$ by

Lemma 4.3. Without loss of generality, we may assume that $a_1 < \frac{2}{3}$. It follows from (4.1) that

$$\begin{aligned} 0 &= L \cdot \left(D - \frac{2}{3}(L_1 + L_2) \right) = L \cdot \left(\left(a_1 - \frac{2}{3} \right) L_1 + \left(a_2 - \frac{2}{3} \right) L_2 + \Delta \right) = \\ &= \left(a_1 - \frac{2}{3} \right) (1 - \epsilon_1) + \left(a_2 - \frac{2}{3} \right) (1 - \epsilon_2) + \Delta \cdot L \geq \left(a_1 - \frac{2}{3} \right) (1 - \epsilon_1) + \left(a_2 - \frac{2}{3} \right) (1 - \epsilon_2). \end{aligned}$$

This gives us

$$(4.4) \quad a_1 + a_2 \leq \frac{4}{3} - \frac{2}{3}\epsilon + a_1\epsilon_1 + a_2\epsilon_2.$$

Let $m = \text{mult}_P(\Delta)$. Then we get two inequalities:

$$(4.5) \quad 1 - \epsilon_1 + a_1 - 2a_2 = \Delta \cdot L_1 \geq m,$$

and $1 - \epsilon_2 - 2a_1 + a_2 = \Delta \cdot L_2 \geq m$. Adding them, we get $2 - \epsilon \geq a_1 + a_2 + 2m$, so that $m \leq 1 - \frac{\epsilon}{2}$. Since $a_1 < \frac{2}{3}$, it follows from (4.5) that $2a_2 \leq 1 - \epsilon_1 + a_1 \leq 1 + a_1 < \frac{5}{3}$, so that $a_2 < \frac{5}{6}$.

Denote by \tilde{L}_1, \tilde{L}_2 and $\tilde{\Delta}$ the proper transforms on \tilde{S} of L_1, L_2 and Δ , respectively. Then the log pair $(\tilde{S}, \lambda a_1 \tilde{L}_1 + \lambda a_2 \tilde{L}_2 + \lambda \tilde{\Delta} + (\lambda(a_1 + a_2 + m) - 1)F)$ is not log canonical at the point Q . If $Q \notin \tilde{L}_1 \cup \tilde{L}_2$, then Lemma 2.4 and (4.2) give $m = \tilde{\Delta} \cdot F > \frac{1}{\lambda} > \frac{1}{\gamma} \geq \frac{2-\epsilon}{2}$, which is impossible, since $m \leq 1 - \frac{\epsilon}{2}$. Thus, we have $Q \in \tilde{L}_1 \cup \tilde{L}_2$. If $Q \notin \tilde{L}_1$, then Lemma 2.4 gives

$$\frac{2}{\lambda} - (a_1 + a_2 + m) < \tilde{L}_2 \cdot \tilde{\Delta} = 1 - \epsilon_2 - 2a_1 + a_2 - m,$$

so that $1 - \epsilon_2 + 2a_2 - a_1 > \frac{2}{\lambda}$. Adding this inequality to (4.5), we get $2 - \epsilon > \frac{2}{\lambda} + m \geq \frac{2}{\lambda}$, which contradicts (4.2). Thus, we see that $Q \in \tilde{L}_1$. Similarly, it can be shown that $Q \in \tilde{L}_2$.

We have $Q = F \cap \tilde{L}_1 \cap \tilde{L}_2$. Then L_1 is tangent to L_2 at the point P . Let $\tilde{m} = \text{mult}_Q(\tilde{\Delta})$. Then $\tilde{\Delta} \cdot \tilde{L}_1 \geq \tilde{m}$ and $\tilde{\Delta} \cdot \tilde{L}_2 \geq \tilde{m}$. This gives

$$(4.6) \quad \begin{cases} 1 - \epsilon_1 + a_1 - 2a_2 - m \geq \tilde{m}, \\ 1 - \epsilon_2 - 2a_1 + a_2 - m \geq \tilde{m}. \end{cases}$$

In particular, adding the two inequalities, we obtain

$$(4.7) \quad 2 - \epsilon \geq a_1 + a_2 + 2m + 2\tilde{m}.$$

Denote by $\hat{L}_1, \hat{L}_2, \hat{F}$ and $\hat{\Delta}$ the proper transforms of $\tilde{L}_1, \tilde{L}_2, F$ and $\tilde{\Delta}$ on \hat{S} , respectively. Then $(\hat{S}, \lambda a_1 \hat{L}_1 + \lambda a_2 \hat{L}_2 + \lambda \hat{\Delta} + (\lambda(a_1 + a_2 + m) - 1)\hat{F} + (\lambda(2a_1 + 2a_2 + m + \tilde{m}) - 1)G)$ is not log canonical at the point O . Moreover, we have $2a_1 + 2a_2 + m + \tilde{m} \leq 1 - \frac{\epsilon}{2} + \frac{3}{2}(a_1 + a_2)$ by (4.7). Now using (4.2), (4.4), $a_1 < \frac{2}{3}$ and $a_2 < \frac{5}{6}$, we get

$$\begin{aligned} 2a_1 + 2a_2 + m + \tilde{m} &\leq 1 - \frac{\epsilon}{2} + \frac{3}{2} \left(\frac{4}{3} - \frac{2}{3}\epsilon + a_1\epsilon_1 + a_2\epsilon_2 \right) = 3 - \frac{3}{2}(\epsilon - a_1\epsilon_1 - a_2\epsilon_2) = \\ &= 3 + \frac{3}{2}(a_1 - 1)\epsilon_1 + \frac{3}{2}(a_2 - 1)\epsilon_2 < 3 - \frac{1}{2}\epsilon_1 - \frac{1}{4}\epsilon_2 \leq 3 - \frac{1}{4}\epsilon_1 - \frac{1}{4}\epsilon_2 = 3 - \frac{1}{4}\epsilon = \frac{12 - \epsilon}{4} \leq \frac{3}{\gamma} < \frac{3}{\lambda}. \end{aligned}$$

If $O \notin \hat{L}_1 \cup \hat{L}_2 \cup \hat{F}$, then Lemma 2.4 and (4.2) give $\tilde{m} = \hat{\Delta} \cdot G > \frac{1}{\lambda} > \frac{2-\epsilon}{2}$, which is impossible, because $\tilde{m} \leq m \leq 1 - \frac{\epsilon}{2}$. If $O = \hat{F} \cap G$, then Lemma 2.4 gives $\frac{2}{\lambda} - (a_1 + a_2 + m) < G \cdot \hat{\Delta} = \tilde{m}$, so that $a_1 + a_2 + m + \tilde{m} > \frac{2}{\lambda}$. This together with (4.7) imply $2 - \epsilon \geq a_1 + a_2 + 2m + 2\tilde{m} \geq a_1 + a_2 + m + \tilde{m} > \frac{2}{\lambda}$, which contradicts (4.2). Thus, we see that either $O = \hat{L}_1 \cap G$ or $O = \hat{L}_2 \cap G$.

Suppose that $O = \widehat{L}_1 \cap G$. Then $\widetilde{m} = G \cdot \widehat{\Delta} > \frac{1}{\lambda} - a_1$ by Lemma 2.4. Plugging this into (4.6), we obtain $1 - \epsilon_2 - 2a_1 + a_2 \geq m + \widetilde{m} \geq 2\widetilde{m} > \frac{2}{\lambda} - 2a_1$, so that $1 - \epsilon_2 + a_2 > \frac{2}{\lambda}$. But

$$\frac{3}{\lambda} - (2a_1 + 2a_2 + m + \widetilde{m}) < \widehat{L}_1 \cdot \widehat{\Delta} = 1 - \epsilon_1 + a_1 - 2a_2 - m - \widetilde{m}$$

by Lemma 2.4, so that $1 - \epsilon_1 + 3a_1 > \frac{3}{\lambda}$. Adding these inequalities, we get $3a_1 + a_2 + 2 - \epsilon > \frac{5}{\lambda}$. Now using $a_1 < \frac{2}{3}$, $a_2 < \frac{5}{6}$ and (4.4), we get

$$\begin{aligned} \frac{5}{\lambda} < 3a_1 + a_2 + 2 - \epsilon &< \frac{10}{3} + a_1 + a_2 - \epsilon \leq \frac{10}{3} + \left(\frac{4}{3} - \frac{2}{3}\epsilon + a_1\epsilon_1 + a_2\epsilon_2\right) - \epsilon = \\ &= \frac{14}{3} - \frac{5}{3}\epsilon + a_1\epsilon_1 + a_2\epsilon_2 < \frac{14}{3} - \frac{5}{3}\epsilon + \frac{2}{3}\epsilon_1 + \frac{5}{6}\epsilon_2 = \frac{14}{3} - \epsilon_1 - \frac{5}{6}\epsilon_2 \leq \frac{14}{3} - \frac{5}{6}\epsilon, \end{aligned}$$

which implies that $\gamma > \lambda > \frac{30}{28-5\epsilon}$. This is impossible by (4.2). We conclude that $O \neq \widehat{L}_1 \cap G$.

We have $O = \widehat{L}_2 \cap G$. By (4.7), we also have $4\widetilde{m} \leq 2m + 2\widetilde{m} \leq a_1 + a_2 + 2m + 2\widetilde{m} \leq 2 - \epsilon$, so that $\widetilde{m} \leq \frac{1}{2}$. Then (4.2) gives $\text{mult}_O(\lambda\widehat{\Delta}) \leq \lambda\widetilde{m} \leq \frac{\lambda}{2} < 1$. Thus, applying Theorem 2.5 either we obtain $2(\frac{1}{\lambda} - a_2) < \widehat{\Delta} \cdot G = \widetilde{m}$, or we obtain

$$(4.8) \quad 2\left(\frac{3}{\lambda} - (2(a_1 + a_2) + m + \widetilde{m})\right) < \widehat{\Delta} \cdot \widehat{L}_2 = 1 - \epsilon_2 - 2a_1 + a_2 - m - \widetilde{m}.$$

If $\widetilde{m} > 2(\frac{1}{\lambda} - a_2)$, then (4.2) and (4.6) imply

$$1 + a_1 - 2a_2 \geq 1 - \epsilon_1 + a_1 - 2a_2 \geq m + \widetilde{m} \geq 2\widetilde{m} > 4\left(\frac{1}{\lambda} - a_2\right) = \frac{4}{\lambda} - 4a_2 \geq \frac{11}{3} - 4a_2,$$

which gives $a_1 + 2a_2 > \frac{8}{3}$. But $a_1 + 2a_2 < \frac{7}{3}$, since $a_1 < \frac{2}{3}$ and $a_2 < \frac{5}{6}$. We see that (4.8) holds. It gives us $1 - \epsilon_2 + 2a_1 + 5a_2 + m + \widetilde{m} > \frac{6}{\lambda}$. But $m + \widetilde{m} \leq 1 - \epsilon_1 + a_1 - 2a_2$ by (4.6), so that we obtain $2 - \epsilon + 3(a_1 + a_2) \geq 1 - \epsilon_2 + 2a_1 + 5a_2 + m + \widetilde{m} > \frac{6}{\lambda}$. Now using (4.4), we get

$$\frac{6}{\lambda} < 2 - \epsilon + 3(a_1 + a_2) \leq 6 - 3\epsilon + 3a_1\epsilon_1 + 3a_2\epsilon_2 < 6 - 3\epsilon + 2\epsilon_1 + \frac{5}{2}\epsilon_2 \leq 6 - \frac{\epsilon}{2},$$

because $a_1 < \frac{2}{3}$ and $a_2 < \frac{5}{6}$. Thus, we see that $\lambda > \frac{12}{12-\epsilon}$, which is impossible by (4.2). \square

Let \widetilde{C} be a curve in the pencil $|-K_{\widetilde{S}}|$ that passes through Q . Denote by C its image on the surface S . Then $P \in C$ and $C \in |-K_S|$. Moreover, the curve C is smooth at P , because P is not contained in C_4 by Lemma 4.6. Furthermore, we have

Lemma 4.7. *The curve C is reducible.*

Proof. Suppose that C is irreducible. Let us write $D = aC + \Delta$, where a is a non-negative rational number, and Δ is an effective \mathbb{Q} -divisor on the surface S , whose support does not contain the curve C . Then $a < \frac{2}{3}$ by Lemma 4.3.

Denote by $\widetilde{\Delta}$ the proper transform of the divisor Δ on the surface \widetilde{S} . Let $m = \text{mult}_P(\Delta)$. Then $(\widetilde{S}, \lambda a\widetilde{C} + \lambda\widetilde{\Delta} + (\lambda(a+m) - 1)F)$ is not log canonical at Q . Now, applying Lemma 2.4, we get $\frac{2}{\lambda} - (a+m) < \widetilde{C} \cdot \widetilde{\Delta} = 2 - 2a - \epsilon - m$, so that $\frac{2}{\lambda} < 2 - \epsilon$, which contradicts (4.2). \square

Thus, we have $C = L_1 + L_2$, where L_1 and L_2 are irreducible curves such that $L_1^2 = L_2^2 = -1$ and $L_1 \cdot L_2 = 2$. Since C is smooth at P , we have $P \notin L_1 \cap L_2$. Without loss of generality, we may assume that $P \in L_1$. Let $\epsilon_1 = L_1 \cdot R$ and $\epsilon_2 = L_2 \cdot R$. Then $\epsilon = \epsilon_1 + \epsilon_2$.

Write $D = a_1L_1 + a_2L_2 + \Delta$, where a_1 and a_2 are non-negative rational numbers, and Δ is an effective \mathbb{Q} -divisor, whose support contains neither L_1 nor L_2 . Then $a_1 \leq \frac{1+a_2}{2}$, since

$$(4.9) \quad 1 - \epsilon_2 - 2a_1 + a_2 = L_2 \cdot \Delta \geq 0.$$

If $a_2 < \frac{2}{3}$, then $a_1 \leq \frac{1+a_2}{2} < \frac{5}{6}$. Vice versa, if $a_2 \geq \frac{2}{3}$, then $a_1 < \frac{2}{3}$ by Lemma 4.3. In both cases we have $a_1 < \frac{5}{6}$. Then $\lambda a_1 \leq \frac{10}{11}$ by (4.2).

Denote by \tilde{L}_1 , \tilde{L}_2 and $\tilde{\Delta}$ the proper transforms on \tilde{S} of L_1 , L_2 and Δ , respectively. Then the log pair $(\tilde{S}, \lambda a_1 \tilde{L}_1 + \lambda \tilde{\Delta} + (\lambda(a_1 + m) - 1)F)$ is not log canonical at $Q = \tilde{L}_1 \cap F$, by construction. Since $\lambda a_1 \leq 1$, we can apply Lemma 2.4 to this log pair and the curve \tilde{L}_1 . This gives

$$\frac{2}{\lambda} - (a_1 - m) < \tilde{L}_1 \cdot \tilde{\Delta} = 1 - \epsilon_1 + a_1 - 2a_2 - m,$$

so that $\frac{2}{\lambda} < 1 - \epsilon_1 + 2a_1 - 2a_2$. Using (4.9), we get $\frac{2}{\lambda} < 2 - \epsilon_1 - \epsilon_2 - a_2 \leq 2 - \epsilon_1 - \epsilon_2 = 2 - \epsilon$, which implies that $\lambda > \frac{2}{2-\epsilon}$. This is impossible by (4.2).

The obtained contradiction completes the proof of Theorem 4.1.

5. CUBIC SURFACES

Let S be a smooth cubic surface in \mathbb{P}^3 . Fix six disjoint lines $E_1, E_2, E_3, E_4, E_5, E_6$ in S , and fix a positive rational number $x \leq \frac{1}{10}$. Let $L = -K_S + x \sum_{i=1}^6 E_i$. Then L is ample.

The goal of this section is to prove Theorem 1.5. By Theorem 1.2, to do this it is enough to show that $\alpha(S, L) \geq \frac{2}{3+3x}$. Suppose that this is not true. Then there exist an effective \mathbb{Q} -divisor D on the surface S and a positive rational number $\lambda < \frac{2}{3+3x}$ such that $D \sim_{\mathbb{Q}} L$, and $(S, \lambda D)$ is not log canonical at some point $P \in S$. In this section we seek for a contradiction.

Lemma 5.1. *The log pair $(S, \lambda D)$ is log canonical outside of finitely many points.*

Proof. Suppose that this is not true. Then $D = aC + \Delta$, where C is an irreducible curve, a is a positive rational number such that $a > \frac{1}{\lambda}$, and Δ is an effective \mathbb{Q} -divisor, whose support does not contain the curve C . Denote by d the degree of the curve C in \mathbb{P}^3 . Then

$$3 + 6x = -K_S \cdot D = -K_S \cdot (aC + \Delta) = ad - K_S \cdot \Delta \geq ad > \frac{d}{\lambda} > \frac{3d + 3xd}{2}.$$

which implies that $d \leq 2$. Then C is a line or a conic. Denote by Z a general curve in $|-K_S - C|$.

Suppose that C is a line. Then Z is a smooth conic, $C \cdot Z = 2$, and Z is not contained in the support of Δ . If C is one of the lines $E_1, E_2, E_3, E_4, E_5, E_6$, then $2 + 7x - 2a \geq \Delta \cdot Z \geq 0$, so that $3 + 3x < \frac{2}{\lambda} < 2a \leq 2 + 7x$, which is impossible, because $x \leq \frac{1}{10}$. Then

$$2 + 6x - 2a \geq \left(-K_S + x \sum_{i=1}^6 E_i - aC \right) \cdot Z = (L - aC) \cdot Z = \Delta \cdot Z \geq 0,$$

so that $2a \leq 2 + 6x$, which is impossible, because $2a > \frac{2}{\lambda} > 3 + 3x$ and $x \leq \frac{1}{10}$.

We see that C is a conic. Then Z is a line. Write $\Delta = bZ + \Omega$, where b is a non-negative rational number, and Ω is an effective \mathbb{Q} -divisor, whose support contains no Z . If Z is one of the lines $E_1, E_2, E_3, E_4, E_5, E_6$, then $1 - x - 2a + b = Z \cdot \Omega \geq 0$ and $2 + 7x - 2b \geq C \cdot \Omega \geq 0$, so that $4 + 5x \geq 4a > \frac{4}{\lambda} > 6 + 6x$, which is absurd. Then $(C + Z) \cdot E_i = 1$ for each E_i . But

$$1 + x \sum_{i=1}^6 Z \cdot E_i - 2a + b = Z \cdot \Omega \geq 0$$

and $2 + x \sum_{i=1}^6 C \cdot E_i - 2b = C \cdot \Omega \geq 0$. This gives

$$4 + 12x \geq 4 + x \left(6 + \sum_{i=1}^6 Z \cdot E_i \right) = 4 + x \left(2 \sum_{i=1}^6 Z \cdot E_i + \sum_{i=1}^6 C \cdot E_i \right) \geq 4a > \frac{4}{\lambda} > 6 + 6x,$$

so that $x > \frac{1}{3}$, which is impossible, because $x \leq \frac{1}{10}$. \square

Let $h: S \rightarrow \mathbb{P}^2$ be the contraction of the lines $E_1, E_2, E_3, E_4, E_5, E_6$,

Lemma 5.2. *The point P is contained in $E_1 \cup E_2 \cup E_3 \cup E_4 \cup E_5 \cup E_6$.*

Proof. Suppose that $P \notin E_1 \cup E_2 \cup E_3 \cup E_4 \cup E_5 \cup E_6$. and let ℓ be a line in \mathbb{P}^2 such $h(P) \notin \ell$. Let $\overline{D} = h(D)$, and denote by $\text{LCS}(\mathbb{P}^2, \lambda\overline{D} + \ell)$ the locus in \mathbb{P}^2 consisting of all points where the log pair $(\mathbb{P}^2, \lambda\overline{D} + \ell)$ is not Kawamata log terminal. Then this set contains both $h(P)$ and ℓ , so that it is not connected by Lemma 5.1. But it follows from $\lambda < \frac{2}{3+3x} \leq \frac{2}{3}$ that the divisor

$$-K_{\mathbb{P}^2} + \lambda\overline{D} + \ell \sim_{\mathbb{Q}} (-3 + 3\lambda + 1)\ell$$

is ample. This contradicts Shokurov's connectedness principle [12, Theorem 6.9]. \square

We may assume that $P \in E_1$. Then E_1 is contained in the support of the divisor D , because otherwise we would have $1 - x = E_1 \cdot D \geq \text{mult}_P(D) > 1$ by Lemma 2.3.

Lemma 5.3. *The point P is an Eckardt point.*

Proof. Let T be the plane section of S that is singular at P , let Z_1, Z_2, Z_3, Z_4, Z_5 , and Z_6 be general conics in the pencils $|-K_S - E_1|, |-K_S - E_2|, |-K_S - E_3|, |-K_S - E_4|, |-K_S - E_5|$, and $|-K_S - E_6|$, respectively. Let

$$\Upsilon = \left(\frac{1-\lambda}{6} - \lambda x \right) \sum_{i=1}^6 E_i + \frac{1-\lambda}{6} (Z_1 + Z_2 + Z_3 + Z_4 + Z_5 + Z_6).$$

Then Υ is an effective \mathbb{Q} -divisor such that $\Upsilon + \lambda D \sim_{\mathbb{Q}} -K_S$, and the pair $(S, \lambda D + \Upsilon)$ is not log canonical at P . Then P is not an ordinary double point of the curve T by [5, Corollary 1.24]. Thus, either P is an Eckardt point or T has a tacknodal singularity at the point P .

Suppose that P is not an Eckardt point. Then $T = E_1 + C$, where C is smooth conic such that E_1 is tangent to C at the point P . We have

$$\lambda D \sim_{\mathbb{Q}} (\lambda + \lambda x)E_1 + \lambda C + \lambda x(E_2 + E_3 + E_4 + E_5 + E_6),$$

and $(S, (\lambda + \lambda x)E_1 + \lambda C + \lambda x(E_2 + E_3 + E_4 + E_5 + E_6))$ is log canonical. Therefore, by Remark 2.2, we may assume that the support of the divisor D does not contain either C or E_2 (or both).

Write $D = aE_1 + bC + \Delta$, where a and b are non-negative rational numbers, and Δ is an effective \mathbb{Q} -divisor, whose support does not contain the curves E_1 and C . We know that $a > 0$. If $b > 0$, then $1 - x - b \geq E_2 \cdot \Delta \geq 0$, so that $b \leq 1 - x$. Thus, we always have $b \leq 1 - x$.

Let $m = \text{mult}_P(\Delta)$. Then $1 - x + a - 2b = E_1 \cdot \Delta \geq m$ and $2 + 7x - 2a = C \cdot \Delta \geq m$, which gives $a \leq 1 + \frac{7}{2}x$ and

$$m + a + b \leq \frac{5 + 13x - 3a - 2b}{3} + a + b = \frac{5 + 13x}{3} + \frac{b}{3} \leq \frac{5 + 13x}{3} + \frac{1 - x}{3} = 2 + 4x < \frac{2}{\lambda}$$

and $m \leq \frac{2-2x+2a-4b}{3} + \frac{2+7x-2a}{3} = \frac{4+5x-4b}{3} \leq \frac{4+5x}{3} < \frac{1}{\lambda}$.

Let $f: \tilde{S} \rightarrow S$ be the blow up at P , and let F be its exceptional curve. Denote by \tilde{E}_1, \tilde{C} and $\tilde{\Delta}$ the proper transforms on \tilde{S} of the curve E_1 , the curve C and the divisor Δ , respectively. Then $(\tilde{S}, \lambda a \tilde{E}_1 + \lambda b \tilde{C} + \lambda \tilde{\Delta} + (\lambda(a + b + m) - 1)F)$ is not log canonical at some point $Q \in F$.

If $Q \neq \tilde{E}_1 \cap \tilde{C}$, then Lemma 2.4 gives $m = F \cdot \tilde{\Delta} > \frac{1}{\lambda}$, which is impossible, as $m < \frac{1}{\lambda}$. Let $\tilde{m} = \text{mult}_Q(\tilde{\Delta})$. Then $1 - x + a - 2b - m = \tilde{E}_1 \cdot \tilde{\Delta} \geq \tilde{m}$ and $2 + 7x - 2a - m \geq \tilde{C} \cdot \tilde{\Delta} \geq \tilde{m}$. Since $b \leq 1 - x$, the latter inequality gives

$$m + \tilde{m} + 2a + 2b \leq 2 + 7x + 2b \leq 2 + 7x + 2(1 - x) \leq 4 + 5x < \frac{3}{\lambda}.$$

Let $g: \widehat{S} \rightarrow \widetilde{S}$ be the blow up at Q , and let G be its exceptional curve. Denote by $\widehat{E}_1, \widehat{C}, \widehat{F}$ and $\widehat{\Delta}$ the proper transforms on \widehat{S} of $\widetilde{E}_1, \widetilde{C}, F$ and $\widetilde{\Delta}$, respectively. Then the singularities of the log pair $(\widehat{S}, \lambda a \widehat{E}_1 + \lambda a \widehat{C} + \lambda \widehat{\Delta} + (\lambda(a+b+m) - 1)\widehat{F} + (\lambda(2a+2b+m+\widetilde{m}) - 2)G)$ are not log canonical at some point $O \in G$. Note that $1 \geq \lambda(2a+2b+m+\widetilde{m}) - 2 > 0$.

The curves $\widehat{E}_1, \widehat{C}$, and \widehat{F} are disjoint. If $O \notin \widehat{E}_1 \cup \widehat{C} \cup \widehat{F}$, then $\widetilde{m} = G \cdot \widehat{\Delta} > \frac{1}{\lambda}$ by Lemma 2.4. But $\widetilde{m} \leq m < \frac{1}{\lambda}$, so that $O \in \widehat{E}_1 \cup \widehat{C} \cup \widehat{F}$. If $O = \widehat{E}_1 \cap G$, then Lemma 2.4 implies

$$1 - x + a - 2b - m - \widetilde{m} = \widehat{E}_1 \cdot \widehat{\Delta} > \frac{3}{\lambda} - 2a - 2b - m - \widetilde{m},$$

so that we have $3\frac{2+7x}{2} + 1 - x \geq 3a + 1 - x > \frac{3}{\lambda} > \frac{3(3+3x)}{2}$, which is impossible, because $x \leq \frac{1}{10}$. If $O = \widehat{C} \cap G$, then Lemma 2.4 implies

$$2 + 7x - 2a - m - \widetilde{m} \geq \widehat{C} \cdot \widehat{\Delta} > \frac{3}{\lambda} - 2a - 2b - m - \widetilde{m},$$

so that $4+5x = 2+7x+2(1-x) \geq 2+7x+2b > \frac{3}{\lambda} > \frac{3(3+x)}{2}$, which is impossible, because $x \leq \frac{1}{10}$. Finally, if we have $O = \widehat{F} \cap G$, then Lemma 2.4 implies $m - \widetilde{m} = \widehat{F} \cdot \widehat{\Delta} > \frac{3}{\lambda} - 2a - 2b - m - \widetilde{m}$, so that $2m + 2a + 2b > \frac{3}{\lambda}$, which is impossible because $m + a + b \leq 2 + 4x$ and $x \leq \frac{1}{10}$. \square

We see that S contains two more lines (except the line E_1) that pass through the point P . Denote these two lines by L_1 and L_2 . Recall that E_1 is contained in the support of D . If L_1 is not contained in the support of D , then Lemma 2.3 gives

$$1 + 6x \geq 1 + xL_1 \cdot (E_1 + E_2 + E_3 + E_4 + E_5 + E_6) = L_1 \cdot D \geq \text{mult}_P(D) > \frac{1}{\lambda} > \frac{3+3x}{2}.$$

However, $1 + 6x < \frac{3+3x}{2}$, because $x \leq \frac{1}{10}$. Thus, we see that L_1 is also contained in the support of D . Similarly, we see that L_2 is contained in the support of D as well.

As usual, we write $D = aE_1 + b_1L_1 + b_2L_2 + \Delta$, where a, b_1, b_2 are positive rational numbers, and Δ is an effective \mathbb{Q} -divisor whose support does not contain E_1, L_1 , or L_2 . We have

$$\lambda D \sim_{\mathbb{Q}} (\lambda + \lambda x)E_1 + \lambda L_1 + \lambda L_2 + x(E_2 + E_3 + E_4 + E_5 + E_6),$$

and the singularities of the log pair $(S, (\lambda + \lambda x)E_1 + \lambda L_1 + \lambda L_2 + x(E_2 + E_3 + E_4 + E_5 + E_6))$ are log canonical. Hence, by Remark 2.2, we may assume that $\text{Supp}(\Delta)$ does not contain at least one line among E_2, E_3, E_4, E_5, E_6 .

Lemma 5.4. *Either $b_1 \leq 1 - x$ or $b_2 \leq 1 - x$ (or both).*

Proof. Without loss of generality, we may assume that $E_2 \notin \text{Supp}(D)$. Then

$$1 - x - E_2 \cdot (b_1L_1 + b_2L_2) = E_2 \cdot \Delta \geq 0.$$

But $E_2 \cdot (L_1 + L_2) = 1$, so that either $E_2 \cdot L_1 = 1$ and $E_2 \cdot L_2 = 0$, or $E_2 \cdot L_1 = 0$ and $E_2 \cdot L_2 = 1$. In the former case, we get $b_1 \leq 1 - x$. In the latter case, we get $b_2 \leq 1 - x$. \square

We may assume that $h(L_1)$ is a line in \mathbb{P}^2 , and $h(L_2)$ is a conic. Then $h(L_1)$ is tangent to the conic $h(L_2)$ at the point $h(E_1)$. We may also assume that $h(L_1)$ contains the point $h(E_6)$. Then $L_1 \cdot E_1 = L_1 \cdot E_6 = 1$, $L_1 \cdot E_2 = L_1 \cdot E_3 = L_1 \cdot E_4 = L_1 \cdot E_5 = 0$, $L_2 \cdot E_6 = 0$ and $L_2 \cdot E_1 = L_2 \cdot E_2 = L_2 \cdot E_3 = L_2 \cdot E_4 = L_2 \cdot E_5 = 1$.

Let $m = \text{mult}_P(\Delta)$. Then $E_1 \cdot \Delta \geq m$, $L_1 \cdot \Delta \geq m$ and $L_1 \cdot \Delta \geq m$. This gives

$$(5.1) \quad \begin{cases} 1 - x + a - b_1 - b_2 = m, \\ 1 + 2x - a + b_1 - b_2 = m, \\ 1 + 5x - a - b_1 + b_2 = m. \end{cases}$$

Lemma 5.5. *One has $m+a \leq 1+\frac{7}{2}x$, $m+b_1 \leq 1+2x$, $m+b_2 \leq 1+\frac{x}{2}$ and $a+b_1+b_2+m \leq 3+3x$.*

Proof. The first three inequalities directly follow from (5.1). To prove the fourth inequality, let us recall that $b_1 \leq 1-x$ or $b_2 \leq 1-x$ by Lemma 5.4. Thus, if $b_1 \leq 1-x$, then

$$3+x \geq 3 \geq 1+2x+2b_1 \geq m+a+b_2+b_1$$

by (5.1). Similarly, if $b_2 \leq 1-x$, then (5.1) gives $3+3x \geq 1+5x+2b_2 \geq m+a+b_2+b_1$. \square

Let $f: \tilde{S} \rightarrow S$ be a blow up of the point P . Denote by F be the f -exceptional curve, and denote by \tilde{E}_1 , \tilde{L}_1 and \tilde{L}_2 the proper transforms on \tilde{S} of the lines E_1 , L_1 and L_2 , respectively. Similarly, denote by $\tilde{\Delta}$ the proper transform on \tilde{S} of the \mathbb{Q} -divisor Δ . Then the singularities of the log pair $(\tilde{S}, \lambda a \tilde{E}_1 + \lambda b_1 \tilde{L}_1 + \lambda b_2 \tilde{L}_2 + \lambda \tilde{\Delta} + (\lambda(a+b_1+b_2+m)-1)F)$ are not log canonical at some point $Q \in F$. By Lemma 5.5, we have $m+a+b_1+b_2 \leq 3+3x < \frac{2}{\lambda}$, since $\lambda < \frac{2}{3+3x}$.

If Q is one of the points $F \cap \tilde{L}_1$, $F \cap \tilde{L}_2$ or $F \cap \tilde{E}_1$, then Lemma 2.4 gives

$$m = F \cdot \tilde{\Delta} > \begin{cases} \frac{3}{3+3x} - a & \text{if } Q = F \cap \tilde{E}_1, \\ \frac{3}{3+3x} - b_1 & \text{if } Q = F \cap \tilde{L}_1, \\ \frac{3}{3+3x} - b_2 & \text{if } Q = F \cap \tilde{L}_2, \\ \frac{3}{3+3x} & \text{if } Q \notin \tilde{E}_1 \cup \tilde{L}_1 \cup \tilde{L}_2, \end{cases}$$

because $\lambda < \frac{2}{3+3x}$. But $m+a \leq 1+\frac{7}{2}x$, $m+b_1 \leq 1+2x$ and $m+b_2 \leq 1+\frac{x}{2}$ by Lemma 5.5. This immediately leads to a contradiction, because $0 \leq x \leq \frac{1}{10}$.

The obtained contradiction completes the proof of Theorem 1.5.

6. DEL PEZZO SURFACES OF DEGREE FOUR AND HIGHER

Let S be a smooth del Pezzo surface such that $K_S^2 \geq 4$. Let $d = K_S^2$. Let L be an ample \mathbb{Q} -divisor on the surface S . Let $\nu(L) = \frac{-K_S \cdot L}{L^2}$. The goal of this section is to prove Theorem 1.4.

If S is one of the surfaces \mathbb{P}^2 , $\mathbb{P}^1 \times \mathbb{P}^1$ or \mathbb{F}_1 , then one easily sees that $\alpha(S, L) < \frac{2}{3}\nu(L)$. Thus, to prove Theorem 1.4, we may assume $d \leq 7$. In this case the Mori cone $\overline{\text{NE}}(S)$ is a polyhedral cone that is generated by all (-1) -curves in S . Let

$$\mu_L = \inf \left\{ \lambda \in \mathbb{Q}_{\geq 0} \mid \text{the } \mathbb{Q}\text{-divisor } K_S + \lambda L \in \overline{\text{NE}}(S) \right\}.$$

Replacing L by $\mu_L L$, we may assume that $\mu_L = 1$. Denote by Δ_L the largest extremal face of the Mori cone $\overline{\text{NE}}(S)$ that contains the divisor $K_S + L$, and denote its dimension by r_L .

If $\Delta_L = 0$ and $d = 4$, then $\alpha(S, L) = \frac{2}{3}\nu(L)$ by [3, Theorem 1.7]. Similarly, we see that if $\Delta_L = 0$ and $d \geq 5$, then $\alpha(S, L) \leq \frac{1}{2}\nu(L)$. Thus, we may assume that $\Delta_L \neq 0$, so that $r_L > 0$.

Let $\phi_L: S \rightarrow Z$ be the contraction of the face Δ_L . Then either ϕ_L is a birational morphism, or ϕ_L is a conic bundle with $Z = \mathbb{P}^1$.

Lemma 6.1. *Suppose that $Z = \mathbb{P}^2$. Then $\alpha(S, L) < \frac{2}{3}\nu(L)$.*

Proof. The face Δ_L is generated by $r_L = 9-d$ disjoint (-1) -curves E_1, \dots, E_{r_L} . Then

$$L \sim_{\mathbb{Q}} -K_S + \sum_{i=1}^{r_L} a_i E_i$$

for some positive rational numbers a_1, \dots, a_{r_L} . We may assume that $a_1 \geq \dots \geq a_{r_L}$. Then we must have $a_1 < 1$, because $L \cdot E_1 > 0$. We also have

$$\nu(L) = \frac{d + \sum_{i=1}^{r_L} a_i}{d + 2 \sum_{i=1}^{r_L} a_i - \sum_{i=1}^{r_L} a_i^2}.$$

For every $i \leq r_L$, denote by L_{1i} the proper transform on S of the line in \mathbb{P}^2 that passes through the points $\phi_L(E_1)$ and $\phi_L(E_i)$, e.g., $\phi_L(L_{12})$ is the line that contains $\phi_L(E_1)$ and $\phi_L(E_2)$.

If $d = 7$, then $L \sim_{\mathbb{Q}} 3L_{12} + (2 + a_1)E_1 + (2 + a_2)E_2$, so that

$$\alpha(S, L) \leq \frac{1}{3} < \frac{14 + 2a_1 + 2a_2}{21 + 6a_1 + 6a_2 - 3a_1^2 - 3a_2^2} = \frac{2}{3}\nu(L).$$

Similarly, if $d = 6$, then $L \sim_{\mathbb{Q}} 2L_{12} + L_{13} + (2 + a_1)E_1 + (1 + a_2)E_2 + a_3E_3$, which implies that

$$\alpha(S, L) \leq \frac{1}{2 + a_1} < \frac{12 + 2a_1 + 2a_2 + 2a_3}{18 + 6a_1 + 6a_2 + 6a_3 - 3a_1^2 - 3a_2^2 - 3a_3^2} = \frac{2}{3}\nu(L),$$

because otherwise we would have

$$0 \geq 6 - 2 \sum_{i=1}^3 a_i + 12a_1 + 2a_1 \sum_{i=1}^3 a_i + 3 \sum_{i=1}^3 a_i^2 \geq 6 + 2 \sum_{i=1}^3 a_i + 2a_1 \sum_{i=1}^3 a_i + 3 \sum_{i=1}^3 a_i^2,$$

which is absurd. If $d = 5$, then $L \sim_{\mathbb{Q}} L_{12} + L_{13} + L_{14} + (2 + a_1)E_1 + a_2E_2 + a_3E_3 + a_4E_4$, which implies that

$$\alpha(S, L) \leq \frac{1}{2 + a_1} < \frac{10 + 2a_1 + 2a_2 + 2a_3 + 2a_4}{15 + 6a_1 + 6a_2 + 6a_3 + 6a_4 - 3a_1^2 - 3a_2^2 - 3a_3^2 - 3a_4^2} = \frac{2}{3}\nu(L).$$

Indeed, if this inequality does not hold, then

$$2 \sum_{i=1}^4 a_i \geq 5 + 3 \sum_{i=1}^4 a_i^2 + 10a_1 + 2a_1 \sum_{i=1}^4 a_i > 5a_1 + 3a_1^2 + 10a_1 + 2a_1^2,$$

so that $8a_1 \geq 2 \sum_{i=1}^4 a_i > 15a_1 + 5a_1^2$, which is absurd.

We may assume that $d = 4$. Let Z the proper transform on the surface S of the conic in \mathbb{P}^2 that passes through the points $\phi_L(E_1)$, $\phi_L(E_2)$, $\phi_L(E_3)$, $\phi_L(E_4)$, and $\phi_L(E_5)$. Then

$$L \sim_{\mathbb{Q}} \frac{3 + 2a_1}{2}E_1 + \frac{1}{2}Z + \frac{1}{2}L_{12} + \frac{1}{2}L_{13} + \frac{1}{2}L_{14} + \frac{1}{2}L_{15} + a_2E_2 + a_3E_3 + a_4E_4 + a_5E_5,$$

which implies that $\alpha(S, L) \leq \frac{2}{3+2a_1}$. Let N be the largest number among a_2 , $a_2 + a_3$, $a_2 + a_4$, $a_2 + a_5$, $a_3 + a_4$, $a_3 + a_5$, $a_4 + a_5$, $a_2 + a_3 + a_4$, $a_2 + a_3 + a_5$, $a_2 + a_4 + a_5$, $a_3 + a_4 + a_5$, and $a_2 + a_3 + a_4 + a_5$ that does not exceed 1. We claim that $\alpha(S, L) \leq \frac{2}{3+2a_1+N}$. Indeed, we have $E_2 \sim_{\mathbb{Q}} \frac{1}{2}E_1 - \frac{1}{2}Z - \frac{1}{2}L_{12} + \frac{1}{2}L_{13} + \frac{1}{2}L_{14} + \frac{1}{2}L_{15}$, so that we have

$$L \sim_{\mathbb{Q}} \frac{3 + 2a_1 + a_2}{2}E_1 + \frac{1 - a_2}{2}Z + \frac{1 - a_2}{2}L_{12} + \frac{1 + a_2}{2}L_{13} + \frac{1 + a_2}{2}L_{14} + \frac{1 + a_2}{2}L_{15} + \sum_{i=3}^5 a_i E_i.$$

In particular, if $N = a_2$, then $\alpha(S, L) \leq \frac{2}{3+2a_1+N}$ as claimed. We also have

$$\begin{cases} E_3 \sim_{\mathbb{Q}} \frac{1}{2}E_1 - \frac{1}{2}Z + \frac{1}{2}L_{12} - \frac{1}{2}L_{13} + \frac{1}{2}L_{14} + \frac{1}{2}L_{15}, \\ E_4 \sim_{\mathbb{Q}} \frac{1}{2}E_1 - \frac{1}{2}Z + \frac{1}{2}L_{12} + \frac{1}{2}L_{13} - \frac{1}{2}L_{14} + \frac{1}{2}L_{15}, \\ E_5 \sim_{\mathbb{Q}} \frac{1}{2}E_1 - \frac{1}{2}Z + \frac{1}{2}L_{12} + \frac{1}{2}L_{13} + \frac{1}{2}L_{14} - \frac{1}{2}L_{15}. \end{cases}$$

For any other value of N , Table 1 provides an effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} L$ that shows the inequality $\alpha(S, L) \leq \frac{2}{3+2a_1+N}$ as claimed.

N	Effective \mathbb{Q} -divisor $D \sim_{\mathbb{Q}} L$
$a_2 + a_3$	$\frac{3+2a_1+N}{2} E_1 + \frac{1-N}{2} Z + \frac{1-a_2+a_3}{2} L_{12} + \frac{1+a_2-a_3}{2} L_{13} + \sum_{i=4,5} \left(\frac{1+a_2+a_3}{2} L_{1i} + a_i E_i \right)$
$a_2 + a_4$	$\frac{3+2a_1+N}{2} E_1 + \frac{1-N}{2} Z + \frac{1-a_2+a_4}{2} L_{12} + \frac{1+a_2-a_4}{2} L_{14} + \sum_{i=3,5} \left(\frac{1+a_2+a_4}{2} L_{1i} + a_i E_i \right)$
$a_2 + a_5$	$\frac{3+2a_1+N}{2} E_1 + \frac{1-N}{2} Z + \frac{1-a_2+a_5}{2} L_{12} + \frac{1+a_2-a_5}{2} L_{15} + \sum_{i=3,4} \left(\frac{1+a_2+a_5}{2} L_{1i} + a_i E_i \right)$
$a_3 + a_4$	$\frac{3+2a_1+N}{2} E_1 + \frac{1-N}{2} Z + \frac{1-a_3+a_4}{2} L_{13} + \frac{1+a_3-a_4}{2} L_{14} + \sum_{i=2,5} \left(\frac{1+a_3+a_4}{2} L_{1i} + a_i E_i \right)$
$a_3 + a_5$	$\frac{3+2a_1+N}{2} E_1 + \frac{1-N}{2} Z + \frac{1-a_3+a_5}{2} L_{13} + \frac{1+a_3-a_5}{2} L_{15} + \sum_{i=2,4} \left(\frac{1+a_3+a_5}{2} L_{1i} + a_i E_i \right)$
$a_4 + a_5$	$\frac{3+2a_1+N}{2} E_1 + \frac{1-N}{2} Z + \frac{1-a_4+a_5}{2} L_{14} + \frac{1+a_4-a_5}{2} L_{15} + \sum_{i=2,3} \left(\frac{1+a_4+a_5}{2} L_{1i} + a_i E_i \right)$
$a_2 + a_3 + a_4$	$\frac{3+2a_1+N}{2} E_1 + \frac{1-N}{2} Z + \sum_{i=2,3,4} \left(\frac{1+\sum_{j=2,3,4}(\delta_{ij}a_j)}{2} L_{1i} \right) + \frac{1+N}{2} L_{15} + a_5 E_5$
$a_2 + a_4 + a_5$	$\frac{3+2a_1+N}{2} E_1 + \frac{1-N}{2} Z + \sum_{i=2,4,5} \left(\frac{1+\sum_{j=2,4,5}(\delta_{ij}a_j)}{2} L_{1i} \right) + \frac{1+N}{2} L_{13} + a_3 E_3$
$a_3 + a_4 + a_5$	$\frac{3+2a_1+N}{2} E_1 + \frac{1-N}{2} Z + \sum_{i=3,4,5} \left(\frac{1+\sum_{j=3,4,5}(\delta_{ij}a_j)}{2} L_{1i} \right) + \frac{1+N}{2} L_{12} + a_2 E_2$
$\sum_{i=2}^5 a_i$	$\frac{3+2a_1+N}{2} E_1 + \frac{1-N}{2} Z + \sum_{i=2,3,4,5} \left(\frac{1+\sum_{j=2,3,4,5}(\delta_{ij}a_j)}{2} L_{1i} \right)$

TABLE 1. Effective \mathbb{Q} -divisors, where $\delta_{ij} = 1$ if $i \neq j$ and $\delta_{ij} = -1$ if $i = j$.

Now Proposition A.1 gives $\alpha(S, L) < \frac{2}{3}\nu(L)$. \square

By Lemma 6.1, we may assume that $Z \neq \mathbb{P}^2$ in order to complete the proof of Theorem 1.4. Then the surface S contains $8-d \geq r_L$ disjoint (-1) -curves E_1, \dots, E_{8-d} and a smooth rational curve C such that $C^2 = 0$, the curve C is disjoint from E_1, \dots, E_{8-d} , and

$$L \sim_{\mathbb{Q}} -K_S + \delta C + \sum_{i=1}^{8-d} a_i E_i$$

for some non-negative rational numbers $\delta, a_1, \dots, a_{8-d}$. We may assume that $a_1 \geq \dots \geq a_{8-d}$. Then $a_1 < 1$, since $L \cdot E_1 > 0$ and $C \cdot E_1 = 0$. If $a_i > 0$, then $E_i \in \Delta_L$. Similarly, the morphism ϕ_L is a conic bundle if and only if $\delta > 0$. We have

$$\nu(L) = \frac{d + 2\delta + \sum_{i=1}^{8-d} a_i}{d + 4\delta + 2 \sum_{i=1}^{8-d} a_i - \sum_{i=1}^{8-d} a_i^2}.$$

Let $h: S \rightarrow \bar{S}$ be the contraction of the curves E_1, \dots, E_{8-d} . Then $\bar{S} = \mathbb{F}_1$ or $\bar{S} = \mathbb{P}^1 \times \mathbb{P}^1$. The linear system $|C|$ is a free pencil. It gives a conic bundle $\iota: S \rightarrow \mathbb{P}^1$ such that there exists

a commutative diagram

$$\begin{array}{ccc} S & \xrightarrow{h} & \bar{S} \\ & \searrow \iota & \swarrow v \\ & \mathbb{P}^1 & \end{array}$$

where v is a natural projection. Then the curves E_1, \dots, E_{8-d} lie in the reducible fibers of ι .

Lemma 6.2. *Suppose that $\bar{S} = \mathbb{F}_1$. Then $\alpha(S, L) < \frac{2}{3}\nu(L)$.*

Proof. Let \bar{E}_{9-d} be the (-1) -curve on \bar{S} , and let E_{9-d} be its proper transform on the surface S . Let $f: \bar{S} \rightarrow \mathbb{P}^2$ be the contraction of the curve \bar{E}_{9-d} . Let $\pi = f \circ g$. Denote by L_{1i} the proper transform on S of the line in \mathbb{P}^2 that passes through $\pi(E_1)$ and $\pi(E_i)$ (for every $i \leq 9-d$).

If $d = 7$, then $L \sim_{\mathbb{Q}} (3 + \delta)L_{12} + (2 + \delta + a_1)E_1 + 2E_2$ and

$$\alpha(S, L) \leq \frac{1}{3 + \delta} < \frac{14 + 4\delta + 2a_1}{21 + 12\delta + 6a_1 - 3a_1^2} = \frac{2}{3}\nu(L),$$

since otherwise we would have $0 \geq 21 + 14\delta + 4\delta^2 + 2\delta a_1 + 3a_1^2$.

If $d = 6$, then $L \sim_{\mathbb{Q}} 2L_{12} + (1 + \delta)L_{13} + (2 + \delta + a_1)E_1 + (1 + a_2)E_2$, which gives

$$\alpha(S, L) \leq \frac{1}{2 + \delta + a_1} < \frac{12 + 4\delta + 2a_1 + 2a_2}{18 + 12\delta + 6a_1 + 6a_2 - 3a_1^2 - 3a_2^2} = \frac{2}{3}\nu(L),$$

because otherwise we would have

$$0 \geq 6 - 2(a_1 + a_2) + 3(a_1^2 + a_2^2) + 8\delta + 12a_1 + (\delta + a_1)(4\delta + 2a_1 + 2a_2),$$

which gives a contradiction, since $a_1 \geq a_2$.

If $d = 5$, then $L \sim_{\mathbb{Q}} L_{12} + L_{13} + (1 + \delta)L_{14} + (2 + \delta + a_1)E_1 + a_2E_2 + a_3E_3$, which gives

$$\alpha(S, L) \leq \frac{1}{2 + \delta + a_1} < \frac{10 + 4\delta + 2a_1 + 2a_2 + 2a_3}{15 + 12\delta + 6a_1 + 6a_2 + 6a_3 - 3a_1^2 - 3a_2^2 - 3a_3^2} = \frac{2}{3}\nu(L).$$

Indeed, if this inequality does not hold, then we have

$$0 \geq 5 + 6\delta - 2 \sum_{i=1}^3 a_i + 3 \sum_{i=1}^3 a_i^2 + 10a_1 + 4\delta(\delta + a_1) + 2(\delta + a_1) \sum_{i=1}^3 a_i,$$

which is impossible since $-2 \sum_{i=1}^3 a_i + 10a_1 \geq 4a_1$ as $a_1 \geq a_2 \geq a_3$.

Thus, we may assume that $d = 4$. Denote by Z be the proper transform on the surface S of the unique conic in \mathbb{P}^2 that passes through the points $\pi(E_1), \pi(E_2), \pi(E_3), \pi(E_4), \pi(E_5)$. Then

$$L \sim_{\mathbb{Q}} \delta C + \frac{3 + 2a_1}{2}E_1 + \frac{1}{2}(Z + L_{12} + L_{13} + L_{14} + L_{15}) + a_2E_2 + a_3E_3 + a_4E_4,$$

Moreover, there exists a (-1) -curve E'_1 such that $E'_1 + E_1 \sim C$, so that

$$L \sim_{\mathbb{Q}} \frac{3 + 2a_1 + 2\delta}{2}E_1 + \delta E'_1 + \frac{1}{2}(Z + L_{12} + L_{13} + L_{14} + L_{15}) + a_2E_2 + a_3E_3 + a_4E_4.$$

Arguing as in the proof of Lemma 6.1, we see that

$$\alpha(S, L) \leq \frac{2}{3 + 2a_1 + 2\delta + N},$$

where N is the largest number among $a_2, a_2 + a_3, a_2 + a_4, a_3 + a_4, a_2 + a_3 + a_4$, that does not exceed 1. Then $\alpha(S, L) < \frac{2}{3}\nu(L)$ by Proposition A.1. \square

The final step of the proof of Theorem 1.4 is the following Lemma.

Lemma 6.3. *Suppose that $\overline{S} = \mathbb{P}^1 \times \mathbb{P}^1$. Then $\alpha(S, L) < \frac{2}{3}\nu(L)$.*

Proof. We may assume that the fibers of the natural projection $v: \overline{S} \rightarrow \mathbb{P}^1$ are curves of bi-degree $(1, 0)$. For every $i \leq 8 - d$, we denote by \overline{F}_i (respectively, by \overline{F}'_i) the curve of bi-degree $(1, 0)$ (respectively, bi-degree $(1, 0)$) in $\mathbb{P}^1 \times \mathbb{P}^1$ that passes through the point $h(E_i)$, and we denote by F_i (respectively, by F'_i) its proper transform on S .

If $d = 7$, then $L \sim_{\mathbb{Q}} (3 + a_1 + \delta)E_1 + (2 + \delta)F_1 + 2F'_1$, which gives

$$\alpha(S, L) \leq \frac{1}{3 + a_1 + \delta} < \frac{14 + 4\delta + 2a_1}{21 + 12\delta + 6a_1 - 3a_1^2} = \frac{2}{3}\nu(L).$$

Similarly, if $d = 6$, then $L \sim_{\mathbb{Q}} (\frac{3}{2} + \delta)F_1 + \frac{3}{2}F'_1 + \frac{1}{2}(F_2 + F'_2) + (2 + \delta + a_1)E_1 + a_2E_2$, which gives

$$\alpha(S, L) < \frac{1}{2 + \delta + a_1} \leq \frac{12 + 4\delta + 2a_1 + 2a_2}{18 + 12\delta + 6a_1 + 6a_2 - 3a_1^2 - 3a_2^2} = \frac{2}{3}\nu(L).$$

For $d \leq 5$, denote by Z_{ij} for $2 \leq i < j \leq 4$ the proper transform on S of the unique irreducible curve of bi-degree $(1, 1)$ on \overline{S} that passes through the points $h(E_i)$ and $h(E_j)$. If $d = 5$, then

$$L \sim_{\mathbb{Q}} (1 + \delta)F_1 + F'_1 + Z_{23} + (2 + \delta + a_1)E_1 + a_2E_2 + a_3E_3,$$

so that, arguing as in the proof of Lemma 6.2, we see that

$$\alpha(S, L) \leq \frac{1}{2 + \delta + a_1} < \frac{10 + 4\delta + 2a_1 + 2a_2 + 2a_3}{15 + 12\delta + 6a_1 + 6a_2 + 6a_3 - 3a_1^2 - 3a_2^2 - 3a_3^2} = \frac{2}{3}\nu(L).$$

Thus, we may assume that $d = 4$. Then $-K_S \sim_{\mathbb{Q}} \frac{3}{2}E_1 + \frac{1}{2}(F_1 + F'_1 + Z_{23} + Z_{24} + Z_{34})$. Hence

$$L \sim_{\mathbb{Q}} \frac{3 + 2a_1 + 2\delta}{2}E_1 + \frac{1 + \delta}{2}F_1 + \frac{1}{2}F'_1 + \frac{1}{2}Z_{23} + \frac{1}{2}Z_{24} + \frac{1}{2}Z_{34} + a_2E_2 + a_3E_3 + a_4E_4,$$

which implies, in particular, that $\alpha(S, L) \leq \frac{2}{3 + 2a_1 + 2\delta}$. Moreover, we have

$$\begin{cases} E_2 \sim_{\mathbb{Q}} \frac{1}{2}E_1 + \frac{1}{2}F_1 + \frac{1}{2}F'_1 - \frac{1}{2}Z_{23} - \frac{1}{2}Z_{24} + \frac{1}{2}Z_{34}, \\ E_3 \sim_{\mathbb{Q}} \frac{1}{2}E_1 + \frac{1}{2}F_1 + \frac{1}{2}F'_1 - \frac{1}{2}Z_{23} + \frac{1}{2}Z_{24} - \frac{1}{2}Z_{34}, \\ E_4 \sim_{\mathbb{Q}} \frac{1}{2}E_1 + \frac{1}{2}F_1 + \frac{1}{2}F'_1 + \frac{1}{2}Z_{23} - \frac{1}{2}Z_{24} - \frac{1}{2}Z_{34}. \end{cases}$$

Therefore, we have

$$\begin{aligned} L \sim_{\mathbb{Q}} \frac{3 + 2a_1 + 2\delta + a_2}{2}E_1 + \frac{1 + \delta + a_2}{2}F_1 + \frac{1 + a_2}{2}F'_1 + \\ + \frac{1 - a_2}{2}Z_{23} + \frac{1 - a_2}{2}Z_{24} + \frac{1 + a_2}{2}Z_{34} + a_3E_3 + a_4E_4, \end{aligned}$$

so that $\alpha(S, L) \leq \frac{2}{3 + 2a_1 + 2\delta + a_2}$. Moreover, if $a_2 + a_3 \leq 1$, then $\alpha(S, L) \leq \frac{2}{3 + 2a_1 + 2\delta + a_2 + a_3}$, since

$$\begin{aligned} L \sim_{\mathbb{Q}} \frac{3 + 2a_1 + 2\delta + a_2 + a_3}{2}E_1 + \frac{1 + \delta + a_2 + a_3}{2}F_1 + \frac{1 + a_2 + a_3}{2}F'_1 + \\ + \frac{1 - a_2 - a_3}{2}Z_{23} + \frac{1 - a_2 + a_3}{2}Z_{24} + \frac{1 + a_2 - a_3}{2}Z_{34} + a_4E_4. \end{aligned}$$

Similarly, if $a_2 + a_4 \leq 1$, then $\alpha(S, L) \leq \frac{2}{3+2a_1+2\delta+a_2+a_4}$, since

$$L \sim_{\mathbb{Q}} \frac{3+2a_1+2\delta+a_2+a_4}{2} E_1 + \frac{1+\delta+a_2+a_4}{2} F_1 + \frac{1+a_2+a_4}{2} F'_1 + \\ + \frac{1-a_2+a_4}{2} Z_{23} + \frac{1-a_2-a_4}{2} Z_{24} + \frac{1+a_2-a_4}{2} Z_{34} + a_3 E_3.$$

And if $a_3 + a_4 \leq 1$, then $\alpha(S, L) \leq \frac{2}{3+2a_1+2\delta+a_3+a_4}$, since

$$L \sim_{\mathbb{Q}} \frac{3+2a_1+2\delta+a_3+a_4}{2} E_1 + \frac{1+\delta+a_3+a_4}{2} F_1 + \frac{1+a_3+a_4}{2} F'_1 + \\ + \frac{1-a_3+a_4}{2} Z_{23} + \frac{1+a_3-a_4}{2} Z_{24} + \frac{1-a_3-a_4}{2} Z_{34} + a_2 E_2.$$

Finally, if $a_2 + a_3 \leq 1 + a_4$, then $\alpha(S, L) \leq \frac{2}{3+2a_1+2\delta+a_2+a_3+a_4}$, since

$$L \sim_{\mathbb{Q}} \frac{3+2a_1+2\delta+a_2+a_3+a_4}{2} E_1 + \frac{1+\delta+a_2+a_3+a_4}{2} F_1 + \frac{1+a_2+a_3+a_4}{2} F'_1 + \\ + \frac{1-a_2-a_3+a_4}{2} Z_{23} + \frac{1-a_2+a_3-a_4}{2} Z_{24} + \frac{1+a_2-a_3-a_4}{2} Z_{34}.$$

Thus, we proved that

$$\alpha(S, L) \leq \begin{cases} \frac{2}{3+2a_1+2\delta+a_2+a_3+a_4} & \text{if } a_2 + a_3 \leq 1 + a_4, \\ \frac{2}{3+2a_1+2\delta+a_2+a_4} & \text{if } a_2 + a_4 \leq 1 \text{ and } a_2 + a_3 > 1 + a_4, \\ \frac{2}{3+2a_1+2\delta+a_3+a_4} & \text{if } a_3 + a_4 \leq 1, a_2 + a_4 > 1 \text{ and } a_2 + a_3 > 1 + a_4, \\ \frac{2}{3+2a_1+2\delta+a_2} & \text{if } a_3 + a_4 > 1, a_2 + a_4 > 1 \text{ and } a_2 + a_3 > 1 + a_4. \end{cases}$$

By Proposition A.1, $\alpha(S, L) < \frac{8+4\delta+2a_1+2a_2+2a_3+2a_4}{12+12\delta+6a_1+6a_2+6a_3+6a_4-3a_1^2-3a_2^2-3a_3^2-3a_4^2} = \frac{2}{3}\nu(L)$. \square

APPENDIX A. INEQUALITIES

Let a_1, a_2, a_3, a_4, a_5 be non-negative rational numbers such that $1 \geq a_1 \geq a_2 \geq a_3 \geq a_4 \geq a_5$, let δ be a non-negative rational number, and let N be the largest number among $a_2, a_2 + a_3, a_2 + a_4, a_2 + a_5, a_3 + a_4, a_3 + a_5, a_4 + a_5, a_2 + a_3 + a_4, a_2 + a_3 + a_5, a_2 + a_4 + a_5, a_3 + a_4 + a_5, a_2 + a_3 + a_4 + a_5$ that does not exceed 1. Let

$$\alpha = \begin{cases} \frac{2}{3+2a_1+2\delta+a_2+a_3+a_4} & \text{if } a_2 + a_3 \leq 1 + a_4, \\ \frac{2}{3+2a_1+2\delta+a_2+a_4} & \text{if } a_2 + a_4 \leq 1 \text{ and } a_2 + a_3 > 1 + a_4, \\ \frac{2}{3+2a_1+2\delta+a_3+a_4} & \text{if } a_3 + a_4 \leq 1, a_2 + a_4 > 1 \text{ and } a_2 + a_3 > 1 + a_4, \\ \frac{2}{3+2a_1+2\delta+a_2} & \text{if } a_3 + a_4 > 1, a_2 + a_4 > 1 \text{ and } a_2 + a_3 > 1 + a_4. \end{cases}$$

Proposition A.1. *One has*

$$\frac{2}{3+2a_1+2\delta+N} \leq \frac{2}{3} \cdot \frac{4+2\delta+a_1+a_2+a_3+a_4+a_5}{4+4\delta+2(a_1+a_2+a_3+a_4+a_5)-a_1^2-a_2^2-a_3^2-a_4^2-a_5^2}$$

and

$$\alpha \leq \frac{8 + 4\delta + 2a_1 + 2a_2 + 2a_3 + 2a_4}{12 + 12\delta + 6a_1 + 6a_2 + 6a_3 + 6a_4 - 3a_1^2 - 3a_2^2 - 3a_3^2 - 3a_4^2}.$$

Moreover, both inequalities are strict unless $a_1 = \delta = 0$.

Proof. Consider the first inequality. If $a_1 = \delta = 0$, then $N = 0$ and both sides of the inequality equal $\frac{2}{3}$. Suppose that either $a_1 > 0$ or $\delta > 0$ (or both). We have to prove that

$$\frac{2}{3 + 2a_1 + 2\delta + N} < \frac{2}{3} \times \frac{4 + 2\delta + a_1 + a_2 + a_3 + a_4 + a_5}{4 + 4\delta + 2(a_1 + a_2 + a_3 + a_4 + a_5) - a_1^2 - a_2^2 - a_3^2 - a_4^2 - a_5^2}$$

Suppose that this inequality does not hold. Then

$$3 \sum_{i=1}^5 a_i - 3 \sum_{i=1}^5 a_i^2 - 2\delta \geq (2a_1 + N + 2\delta) \sum_{i=1}^5 a_i + 8a_1 + 4N + 4a_1\delta + 2N\delta + 4\delta^2.$$

Since either $a_1 > 0$ or $\delta > 0$, this inequality implies that

$$(A.1) \quad 3 \sum_{i=1}^5 a_i > (2a_1 + N) \sum_{i=1}^5 a_i + 8a_1 + 4N.$$

If $N = a_2 + a_3 + a_4 + a_5$, then (A.1) gives

$$3 \sum_{i=1}^5 a_i > (2a_1 + N) \sum_{i=1}^5 a_i + 8a_1 + 4N \geq 8a_1 + 4N > 4 \sum_{i=1}^5 a_i,$$

which is absurd. Thus, $N \neq a_2 + a_3 + a_4 + a_5$, so that $a_2 + a_3 + a_4 + a_5 > 1$. Then (A.1) gives

$$3 \sum_{i=1}^5 a_i > (2a_1 + N) \sum_{i=1}^5 a_i + 8a_1 + 4N > (2a_1 + N)(a_1 + 1) + 8a_1 + 4N,$$

which implies, in particular, that

$$(A.2) \quad 3(a_2 + a_3 + a_4 + a_5) > 7a_1 + 5N,$$

which implies that N is none of the numbers $a_2 + a_3 + a_4$, $a_2 + a_3 + a_5$, $a_2 + a_4 + a_5$, $a_3 + a_4 + a_5$. If N is one of the numbers $a_2 + a_3$, $a_2 + a_4$, $a_2 + a_5$, $a_3 + a_4$, $a_3 + a_5$, $a_4 + a_5$, then (A.2) gives

$$3(a_2 + a_3 + a_4 + a_5) > 7a_1 + 5N \geq 7a_1 + 5(a_4 + a_5),$$

so that $3a_2 + 3a_3 > 7a_1$, which is absurd. We see that $N = a_2$. Then (A.2) gives

$$3(a_3 + a_4 + a_5) > 7a_1 + 2a_2,$$

which is impossible, because $a_1 \geq a_2 \geq a_3 \geq a_4 \geq a_5 \geq 0$.

Consider the second inequality. If $a_1 = \delta = 0$, then both sides of the inequality equal $\frac{2}{3}$. Suppose that $a_1 > 0$ or $\delta > 0$. Take $N \geq 0$ such that $\alpha = \frac{2}{3+2a_1+2\delta+N}$. We have to prove that

$$\alpha < \frac{8 + 4\delta + 2a_1 + 2a_2 + 2a_3 + 2a_4}{12 + 12\delta + 6a_1 + 6a_2 + 6a_3 + 6a_4 - 3a_1^2 - 3a_2^2 - 3a_3^2 - 3a_4^2}.$$

Suppose the latter is false. Then

$$3 \sum_{i=1}^4 a_i - 3 \sum_{i=1}^4 a_i^2 - 2\delta \geq (2a_1 + N + 2\delta) \sum_{i=1}^4 a_i + 8a_1 + 4N + 4a_1\delta + 2N\delta + 4\delta^2.$$

Since either $a_1 > 0$ or $\delta > 0$, this inequality implies that

$$3 \sum_{i=1}^5 a_i > (2a_1 + N) \sum_{i=1}^5 a_i + 8a_1 + 4N.$$

If $N = a_2 + a_3 + a_4$ or $N = a_2 + a_4$, then we get a contradiction

$$3 \sum_{i=1}^4 a_i > (2a_1 + N) \sum_{i=1}^4 a_i + 8a_1 + 4N \geq 8a_1 + 4N > 4 \sum_{i=1}^4 a_i,$$

because $a_1 \geq a_3 \geq a_4$. Thus $N = a_2$, $a_3 + a_4 > 1$, $a_2 + a_4 > 1$ and $a_2 + a_3 > 1 + a_4$. Let $a_5 = 0$. Now the result follows from the first inequality, since a_2 is the largest number not exceeding 1 among those values in the hypothesis. \square

REFERENCES

- [1] C. Arezzo, F. Pacard, *Blowing up Kähler manifolds with constant scalar curvature. II*, Ann. of Math. **170** (2009), no. 2, 685–738.
- [2] R. Berman, T. Darvas, C. Lu, *Regularity of weak minimizers of the K-energy and applications to properness and K-stability*, preprint, arXiv:1602.03114 (2016).
- [3] I. Cheltsov, *Log canonical thresholds of del Pezzo surfaces*, Geom. Funct. Anal., **18** (2008), 1118–1144.
- [4] I. Cheltsov, *Del Pezzo surfaces and local inequalities*, Proceedings of the Trento conference “Groups of Automorphisms in Birational and Affine Geometry”, October 2012, Springer (2014), 83–101.
- [5] I. Cheltsov, J. Park, J. Won, *Affine cones over smooth cubic surfaces*, J. European Math. Soc. **18** (2016), 1537–1564.
- [6] X.-X. Chen, S. Donaldson, S. Sun, *Kähler-Einstein metrics on Fano manifolds. I, II, III*, J. Amer. Math. Soc. **28** (2015), no. 1, 183–197, 199–234, 235–278.
- [7] G. Codogni, J. Stoppa, *Torus equivariant K-stability*, preprint, arXiv 1602.03451 (2016).
- [8] R. Dervan, *Alpha invariants and K-stability for general polarisations of Fano varieties*, Int. Math. Res. Notices **16** (2015), 7162–7189.
- [9] S. Donaldson, *Constant scalar curvature metrics on toric surfaces*, Geom. Funct. Anal. **19** (2009), no. 1, 83–136.
- [10] K. Hong, J. Won, *Alpha invariant for general polarizations of del Pezzo surfaces of degree 1*, preprint, arXiv:1606.01418 (2016).
- [11] J. Ross, R. Thomas, *An obstruction to the existence of constant scalar curvature Kähler metrics*, J. Differential Geom. **72** (2006), 429–466.
- [12] V. Shokurov, *Three-dimensional log perestroikas*, Russian Acad. Sci. Izv. Math. **40** (1993), 95–202.
- [13] G. Tian, *On Calabi’s conjecture for complex surfaces with positive first Chern class*, Invent. Math. **101** (1990), no. 1, 101–172.