

The Signature mod 8

by Frank, David

in Commentarii mathematici Helvetici

volume 48; pp. 520 - 524



Göttingen State and University Library

Terms and Conditions

The Göttingen State and University Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Göttingen State- and University Library.

Each copy of any part of this document must contain these Terms and Conditions. With the usage of the library's online-systems to access or download a digitized document you accept these Terms and Conditions.

Reproductions of materials on the web site may not be made for or donated to other repositories, nor may they be further reproduced without written permission from the Göttingen State- and University Library.

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

Contact:

Niedersächsische Staats- und Universitätsbibliothek Göttingen

Digitalisierungszentrum

37070 Göttingen

Germany

E-Mail: gdz@www.sub.uni-goettingen.de

Purchase a CD-ROM

The Göttingen State and University Library offers CD-ROMs containing whole volumes / monographs in PDF for Adobe Acrobat. The PDF-version contains the table of contents as bookmarks, which allows easy navigation in the document. For availability and pricing, please contact:

Niedersächsische Staats- und Universitätsbibliothek Göttingen

Digitalisierungszentrum


37070 Göttingen

Germany

E-Mail: gdz@www.sub.uni-goettingen.de



Göttingen State and University Library



The Signature mod 8

by DAVID FRANK¹⁾

Let Γ_i be the group of exotic i -dimensional spheres, and let bP_{i+1} be the subgroup of those exotic spheres which bound π -manifolds. There is an exact sequence

$$0 \rightarrow bP_{4k} \rightarrow \Gamma_{4k-1} \rightarrow \pi'_{4k-1} \rightarrow 0,$$

where π'_i is the cokernel of the J -homomorphism $J: \pi_i(SO) \rightarrow \pi_i = \pi_{i+t}(S^t)$, t large. In studying the group bP_{4k} , it was important for Milnor to know

PROPOSITION A. *Let M^{4k} be a smooth, compact, oriented manifold with boundary an exotic sphere. If M is a π -manifold, then the signature of M is divisible by 8.*

In showing that the above exact sequence was split, it was important for Brumfiel [2] and the author [3] to know the stronger

PROPOSITION B. *Let M^{4k} be a smooth, compact, oriented manifold with boundary an exotic sphere. If M is a spin manifold, and if all decomposable Pontrjagin numbers of M are zero, then the signature of M is divisible by 8.*

In this paper we will prove a very general theorem which includes Propositions A and B. Let BSG be the classifying space for stable oriented spherical fibrations. If we kill the second homotopy group of BSG, we obtain a space BSpinG, the classifying space for stable spherical fibrations with a spin structure. Let v_{2k} denote the universal Wu class in either BSG or BSpinG. We first show.

LEMMA 1. *There is a class x_{2k} in $H^{2k}(\text{BSpinG}; Z_4)$ whose mod 2 reduction is v_{2k} in $H^{2k}(\text{BSpinG}; Z_2)$.*

The corresponding statement is of course false in BSG.

We now use the Pontrjagin square cohomology operations. There is a family of such operations; we are interested only in the operations

$$P: H^{2k}(-; Z_4) \rightarrow H^{4k}(-; Z_8).$$

In particular, consider the universal characteristic class $P(x_{2k})$ in $H^{4k}(\text{BSpinG}; Z_8)$. If M is a spin Poincaré complex of dimension $4k$ with fundamental homology class $[M]$, we may consider the characteristic number $P(x_{2k})[M]$, which is an integer modulo 8. Let $\sigma(M)$ denote the signature of M .

¹⁾ Supported by NSF grant No. P 029431000.

THEOREM 2. $P(x_{2k})[M] = \sigma(M)$ modulo 8 for any spin Poincaré complex M^{4k} . For oriented Poincaré complexes, we can show

THEOREM 3. There is a characteristic class y_{4k} in $H^{4k}(\text{BSG}; \mathbb{Z}_8)$ such that $y_{4k}[M] = \sigma(M)$ modulo 8 for any oriented Poincaré complex.

Theorem 3 is the best possible ‘Hirzebruch Signature Theorem’ for Poincaré complexes; the integer 8 cannot be replaced by a larger integer.

The classes y_{4k} are related to the k -invariants of the fibration $F/\text{Top} \rightarrow \text{BStTop} \rightarrow \text{BSG}$.

1. The Spin Case

We begin by considering the Wu class v_{2k} in $H^{2k}(\text{BSpinG}; \mathbb{Z}_2)$. We wish to show v_{2k} is the mod 2 reduction of a \mathbb{Z}_4 cohomology class. Equivalently, we can show $Sq^1(v_{2k}) = 0$, where Sq^i is the i -th Steenrod Square. (In fact, $v_{2k} = 0$ for k odd, but this is irrelevant to what follows.)

We recall one definition of the Wu classes v_i . Let γ be the universal spherical fibration on $\text{BSG}(m)$, m large, let $\text{MSG}(m)$ be the Thom space of γ , and let U be the Thom class. Then define

$$v_i = T^{-1}(\chi(Sq^i U)),$$

where T is the Thom isomorphism ($T: H^*(\text{BSG}(m)) \rightarrow H^*(\text{MSG}(m))$) and χ is the anti-automorphism of the Steenrod Algebra. If M^n is a Poincaré complex, let $h: M \rightarrow \text{BSG}(m)$ be the classifying map of the stable Spivak normal fibration of M . Define $v_i(M)$ as $h^*(v_i)$. It follows from [6, Ch. III] that

$$(v_i(M) \cup x)[M] = Sq^i(x)[M]$$

for all x in $H^{n-i}(M; \mathbb{Z}_2)$. Thus this is an acceptable definition of the Wu classes.

Now note that $Sq^1(v_{2k} \cdot U) = Sq^1(v_{2k}) \cdot U$, since $Sq^1(U) = 0$. Thus $Sq^1(v_{2k}) = 0$ if $Sq^1(v_{2k} \cdot U) = 0$. But

$$\begin{aligned} Sq^1(v_{2k} \cdot U) &= Sq^1(\chi Sq^{2k}) U \\ &= (\chi Sq^1)(\chi Sq^{2k}) U \\ &= \chi(Sq^{2k} Sq^1) U. \end{aligned}$$

Now

$$Sq^{2k} Sq^1 = Sq^2 Sq^{2k-1} + a Sq^1 Sq^{2k}, \quad a \in \mathbb{Z}_2,$$

by an Adem relation, so

$$\chi(Sq^{2k} Sq^1) = \chi(Sq^{2k-1}) \chi(Sq^2) + a \chi(Sq^{2k}) \chi(Sq^1).$$

Since the last expression is zero on the Thom class U of $BSpinG(m)$, we have shown that $Sq^1(v_{2k})=0$ in $BSpinG$. Thus, as claimed in Lemma 1, there is a class x_{2k} in $H^{2k}(BSpinG; Z_4)$ whose mod2 reduction is v_{2k} . If M^{4k} is a spin Poincaré complex, we wish to relate the class $x_{2k}(M)$ to the signature of M . In fact, we will prove Theorem 2 by showing

PROPOSITION 4. *Let M^{4k} be an oriented Poincaré complex and let x be any class in $H^{2k}(M; Z_4)$ whose mod2 reduction is $v_{2k}(M)$. Then $P(x)[M]=\sigma(M) \bmod 8$.*

To prove Proposition 4, we first show

LEMMA 5. *Let M^{4k} be an oriented Poincaré complex and let a and b be classes in $H^{2k}(M; Z_4)$ whose mod2 reduction is $v_{2k}(M)$. Then $P(a)=P(b)$.*

Proof. If t is a cohomology class with Z_{2^n} coefficients, we denote by t' the corresponding class with $Z_{2^{n+1}}$ coefficients (determined by the inclusion homomorphism from Z_{2^n} to $Z_{2^{n+1}}$). Now if $a \bmod 2 = b \bmod 2 = v_{2k}(M)$, then $a = b + d'$, d in $H^{2k}(M; Z_2)$. Then

$$\begin{aligned} P(a) &= P(b + d') \\ &= P(b) + P(d') + (b \cup d')'. \end{aligned}$$

Thus we must show

$$P(d') + (b \cup d')' = 0.$$

But

$$\begin{aligned} (b \cup d')' &= ((b \bmod 2) \cup d)'' \\ &= (v_{2k}(M) \cup d)'' \\ &= (d \cup d)'' , \text{ by definition of the Wu class} \\ &= P(d'). \end{aligned}$$

(The last equality follows immediately from the cochain definition of the Pontrjagin Square.) This verifies Lemma 5.

Thus $P(x)[M]$ is independent of the choice of x (provided $x \bmod 2 = v_{2k}(M)$). A convenient choice for x is given by

LEMMA 6. *If M^{4k} is an oriented Poincaré complex, then $v_{2k}(M)$ is the reduction of an integral cohomology class.*

Proof. (E. Thomas) Let K be the subgroup of $H^{2k}(M; Z_2)$ consisting of all classes which are the mod2 reduction of an integral class. Let L be the subgroup of $H^{2k}(M; Z_2)$ consisting of all classes whose cup product with the mod2 reduction of every

torsion class in $H^{2k}(M; Z)$ is zero. Clearly $K \subseteq L$. But an easy counting argument, using Poincaré duality, shows that $\dim K = \dim L$, so $K = L$. Since $v_{2k}(M)$ is in L , it is in K , and Lemma 6 is proved.

We now prove Theorem 2. Let z in $H^{2k}(M; Z)$ be a class whose mod 2 reduction is $v_{2k}(M)$. By a well-known property of bilinear forms (see [4]), $(z \cup z)[M] = \sigma(M) \pmod 8$. Let x in $H^{2k}(M; Z_4)$ be the mod 4 reduction of z . Then $P(x)[M] = (z \cup z)[M] \pmod 8 = \sigma(M) \pmod 8$, which proves Proposition 4 and Theorem 2.

2. The Oriented Case

Let Ω_{4k}^{PD} be the cobordism group of oriented $4k$ -dimensional Poincaré complexes. There is an exact sequence ([5], [8], [9])

$$0 \rightarrow Z \xrightarrow{i} \Omega_{4k}^{PD} \xrightarrow{j} \pi_{4k} \text{MSG} \rightarrow 0.$$

The infinite cyclic group is generated by the closed Milnor manifold of signature 8. Let $\sigma: \Omega_{4k}^{PD} \rightarrow Z_8$ be the signature homomorphism reduced mod 8. Since $\sigma i = 0$, there is a homomorphism $\bar{\sigma}: \pi_{4k} \text{MSG} \rightarrow Z_8$ such that $\bar{\sigma} j = \sigma$.

Now the spectrum MSG is a product of Eilenberg-MacLane spectra. (We need this only at the prime 2: see [1].) Therefore there is a cohomology class t_{4k} in $H^{4k}(\text{MSG}; Z_8)$ such that for any g in $\pi_{4k} \text{MSG}$,

$$\bar{\sigma}(g) = g^*(t_{4k})[S^{4k}].$$

Let y_{4k} be the class in $H^{4k}(\text{BSG}; Z_8)$ corresponding to t_{4k} under the Thom isomorphism. Then if M^{4k} is an oriented Poincaré complex and $h: M \rightarrow \text{BSG}$ the classifying map for the normal spherical fibration, let c_M denote the cobordism class of M in Ω_{4k}^{PD} . Then $j(c_M) \in \pi_{4k} \text{MSG}$, and $j(c_M)_*[S^{4k}]$, the Hurewicz image of $j(c_M)$, corresponds to $[h_* M]$ under the Thom isomorphism. Hence

$$\begin{aligned} y_{4k}[M] &= h^*(y_{4k})[M] = \langle y_{4k}, h_*[M] \rangle \\ &= \langle t_{4k}, j(c_M)_*[S^{4k}] \rangle \\ &= \langle j(c_M)^*(t_{4k}), [S^{4k}] \rangle \\ &= \bar{\sigma}(j(c_M)) = \sigma(c_M) \\ &= \sigma(M) \pmod 8, \end{aligned}$$

which proves Theorem 3.

3. Proposition B

We show how the techniques of this paper imply Proposition B. Let M^0 be a smooth, compact, spin manifold of dimension $4k$ with boundary an exotic sphere.

Let M be the closed topological manifold formed by attaching a $4k$ -disk along the boundary of M^0 . Also, let $B\text{Spin}$ be the classifying space for stable vector bundles with a spin structure and $h: M^0 \rightarrow B\text{Spin}$ be the classifying map for the stable normal bundle.

According to E. Thomas [7], all 2-torsion in $H^*(B\text{Spin}; \mathbb{Z})$ is of order 2. Let v_{2k} be the Wu class in $H^{2k}(B\text{Spin}; \mathbb{Z}_2)$. Then $Sq^1(v_{2k})=0$ means that v_{2k} is the mod 2 reduction of an integer class z_{2k} in $H^{2k}(B\text{Spin}; \mathbb{Z})$. Define $z_{2k}(M)$ as $h^*(z_{2k})$. Then $(z_{2k}(M))^2[M]=\sigma(M) \bmod 8$. Let q_{2k} in $H^{2k}(B\text{Spin}; \mathbb{Q})$ be the class corresponding to z_{2k} under the inclusion of \mathbb{Z} in \mathbb{Q} . Then $(q_{2k}(M))^2[M]=\sigma(M) \bmod 8$ and q_{2k} is a polynomial (with rational coefficients) in the Pontrjagin classes. Since $(q_{2k})^2$ is decomposable, this proves Proposition B.

REFERENCES

- [1] BROWDER, W., LIULEVICIUS, A., and PETERSON, F. P., *Cobordism theories*, Ann. of Math. 84 (1966), 91–101.
- [2] BRUMFIEL, G., *On the homotopy groups of BPL and PL/O*, Ann. of Math. 88 (1968), 291–311.
- [3] FRANK, D., *The signature defect and the homotopy of BPL and PL/O*, Comment. Math. Helv. 48 (1973), 525.
- [4] HIRZEBRUCH, F., NEUMANN, W. D., and KOH, S. S., *Differentiable manifolds and quadratic forms*, Lecture Notes in Pure and Applied Mathematics, Vol. 4, Marcel Dekker, New York, 1971.
- [5] LEVITT, N., *Poincaré duality cobordism*, Ann. of Math. 96 (1972), 211–244.
- [6] STEENROD, N. E. and EPSTEIN, D. B. A., *Cohomology Operations*, Ann. of Math. Studies 50, Princeton, 1962.
- [7] THOMAS, E., *On the cohomology groups of the classifying space of the stable spinor group*, Bol. Soc. Mat. Mexicana (2) 7 (1962), 57–69.
- [8] JONES, L. *Patch spaces*, Ann. of Math. 97 (1973), 306–343.
- [9] QUINN, F., *Surgery on Poincaré and normal spaces*, Bull. Amer. Math. Soc. 78 (1972), 262–267.

State University of New York at Stony Brook.

Received May 5, 1973.