

AN INVARIANT OF QUADRATIC FORMS MOD 8

BY

F. VAN DER BLIJ

(Communicated by Prof. H. FREUDENTHAL at the meeting of April 25, 1959)

In a short note W. LEDERMANN [1] proved the

Theorem 1. *If f is an integral quadratic form in n variables with odd determinant D , then*

$$(1) \quad f(w, w) \equiv D + \tau - \text{sgn } D \pmod{4},$$

for every integral vector w , which satisfies $f(x, x) \equiv f(x, w) \pmod{2}$ for all integral vectors x . Here τ is the signature of f and $\text{sgn } D = D|D|^{-1}$.

As a special case he found that if f is unimodular we have

$$(2) \quad f(w, w) \equiv \tau \pmod{4},$$

this was a corollary of topological investigations of F. HIRZEBRUCH and H. HOPF [2]. Now it may be readily verified that $f(w, w)$ is an invariant mod 8, since from $f(x, w) \equiv f(x, t) \pmod{2}$ we find $t = w + 2z$, with an integral vector z and $f(t, t) = f(w, w) + 4f(w, z) + 4f(z, z) \equiv f(w, w) \pmod{8}$.

From the theory of the transformation of theta functions [3] one finds if f is definite positive and unimodular that

$$(3) \quad f(w, w) \equiv n \pmod{8}.$$

And now one can ask for a suitable generalization of (1) modulo 8.

We define a gaussian sum by

$$(4) \quad G = \sum_{x \in M'/M} e^{\pi i f(x + \frac{1}{2}w, x + \frac{1}{2}w)},$$

here M is the lattice of all integral vectors and M' is the dual one with respect to f , that is to say M' consists of all vectors z such that $f(z, a)$ is integral for all $a \in M$. Using classical theorems on Fourier expansions one finds

$$G = |D| \sum_{v \in M} \int_T e^{\pi i f(t + \frac{1}{2}w, t + \frac{1}{2}w)} e^{-2\pi i f(t, v)} dt,$$

where T is a fundamental domain of M .

$$G = |D| \sum_{v \in M} \int_T e^{\pi i f(t - v + \frac{1}{2}w, t - v + \frac{1}{2}w)} dt,$$

$$G = |D| \int e^{\pi i f(z, z)} dz.$$

here the integration must be extended for all coordinates of z from $-\infty$ to ∞ . Transforming f in a diagonal form one can easily calculate this integral. We find then

$$(5) \quad G = |D|^{\frac{1}{2}} e^{\pi i \tau / 4}.$$

On the other hand we find

$$G = e^{\frac{1}{2} \pi i f(w, w)} \sum_{x \in M'/M} e^{\pi i f(x, x)} e^{\pi i f(x, w)}$$

and since D is odd we may replace x by $2y$ and thus

$$G = e^{\frac{1}{2} \pi i f(w, w)} \sum_{y \in M'/M} e^{4 \pi i f(y, y)}.$$

Now we write

$$(6) \quad G_0 = \sum_{y \in M'/M} e^{4 \pi i f(y, y)}$$

and we have proved:

Theorem 2.

$$e^{\frac{1}{2} \pi i f(w, w) - \tau} = |D|^{\frac{1}{2}} G_0^{-1}.$$

We consider some special cases. First let $D = \pm 1$, then $G_0 = 1$ and hence

$$(7) \quad f(w, w) \equiv \tau \pmod{8}.$$

Further, one can easily calculate G_0^2 (e.g. by transforming f in a diagonal form modulo D)

$$G_0^2 = \left(\frac{-1}{|D|} \right) |D|$$

and thus we obtain

$$(8) \quad f(w, w) \equiv |D| + \tau - 1 \pmod{4}.$$

Since $|D| + \tau - 1 \equiv D + n - 1 \equiv D + \tau - \text{sgn } D \pmod{4}$ this is equivalent with theorem 1.

Theorem 2 gives a relation between certain local invariants of f ; $f(w, w) \pmod{8}$ is an invariant for the place 2, which can be described as a character of the local Witt group which has on the form αx^2 , with odd α the value $e^{\frac{1}{2} \pi i \alpha}$ and which equals 1 on the even forms with odd determinant.

If $p|D$ we can write f at the place p as $\sum_{i=1}^n f_i p^{\alpha_i} x_i^2$ and this prime gives as contribution to the expression $G_0^{-1} |D|^{\frac{1}{2}}$ a factor (we use the multiplicity properties of gaussian sums)

$$\varphi_p(f) = \prod_{i=1}^n \left(\frac{f_i}{p^{\alpha_i}} \right) e^{\frac{1}{2} \pi i (p^{\alpha_i} - 1)}.$$

We can express φ_p in a simple way in the local invariants of f . With every p -adic form f , there are two forms f_0 and f_1 over the field with p elements, such that the Witt class of f is determined by the form $f_0^* + p f_1^*$ where f_0^* and f_1^* are p -adic forms, which give by restriction modulo p the forms f_0 and f_1 (vid [4]).

The Witt group of the field with p elements consists of four elements. If $p \equiv 3 \pmod{4}$ the Witt group is a cyclic group, if $p \equiv 1 \pmod{4}$ this group is the direct product of two cyclic groups of order two. Let $\varepsilon_p(f_1)$ be the character of this group which equals $e^{i\pi i(p-1)}$ for $f_1(x, x) = x^2$ and in the second case also $-e^{i\pi i(p-1)}$ for εx^2 , where ε is a non-square modulo p . Now it can be readily verified that $\varphi_p(f) = \varepsilon_p(f_1)$. Thus we have proved

Theorem 3.

$$e^{i\pi i(f(w, w) - \tau)} = \prod_{p|D} \varphi_p(f),$$

where $\varphi_p(f) = \varepsilon_p(f_1)$ is a character of the Witt group over the p -adic field.

This is an example of a relation between the local invariants of quadratic forms. Another example is given by EICHLER [5]. It may be an interesting question to ask for other relations of this type.

*Mathematical Institute of the
University of Utrecht.*

REFERENCES

1. LEDERMANN, W., An arithmetical property of quadratic forms. *Commentarii Mathematici Helvetici* 33, 34–37 (1959).
2. HIRZEBRUCH, F. und H. HOFF, Felder von Flächenelementen in 4 dimensionalen Mannigfaltigkeiten. *Math. Ann.* 136 (156–172) (1958).
3. BLIJ, F. VAN DER en J. H. VAN LINT, On some special theta functions. *Proc. Kon. Ak. Amsterdam A* 61 *Ind. Math.* 20, 508–513 (1958).
4. SPRINGER, T. A., Quadratic forms over fields with a discrete valuation. I, II *Proc. Kon. Ak. Amsterdam A* 58, 59, *Ind. Math.* 17, 18, 352–362, 238–246 (1955, 1956).
5. EICHLER, M., *Quadratische Formen und Orthogonale Gruppen*. Berlin 1952, 162.