

Sep. 2019, Bonn

①

## Singular stochastic nonlinear wave equations

### Stochastic nonlinear wave equation (SNLW)

with an additive space-time  
white noise.

$$(SNLW) \quad \left\{ \begin{array}{l} (\partial_t^2 - \Delta) u + u^k = \xi \\ (u, \partial_t u) \Big|_{t=0} = (u_0, u_1) \in N^s(\mathbb{T}^d) \end{array} \right.$$

on  $\mathbb{R}_+ \times \mathbb{T}^d$ ,  $\mathbb{T}^d = (\mathbb{R}/2\pi\mathbb{Z})^d$

$$N^s(\mathbb{T}^d) = H^s(\mathbb{T}^d) \times H^{s-1}(\mathbb{T}^d)$$

- $\xi$  = space-time white noise

= Gaussian process indexed by  $(t, x)$  s.t.

$$\mathbb{E}[\xi(t_1, x_1) \xi(t_2, x_2)] = \delta(t_1 - t_2) \delta(x_1 - x_2)$$

- Itô formulation: With  $v = \partial_t u$ ,

$$d \begin{pmatrix} u \\ v \end{pmatrix} + \left\{ \begin{pmatrix} 0 & -1 \\ -\Delta & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} 0 \\ u^k \end{pmatrix} \right\} dt = \begin{pmatrix} 0 \\ dW \end{pmatrix}$$

(2)

Here,  $W = L^2$ -cylindrical Wiener process

$$W(t) = \sum_{n \in \mathbb{Z}^d} \beta_n(t) e_n$$

$$\cdot e_n(x) = \frac{1}{(2\pi)^{d/2}} e^{inx}$$

$\cdot \{\beta_n\}_{n \in \mathbb{Z}^d}$  = family of mutually independent complex-valued Brownian motions

" $\mathcal{N} = \mathbb{Z}^d / 2$ " conditioned such that  $\beta_{-n} = \overline{\beta_n}, n \in \mathbb{Z}^d$

Convention :  $\text{Var}(\beta_n(t)) = t$ .

$$\text{i.e. } \beta_n = \underbrace{\text{Re } \beta_n + i \text{Im } \beta_n}_{\text{indep real B.M. } / \sqrt{2}}$$

$$\mathcal{N} = \bigcup_{k=0}^{d-1} (\mathbb{Z}^k \times \mathbb{Z}_+ \times \{0\}^{d-k-1})$$

Then,  $\{\beta_n\}_{n \in \mathcal{N}}$  is independent.

(3)

## Duhamel formulation (= mild formulation )

We say that  $u$  is a soln to  $SNLW$  if  $u$  satisfies

$$u(t) = \partial_t S(t) u_0 + S(t) u_1 - \int_0^t S(t-t') u^k(t') dt' \\ + \boxed{\int_0^t S(t-t') dW(t')} \quad \text{stochastic convolution.}$$

where  $S(t) = \frac{\sin(t|\nabla|)}{|\nabla|}$ ,  $|\nabla| = \sqrt{-\Delta}$

$$S(t)f(m) = \begin{cases} \frac{\sin(t|m|)}{|m|} \hat{f}(m), & m \neq 0 \\ t \hat{f}(0), & m = 0. \end{cases}$$

In the following, we actually study

$$\begin{cases} (\partial_t^2 + 1 - \Delta) u + u^k = \mathfrak{I} \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \end{cases}$$

corresponding to the Klein-Gordon eqn but we simply refer to this as  $SNLW$ .

(4)

Duhamel formula:

$$U(t) = 2tS(t)U_0 + S(t)U_1 - \int_0^t S(t-t')U^k(t')dt' \\ + \int_0^t S(t-t')dW(t).$$

Now,  $S(t) = \frac{\sin(t\langle\nabla\rangle)}{\langle\nabla\rangle}$ ,  $\langle\nabla\rangle = \sqrt{1-\Delta}$

$$\langle\cdot\rangle = \sqrt{1-1\cdot 1^2}$$

= Japanese bracket.

The only difference appears at the zeroth freq.

but it does not affect well-posedness theory

• Stochastic damped NLW (SdNLW)

$$\begin{cases} (\partial_t^2 + \underline{\partial_t} + 1 - \Delta)u + u^k = \sqrt{2}\xi \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \end{cases}$$

Itô formulation:

$$d\begin{pmatrix} u \\ v \end{pmatrix} + \begin{Bmatrix} 0 & -1 \\ 1-\Delta & 0 \end{Bmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} 0 \\ u^k \end{pmatrix} dt$$

$$= \begin{pmatrix} 0 \\ -v \end{pmatrix} dt + \begin{pmatrix} 0 \\ \sqrt{2}dW \end{pmatrix}$$

## Duhamel formulation:

(5)

$$u(t) = \mathcal{D}(t) u_0 + \mathcal{D}(t)(u_0 + u_1)$$

$$- \int_0^t \mathcal{D}(t-t') u^k(t') dt'$$

$$+ \sqrt{2} \int_0^t \mathcal{D}(t-t') dW(t'),$$

$$\text{where } \mathcal{D}(t) = \frac{e^{-\frac{t}{2}} \sin(t\sqrt{\frac{3}{4}-\Delta})}{\sqrt{\frac{3}{4}-\Delta}}$$

← In terms of local well-posedness, there is

no difference from SNLW (either wave or KG.)

Why care?

①  $\zeta = dW$  has regularity

$$-\frac{d}{2} - 2 \text{ in } \mathcal{X}$$

very rough.

$$-\frac{1}{2} - \varepsilon \text{ in } t'$$

expect

⇒ stoch. convolution  $\mathbb{F} = \int_0^t \frac{\sin((t-t')\langle \nabla \rangle)}{\langle \nabla \rangle} dW(t')$

has spatial regularity

$$1 - \frac{d}{2} - \varepsilon \quad (< 0 \text{ for } d \geq 2)$$

(6)

$\Rightarrow$  analytically challenging.

(3) SdNLW formally preserves the Gibbs measure  $\mu$ .

( $k \in 2\mathbb{N} + 1$ )

$$d\mu = Z^{-1} e^{-H(u,v)} du \otimes dv$$

$$= Z^{-1} e^{-\frac{1}{k+1} \int_{\mathbb{T}^d} u^{k+1} dx} e^{-\frac{1}{2} \|u\|_{H^1}^2} du$$

$$\times e^{-\frac{1}{2} \|v\|_{L^2}^2} dv$$

$$= Z^{-1} e^{-\frac{1}{k+1} \int_{\mathbb{T}^d} u^{k+1} dx} d\mu_s(u) \otimes d\mu_0(v),$$

where

$\int_{\mathbb{T}^d} u^{k+1} dx$ -measure

$H(u,v) =$  Hamiltonian (= energy) for NLW/NLKG

$$= \frac{1}{2} \int_{\mathbb{T}^d} |\langle \nabla \rangle u|^2 dx + \frac{1}{2} \int_{\mathbb{T}^d} v^2 dx + \frac{1}{k+1} \int_{\mathbb{T}^d} u^{k+1} dx.$$

$\mu_s =$  Gaussian meas with CM space  $H^s(\mathbb{T}^d)$

$$d\mu_s = Z_s^{-1} e^{-\frac{1}{2} \|u\|_{H^s}^2} du.$$

$$\Leftrightarrow u = \sum_{n \in \mathbb{Z}^d} \frac{g_n(\omega)}{\langle n \rangle^s} e^{im \cdot x}$$

$\{g_n\}$ , indep standard  $\mathbb{C}$ -valued Gaussian r.v.'s.  
s.t.  $\overline{g_{-n}} = g_n$ .

SdNLW

NLW in a vectorial form

(7)

$$d \begin{pmatrix} u \\ v \end{pmatrix} + \left\{ \begin{pmatrix} 0 & -1 \\ 1-\Delta & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} 0 \\ u^2 \end{pmatrix} \right\} dt$$

$$= \begin{pmatrix} 0 \\ -v \end{pmatrix} dt + \begin{pmatrix} 0 \\ \sqrt{2} dW \end{pmatrix}$$

Ornstein-Uhlenbeck process for  $v$

- NLW preserves the Gibbs measure  $f$ .

1-d : Friedlander '85

2-d : Oh-Thomann '17.

- OV preserves the white noise  $\mu_0$  for  $v$

and hence preserves the Gibbs measure  
 $f(du, dv)$ .

- $L$  = generator for SdNLW

$$= L_1 + L_2$$

$L_1$  = generator for NLW

$L_2$  = generator for OV.

$f$  invariant  $\Leftrightarrow L^* f = 0$  (8)

i.e.  $\int L F(u, v) d\mu(u, v) = 0$

$\Leftarrow L_1^* f = 0 \text{ AND } L_2^* f = 0.$

b/c  $L^* = L_1^* + L_2^*$

This can be made rigorous by considering the

"finite-dimensional" approximation:

(SdNLW<sub>N</sub>)  $(\partial_t^2 + \partial_x^2 + 1 - \Delta) u^N + P_{\leq N}((P_{\leq N} u^N)^k) = \sqrt{2} \xi.$

$\Leftarrow$  {(i) finite dim'l SdNLW on  $\{ |m| \leq N \}$  decoupled}  
{(ii) linear stochastic damped wave dynamics on high freq}

i.e. this preserves

$$\begin{aligned} d\mu_N &= Z_N^{-1} e^{-\frac{1}{k+1} \int_{\mathbb{T}} (P_{\leq N} u)^{k+1} dx} d\mu_1(u) \otimes d\mu_0(v) \\ &= Z_N^{-1} \left( e^{-\frac{1}{k+1} \int (P_{\leq N} u)^{k+1} dx} d\mu_{1,N}(u) \otimes d\mu_{0,N}(v) \right) \\ &\quad \times \left( d\mu_{1,N}^\perp(u) \otimes d\mu_{0,N}^\perp(v) \right) \end{aligned}$$

(ii)

(9)

## Main Goal:

Prove local well-posedness of  $S^dNLW$  along with a good approximation property by  $S^dNLW_N$ .

$\Rightarrow$  Then, we can apply Bourgain's invariant measure argument to obtain

- almost sure global well-posedness of  $S^dNLW$
- invariance of the Gibbs measure  $\rho$ .

$$\begin{array}{ccc} \rho_N & \xrightarrow{\quad} & \rho \\ | & & : \leftarrow \text{inv.} \\ u^N & \xrightarrow{\quad} & u \end{array}$$

See Bourgain '94, '96, Burq-Tzvetkov '08.

Idea: Use formal invariance of  $\rho$  as a replacement of a conservation law.

(• parabolic  $\Phi_3^4$ -model: Hairer-Matetski '18)

⇒ shows SdNLW is a stochastic quantization equation (SQE) for (10)

$\Phi_d^{k+1}$ -measure for  $u$   $\otimes$  white noise  $\mu_0$  for  $v$ .

- SdNLW is called the canonical stochastic quantization equation. (Ryang, Saito, Shigemoto '85.)  
    $\Leftarrow$  hyperbolic  $\Phi_d^{k+1}$ -model.

Parabolic counterpart:

Parabolic  $\Phi_d^{k+1}$ -model / stochastic quantization eqn

$$(\partial_t - \Delta) u + u^{k+1} = \tilde{\zeta} \text{ on } \mathbb{T}^d$$

- d=2: Da Prato - Debussche '03.
- d=3: (k=3)

Hairer '15: regularity structure

Gubinelli - Imkeller - Perkowski '15:  
paracontrolled distributions  
(Catellier - Chouk '18)

Kupiainen '16: renormalization group method.

. In the following, we focus on local well-posedness

⇒ We only consider the (second) SNLW.

Chap 1: SNLW in 2-d.

Chap 2: quadratic SNLW in 3-d.

Chap 3: cubic SNLW of Hartree-type in 3-d.  
(with Mamoru Okamoto)

← I did not get to cover this part.

## Chapter 0: Preliminaries

(SNLW)

$$U(t) = \mathcal{A}_t S(t) U_0 + S(t) U_1 - \int_0^t S(t-t') U^* H' dt' + \Psi$$

$\Psi$  = stochastic convolution

$$\begin{aligned} &= \int_0^t S(t-t') dW(t') \quad \text{Wiener integral} \\ &= \sum_{m \in \mathbb{Z}^d} e_m \int_0^t \frac{\sin((t-t')\langle m \rangle)}{\langle m \rangle} d\beta_n(t') \end{aligned}$$

• Wiener integral (R-valued case)

Given deterministic  $f \in L^2([a, b])$ ,

define  $I(f) = \int_a^b f dB$  by the left endpt Riemann sum.

$\Rightarrow I(f)$  is a mean 0 Gaussian r.v.

with  $\text{Var}(I(f)) = \|f\|_{L^2([a, b])}^2$ .

i.e.

$I: L^2([a, b]) \rightarrow L^2(\Omega)$  is an isometry  
(onto its image.)

- The following lemma will be useful in studying regularities of various random distributions

- $\mathcal{H}_k$  = homogeneous Wiener chaoses of order  $k$ .

$$\mathcal{H}_{\leq k} = \bigoplus_{j=1}^k \mathcal{H}_j.$$

(Think of the  $L^2(\Omega)$ -completion of polynomials in  $B_n$ 's of degree  $\leq k$ .

## Lemma 1 (Wiener chaos estimate)

Let  $k \in \mathbb{N}$ . Then, we have

$$\|X\|_{L^p(\Omega)} \leq (p-1)^{\frac{1}{2}} \|X\|_{L^2(\Omega)}$$

for any  $p \geq 2$  and  $X \in \mathcal{H}_{\leq k}$ .

( $\Leftarrow$  Nelson's hypercontractivity of the OU semigroup.

Lemma 2: Let  $\{X_N\}_{N \in \mathbb{N}}$  and  $X$  be spatially

homogenous stochastic processes:  $\mathbb{R}_+ \rightarrow \mathcal{D}'(\mathbb{T}^d)$

i.e. for any  $x_0 \in \mathbb{T}^d$ ,

$$\{X(\cdot, t)\}_{t \in \mathbb{R}_+} \text{ and } \{X(x_0 + \cdot, t)\}_{t \in \mathbb{R}_+}$$

have the same law.

- Suppose that  $\exists k \in \mathbb{N}$  s.t.  $X_N(t)$  and  $X(t)$  belong to  $\mathcal{H}_{\leq k}$  for each  $t \in \mathbb{R}_+$ .

Then,

ii)  $t \in \mathbb{R}_+$ . If  $\exists s_0 \in \mathbb{R}$  such that

$$(*) \quad \mathbb{E} [\|\widehat{X}(t, n)\|^2] \lesssim \langle n \rangle^{d-2s_0}$$

for any  $n \in \mathbb{Z}^d$ , then

$$X(t) \in W^{s, \infty}(\mathbb{T}^d), \quad s < s_0, \text{ a.s.}$$

↑  
Bessel potential space

$$\begin{aligned} \|f\|_{W^{s,p}} &= \|\langle \nabla \rangle^s f\|_{L^p} \\ &= \|\mathcal{F}^{-1}(\langle n \rangle^s \widehat{f}(n))\|_{L^p}. \end{aligned}$$

Moreover, if  $\exists \theta > 0$  s.t.

$$\mathbb{E} [\|\widehat{X}_N(t, n) - \widehat{X}(t, n)\|^2] \lesssim N^{-\theta} \langle n \rangle^{d-2s_0}$$

for any  $n \in \mathbb{Z}^d$  and  $N \geq 1$ , then

$$X_N \rightarrow X \text{ in } W^{s, \infty}(\mathbb{T}^d), \quad s < s_0, \text{ a.s.}$$

(ii) Given  $h \in \mathbb{R}$ , define the difference operator  $\delta_h$

$$\text{by } \delta_h X(t) = X(t+h) - X(t).$$

Let  $T > 0$  and suppose (i) holds on  $[0, T]$ .

(15)

If  $\exists \sigma \in (0, 1)$  s.t.

$$\mathbb{E} [|\delta_h \hat{X}(t, m)|^2] \lesssim \langle m \rangle^{-d-2s_0+\sigma} = |h|^\sigma$$

for any  $m \in \mathbb{Z}^d$ ,  $t \in [0, T]$ ,  $|h| \leq 1$  with  
 $0 \leq t+h \leq T$ ,

then,

$$X \in C([0, T]; W^{s, \infty}(\mathbb{T}^d))$$

for  $s < s_0 - \frac{\sigma}{2}$ , a.s.

Furthermore, if  $\exists \theta > 0$  s.t.

$$\begin{aligned} \mathbb{E} [|\delta_h \hat{X}_N(t, m) - \delta_h \hat{X}(t, m)|^2] \\ \lesssim N^{-\theta} \langle m \rangle^{-d-2s_0+\sigma} |h|^\sigma, \end{aligned}$$

for any  $m \in \mathbb{Z}^d$ ,  $t \in [0, T]$ ,  $|h| \leq 1$ , and  $N \geq 1$ ,

then  $X_N$  converges to  $X$  in  $C([0, T]; W^{s, \infty}(\mathbb{T}^d))$

for  $s < s_0$ , a.s.

See Mourrat-Weber-Xu, Oh-Okamoto-Tzvetkov

Idea of the proof:

translation invariance

$$\Rightarrow \mathbb{E} [\hat{X}(t, n) \hat{X}(t, m)] = 0$$

for  $n+m \neq 0$ .

(real-valued setting.)

(16)

$$\cdot \mathbb{E}[\hat{X}(t, n) \hat{X}(t, m)]$$

$$= \iint \underbrace{\mathbb{E}[X(t, x) X(t, y)]}_{= F(t, x-y)} \underbrace{e_{-n}(x) e_{-m}(y)}_{= e_{-(m+n)}(x) e_m(x-y)} dy dx$$

$$= \hat{F}(t, m) \int_{\mathbb{T}^d} e^{-i(m+n) \cdot x} dx$$

$$= 0 \text{ for } n+m \neq 0.$$

Suppose  $\circledast$  holds. Then, for  $p \gg 1$ ,

~~Suppress t~~

$$\| \|X\|_{W^{s, \infty}} \|_{L^p(\Omega)} \stackrel{\text{Sobolev}}{\lesssim} \| \|X\|_{W^{s+\varepsilon, p}} \|_{L^p(\Omega)}$$

$$= \| \| \langle \nabla \rangle^{s+2} X(x) \|_{L^p(\Omega)} \|_{L_x^p}$$

$$\stackrel{\text{Lem } 1}{\leq} p^{k/2} \| \sum_{n \in \mathbb{Z}^d} \langle n \rangle^{s+2} \hat{X}(n) e_n(x) \|_{L_x^2(\Omega)} \|_{L_x^p}$$

$$= \left( \sum_{n \in \mathbb{Z}^d} \langle n \rangle^{2(s+\varepsilon)} |\hat{X}(n)|^2 \right)^{1/2}$$

$$\leq C_p \left( \sum_{n \in \mathbb{Z}^d} \langle n \rangle^{2(s+\varepsilon)-d-2s_0} \right)^{1/2} < \infty$$

iff  $s + \varepsilon - s_0 < 0$  i.e.  $s < s_0$ .  $\square$

(17)

Lemma 3:

$$\Psi \in C(\mathbb{R}_+; W^{1-\frac{d}{2}-, \infty}(\mathbb{T}^d)), \text{ a.s.}$$

Moreover,

$$\Psi_N = P_{\leq N} \Psi \rightarrow \Psi$$

in  $C(\mathbb{R}_+; W^{1-\frac{d}{2}-, \infty}(\mathbb{T}^d)), \text{ a.s.}$ 

↑ compact-open topology

Pf: We only verify  $\otimes$  in Lemma 2.

$$\begin{aligned} \mathbb{E} [|\hat{\Psi}(t, n)|^2] &= \int_0^t (\sin((t-t')\langle n \rangle))^2 dt' \\ &\lesssim_t \langle n \rangle^{-2} = \langle n \rangle^{-d-2(1-\frac{d}{2})} \\ &\quad \underset{= s_0}{=} . \end{aligned}$$

• For temporal regularity (ii), use mean value thm in time  $\square$ • When  $d = 1$ ,

$$\Psi \in C(\mathbb{R}_+; W^{\frac{1}{2}-, \infty}(\mathbb{T}^d)), \text{ a.s.}$$

 $\Rightarrow$  LWP of SNLW is trivial.• For  $d \geq 2$ ,  $\Psi(t)$  is NOT a function.  
only a distribution. $\Rightarrow$  ? issue in making sense of  $(\Psi(t))^k$ .  
(and hence  $u^k$ .) $\Rightarrow$  Need to renormalize the nonlinearity

(18)

## Chap 1: SNLW in 2-d

$$u(t) = \partial_t S(t) u_0 + \delta(t) u_1 - \int_0^t S(t-t') u^k(t') dt' \\ + \Psi.$$

Picard 2<sup>nd</sup> iterate :  $\int_0^t S(t-t') \Psi^k(t') dt'$

$\uparrow$  does NOT make sense.

Triviality:  $u_\varepsilon$  = soln to SNLW with regularized noise  $\tilde{\zeta}_\varepsilon = \gamma_\varepsilon * \tilde{\zeta}$ .

Without a proper renormalization,

$$u_\varepsilon \rightarrow \lim_{\varepsilon} \text{soln (or 0)}$$

Albeverio - Haba - Russo '96  
Oh - Okamoto - Robert '19.

i.e. renormalization is necessary in order to have a non-trivial limit.

Truncated SNLW:

$$(\partial_t^2 + 1 - \Delta) u_N + u_N^k = P_{\leq N} \tilde{\zeta}.$$

$\Rightarrow$  Truncated convolution  $\Psi_N = P_{\leq N} \Psi$ .

$\uparrow$  smooth in  $x$ .

(19)

Fix  $(t, x) \in \mathbb{R}_+ \times \mathbb{T}^2$ .

$\Sigma_N(t, x)$  = mean 0 Gaussian r.v. with

$$\text{variance } \sigma_N(t) = \mathbb{E}[\Sigma_N^2(t, x)]$$

$$= \sum_{|m| \leq N} \int_0^t \frac{\langle m \rangle (t-t') \langle m \rangle^2}{\langle m \rangle^2} dt'$$

$\sim t \log N \rightarrow \infty$  as  $N \rightarrow \infty$ .

Da Prato - Debussche trick ('02-'03)

McKean '95

Bourgain '96

$$\text{Duhamel} \Rightarrow u_N = v_N + \Sigma_N$$

↑ postulate  $v_N$  is smoother.

Binomial:

$$u_N^k = \sum_{j=0}^k \binom{k}{j} \Psi_N^j v_N^{k-j}$$

↑ does NOT have a limit (as  $N \rightarrow \infty$ )

Wick renormalization: Replace  $\Psi_N^j$  by

$$:\Psi_N^j(t, x): = H_j(\Sigma_N(t, x); \sigma_N(t))$$

↑ pairwise operation .

↑ time-dependent

(20)

$H_j(x; \sigma)$  = Hermite poly of deg  $j$ .

defined by the generating function:

$$\underline{G(t, x; \sigma)} = e^{tx - \frac{1}{2}\sigma t^2} = \sum_{j=0}^{\infty} \frac{t^j}{j!} H_j(x; \sigma)$$

Prop 4:  $T > 0$ .

$\left\{ \mathbb{E}_N^j \right\}_{N \in \mathbb{N}}$  is Cauchy in  $L^p(\Omega; C_T W_x^{-\varepsilon, \infty})$   
 $p < \infty$

also almost surely in  $C([0, T]; W^{-\varepsilon, \infty}(T^2))$

Pf: We only verify  $\otimes$  in Lemma 2.  
 (unif in  $N \geq 1$ .)

Recall: For mean 0 Gaussian r.v.'s  $f$  and  $g$  with variances  $\sigma_f^2$  and  $\sigma_g^2$ , we have

$$\begin{aligned} \mathbb{E}[H_j(f; \sigma_f) H_k(g; \sigma_g)] \\ = \delta_{jk} \cdot j! \left\{ \mathbb{E}[fg] \right\}^j. \end{aligned}$$

$\Leftarrow$  follows from computing

$$\int_{\Omega} G(t, f; \sigma_f) G(s, g; \sigma_g) dP$$

- directly
- in terms of Hermite poly } compare coeff.

(21)

$$\mathbb{E} \left[ \left| \Psi_N^j(t, m) \right|^2 \right]$$

$$= \iint \mathbb{E} \left[ \left| \Psi_N^j(t, x) \right|^2 \right] e_n(y-x) dx dy$$

$$= j! \left\{ \mathbb{E} [ \Psi_N(t, x) \Psi_N(t, y) ] \right\}^j$$

$$= \sum_{\substack{m \in \mathbb{Z}^2 \\ |m| \leq N}} \int_0^t \frac{(\sin((t-t')\langle m \rangle))^2}{\langle m \rangle^2} dt' e_m(x-y)$$

$$\lesssim \sum_{n_1 + \dots + n_j = n} \frac{1}{\langle m_1 \rangle^2 \dots \langle m_j \rangle^2} \lesssim \frac{1}{\langle m \rangle^{2-\varepsilon}}$$

As before, the time difference in (ii) can be established by MVT  $\square$

Define

$$:u_N^k: = \sum_{j=0}^k \binom{k}{j} : \Psi_N^j: v_N^{k-j}$$



$$:u^k: = \sum_{j=0}^k \binom{k}{j} : \Psi^j: v^{k-j}$$

smoother.

for  $u = \Psi + v$

(22)

$\Rightarrow$  Renormalized SNLW (rSNLW)

$$\begin{cases} (\partial_t^2 + 1 - \Delta) u + :u^k: = \xi \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \end{cases}$$

Really means

XX

$$\begin{cases} (\partial_t^2 + 1 - \Delta) v + \sum_{j=0}^k \binom{k}{j} :v^j: v^{k-j} = 0 \\ (v, \partial_t v)|_{t=0} = (u_0, u_1). \end{cases}$$

Theorem 5 : (Gubinelli-Koch-Oh '18)

Let  $k \geq 2$ . Then, rSNLW is locally well-posed  
in  $H^\delta(\mathbb{T})$  for

scaling conformal

$$(i) k \geq 4 : \delta \geq s_{\text{crit}} = \max\left(1 - \frac{2}{k-1}, \frac{3}{4} - \frac{1}{k-1}, 0\right)$$

$$(ii) k = 2, 3 : \delta > s_{\text{crit}}.$$

Idea: Solve the fixed pt problem XX for  $v$  by

① Strichartz estimates (on lin solns)

② Prop 4 on  $:v^j:$

③ product estimates.

(23)

(i) fractional Leibniz rule.

$$(ii) \quad \frac{1}{p} + \frac{1}{q} \leq \frac{1}{r} + \frac{s}{d}$$

$$\Rightarrow \|\langle \nabla \rangle^{-s} (fg)\|_{L^r} \lesssim \|\langle \nabla \rangle^{-s} f\|_{L^p} \|\langle \nabla \rangle^s g\|_{L^q}$$

A typical term

$$\|\langle \nabla \rangle^{-s} (\Psi^j v^{k-j})\|$$

$$\begin{aligned} &\stackrel{(ii)}{\lesssim} \|\langle \nabla \rangle^{-s} : \Psi^j : \| \underbrace{\|\langle \nabla \rangle^s (v^{k-j})\|}_{\stackrel{(ii)}{\lesssim} \|\langle \nabla \rangle^s v\| \|v\|^{k-j-1}} \\ &\quad \stackrel{(ii)}{\lesssim} \|\langle \nabla \rangle^s v\| \|v\|^{k-j-1} \end{aligned}$$

$$\begin{aligned} u &= \Psi + v \\ &\in \Psi + C([0, T]; H^r(\mathbb{T}^2)), \quad r = s \wedge (1 - \varepsilon) \\ &\subset C([0, T]; H^{-\varepsilon}(\mathbb{T}^2)). \quad \# \varepsilon > 0 \end{aligned}$$

Back to page ⑨ : LWP of renormalized SdNLW

$\Rightarrow$  a.s. GWP of renormalized SdNLW  
 $(=$  hyperbolic  $\Phi_2^{k+1}$ -model  $)$

and invariance of the Gibbs measure

(by applying Bourgain's inv. meas. argument.)

- unique ergodicity (cubic) : L. Tolomeo '19.

Q: What about (deterministic)  
global well-posedness?

- In the heat case, one simply estimates  $\partial_t \|v\|_{L^p}^p$ .
- In the dispersive setting, we need to rely on the energy but the situation is much more intricate.

We only consider the defocusing cubic case.

$$(\partial_t^2 + 1 - \Delta)v + v^3 + 3v^2\Psi + 3v:\Psi^2 + :\Psi^3 = 0$$

rough perturbation

Two difficulties:

①  $v(t) \notin H^1(\mathbb{T}^2)$ .

⇒ can not use the energy

$$H(v, \partial_t v) = \frac{1}{2} \int |\nabla v|^2 dx + \frac{1}{2} \int (\partial_t v)^2 + \frac{1}{4} \int v^4 dx.$$

• We need to smooth  $v$ .

⇒ I-method in the stochastic setting.

② Even if  $v$  were in  $H^1$ ,  $v$  does not satisfy

(deterministic) NLW, i.e.  $H(v)$  is NOT conserved.

(25)

- If a noise is a bit smoother

(think of  $\langle \nabla \rangle^2 \Psi \in C_t L_x^2$ ) s.t.  $v \in C_t H_x^1$

then

$$\begin{aligned} \partial_t H(v) &= \int_{\mathbb{T}^2} 2\pi v \underbrace{\left( (\partial_t^2 + 1 - \Delta)v + v^3 \right)}_{= -(v + \Psi)^3} dx \\ &\quad \leftarrow \text{No need for renormalization} \end{aligned}$$

$\sim v^2 \Psi + v \Psi^2 + \Psi^3$

$v^3$ -term is gone!!

$$\stackrel{C-S}{\lesssim} \left( H(v) \right)^{1/2} \left( \|\Psi\|_{C_t L_x^\infty}^2 \int v^4 dx + \|\Psi\|_{C_t L_x^6}^6 \right)^{1/2}$$

$$\leq C(T, \Psi) H(v)$$

$\Rightarrow$  Gronwall. (Bing-Tzvetkov '14 in the random data well-posedness theory on 3-d cubic NLW)

• I-method (= method of almost conservation law) (26)


• CKSTT '02

(after Bourgain's high-low method '98)

Let  $N \in \mathbb{N}$ . Define

$$0 < s < 1$$

$$m_N(m) = \begin{cases} 1, & |m| \leq N \\ \frac{N^{1-s}}{|m|^{1-s}}, & |m| > 2N \end{cases}$$



Set

$I = I_N$  = Fourier multiplier op with  $m_N$ .

low freq : identity

high freq : fractional integration

def & Lp theory

$$\cdot \|If\|_{W^{a+\sigma, p}} \lesssim N^\sigma \|f\|_{W^{a, p}}$$

#  $0 \leq \sigma \leq 1-s$   
#  $1 < p < \infty$

$$\cdot \|f\|_{H^s} \lesssim \|If\|_{H^1} \lesssim N^{1-s} \|f\|_{H^s}$$

(27)

$\Rightarrow Iv$  does not satisfy NLW in 2 ways

① b/c of I

② b/c of the terms with  $\Psi$ .

$$H(Iv)(t) - H(Iv)(0)$$

$$= \int_0^t \int (\partial_t Iv) \underbrace{\left( (\partial_t^2 + 1 - \Delta) Iv + (Iv)^3 \right)}_{= -I(v + \Psi)^3}$$

$$= -I(v^3 + 3v^2\Psi + 3v\Psi^2 + \Psi^3)$$

$$= -I(v^3 + 3v^2\Psi + 3v\Psi^2 + \Psi^3)$$

$$= \int_0^t \int_x (\partial_t Iv) \left( -Iv^3 + (Iv)^3 \right)$$

↗ I-method part  
(commutator part)

$$- 3 \int_0^t \int_x (\partial_t Iv) I(v^2\Psi)$$

$$- 3 \int_0^t \int_x (\partial_t Iv) I(v\Psi^2)$$

$$- \int_0^t \int_x (\partial_t Iv) I(\Psi^3)$$

Gronwall part

$$=: A_1 + A_2 + A_3 + A_4$$

## Theorem 6: (GKO-Tolomeo '18)

The defocusing cubic  $r\text{-SNLW}$  is globally well-posed in  $\mathcal{H}^s(\mathbb{T}^2)$ ,  $s > \frac{4}{5}$ .

### Sketch of the proof:

#### ① Commutator estimates

Lemma 7:  $k \leq 3$

$$\| (If)^k - I(f^k) \|_{L^2} \lesssim N^{-\frac{1-k(1-s)}{2}} \| If \|_{H^1}^k$$

one of the freq. must be high.

Lemma 8: Given  $\gamma > 0$ ,  $0 < \sigma < 1$ ,

$\exists \gamma(\gamma) > 0$  and  $p(\gamma) \gg 1$  s.t.

$$\| (If)(Ig) - I(fg) \|_{L^2} \lesssim N^{\gamma - \frac{1-\sigma}{2}} \| f \|_{H^{1-\sigma}} \| g \|_{W^{-\gamma, p}}$$

Lemma 9:  $k = 0, 1, 2$ .  $\forall \gamma, \exists \gamma, p$  s.t.

$$\| I(v^k : \Psi^{3-k} : ) - (Iv)^k I(: \Psi^{3-k} :) \|_{L^2}$$

$$\lesssim N^{-\frac{1-k(1-s)}{2} + \gamma} \| Iv \|_{H^1}^k \| : \Psi^{3-k} : \|_{W^{-\gamma, p}}$$

( $\Leftarrow$  Lemmas 7 & 8)

Note: We may lower the regularity by using spacetime estimates.

(2) On the stochastic term:

Lemma 10:  $p < \infty$

$$\left( \mathbb{E} \| I\Psi \|_{L^p_{T,x}}^p \right)^{\frac{1}{p}} \lesssim p^{1/2} T^{1/2 + 1/p} (\log N)^{1/2}.$$

Pf: Separately estimate  $\underbrace{I P_{\leq N} \Psi}_{= P_{\leq N} \Psi}$  and  $I P_{> N} \Psi$ .

$\Rightarrow$  Fubini & Chebyshov,

$$P \left( \| I\Psi \|_{L^p_{T,x}} > \lambda \right) \lesssim \frac{p^{p/2} T^{\frac{p}{2} + 1} (\log N)^{p/2}}{\lambda^p}.$$

□

For Gronwall part, we have

Lemma 11 : (i)  $k=0, 1$ ,  $\forall 0 < \theta \leq 1-s$

(30)

$$\left| \int_{\mathbb{T}^2} (\partial_t IV) (IV)^k I \left( : \Psi^{3-k} : \right) \right|$$

$L_x^2 \nearrow \quad \searrow L_x^4$

$\approx N^\theta (1 + H(IV)^{\frac{3}{4}}) \| : \Psi^{3-k} : \|_{L_T^\infty W_x^{-\theta, 4}}$

BAD mapping property of  $I$ .

(ii) ( $k=2$ )  $\exists c > 0$  s.t.  $\forall 0 < \gamma < 1/8$ ,

$$\left| \int_{t_1}^{t_2} \int_{\mathbb{T}^2} (\partial_t IV) (IV)^2 (I\Psi) dx dt \right|$$

$$\lesssim \left( 1 + (t_2 - t_1) + \int_{t_1}^{t_2} H(V)^{1+c\gamma} dt \right) \| I\Psi \|_{L_{(t_1, t_2), x}^{\gamma-1}}$$

$\leftarrow$  interpolation & Sobolev

(31)

Putting together, we have

$$H(IV)(t) - H(IV)(t_0)$$

$$= \int_{t_0}^t \int_x (\partial_t IV) \underbrace{\left( -IV^3 + (IV)^3 \right)}_{\text{Lemma 7}} \, dx \, dt \quad \xleftarrow{C-S}$$

$$- 3 \int_{t_0}^t \int_x (\partial_t IV) \left( \underbrace{I(v^2 \Psi) - (IV)^2 I\Psi}_{\text{Lemma 9}} \right) \, dx \, dt \quad \xleftarrow{\text{commutator terms}}$$

$$- 3 \int_{t_0}^t \int_x (\partial_t IV) (IV)^2 I\Psi \, dx \, dt \quad \xleftarrow{\text{Lemma II(ii)}}$$

$$- 3 \int_{t_0}^t \int_x (\partial_t IV) \left( \underbrace{I(v : \Psi^2 : ) - (IV) I(: \Psi^2 :)}_{\text{Lemma 9}} \right) \, dx \, dt$$

$$- 3 \int_{t_0}^t \int_x (\partial_t IV) (IV) I(: \Psi^2 : ) \, dx \, dt \quad \xleftarrow{\text{Gronwall part}}$$

$$- \int_{t_0}^t \int_x (\partial_t IV) I(: \Psi^3 : ) \, dx \, dt \quad \left. \vphantom{\int_{t_0}^t \int_x (\partial_t IV) (IV) I(: \Psi^2 : )} \right\} \text{Lemma II(i)}$$

(32)

Given  $T > 0$  and  $\theta > 0$ , define

$$A(N) = \frac{\|I\Psi\|_{L_{T,x}^{\log N}}}{\log N}$$

$$\Omega_{M,\gamma,\theta} = \left\{ \max_{k=1,2} \| : \Psi^{3-k} : \|_{L_T^\infty W_x^{-\gamma(\cdot), \rho(\cdot)}} \right.$$

$$\left. + \max_{k=0,1} \| : \Psi^{3-k} : \|_{L_T^\infty W_x^{-\theta,4}} \leq M \right\}$$

↑ Gronwall part.

Prop 12:  $\exists \gamma = \gamma(s) > 0, \alpha = \alpha(s) > 0$

$$\theta = \theta(s) > 0,$$

$$\beta = \beta(s) > 0 \text{ with } \beta < \alpha \leq 1 - \beta(1-s)$$

s.t. if  $\omega \in \Omega_{M,\gamma,\theta}$ ,  $\underline{H(Iv)(t_0)} \leq N^\beta$   
 $|t_0| \leq T,$

then  $\exists I = I(M,T,s)$

$$\delta = f(s) > 0, N_0 = N_0(T,s)$$

s.t.

$$\underline{H(Iv)(t)} \leq N^\alpha$$

provided that  $|t - t_0| \leq \tau / \underline{A(N)}$

$$|t| < T$$

$$\underline{A(N)} \leq N^\delta, N \geq N_0$$

$$\begin{aligned}
 \text{Pf: } H(t) - H(t_0) &\lesssim \int_{t_0}^t N^{-(1-3(1-s))} H^2 \quad (33) \\
 &+ \sum_{k=1}^2 \int_{t_0}^t N^{-\frac{1-k(1-s)}{2} + r} H^{\frac{k+1}{2}} \| \Psi^{3-k} \|_{L_T^\infty W_x^{-y(r), p(r)}} \\
 &+ \sum_{k=0}^1 \int_{t_0}^t N^0 (1 + H^{3/4}) \| \Psi^{3-k} \|_{L_T^\infty W_x^{-0, 4}} \\
 &+ \left( 1 + (t - t_0) + \int_{t_0}^t H^{1+c\gamma} \right) \| \Psi \|_{L_{T,x}^{\gamma-1}}
 \end{aligned}$$

$$\text{choose } \gamma = (\log N)^{-1}$$

$$F(t) \geq H(t) \text{ and } N^\beta \leq F \leq N^\alpha.$$

$$\begin{aligned}
 (i) \quad &N^{-(1-3(1-s))} H^2 \\
 &\leq N^{-\alpha} F \cdot F \leq F, \quad \alpha \leq 1-3(1-s)
 \end{aligned}$$

$$\begin{aligned}
 (ii) \quad &\underline{k=2}: \quad N^{-\frac{1-2(1-s)}{2} + r} H^{\frac{1}{2}} H \\
 &\leq N^{-\frac{\alpha}{2}} F^{\frac{1}{2}} F \leq F \quad \text{choose } r \leq \frac{1-s}{2}
 \end{aligned}$$

$$\begin{aligned}
 &\underline{k=1}: \quad N^{-\frac{1-(1-s)}{2} + r} H \\
 &\leq H \leq F.
 \end{aligned}$$

$$(iii) N^\theta (1 + H^{\frac{3}{4}})$$

(34)

$$\approx N^\theta F^{\frac{3}{4}} = \underbrace{N^\theta F^{-\frac{1}{4}}}_{\approx 1} \cdot F$$

if  $\theta \leq \frac{\beta}{4}$  i.e. given  $\beta > 0$ ,  
choose  $\theta \ll 1$ .

$$\Rightarrow H(t) \leq H(t_0) + C_{M,T,S} \left( 1 + \int_0^t F^{1+c(\log N)^{-1}} \right) A \log N$$

- Let  $\beta < \tilde{\beta} < \alpha$

Consider  $F(t) = N^{\tilde{\beta} + A(t-t_0)}$

- $F(t_0) = N^{\tilde{\beta}} > N^\beta \geq H(t_0)$

- Suppose that  $\exists t < T$  s.t.  $t - t_0 < \frac{\alpha - \tilde{\beta}}{\lambda}$

and  $F(t) \leq H(t)$ )

(Otherwise, we would have

$$H(t) \leq F(t) \leq N^{\tilde{\beta} + \alpha - \tilde{\beta}} = N^\alpha, \quad \forall t - t_0 < \frac{\alpha - \tilde{\beta}}{\lambda}$$

- By continuity of  $H$  and  $F$ ,

$\exists t_*$  with  $t_* - t_0 < \frac{\alpha - \tilde{\beta}}{\lambda}$

s.t.  $H(t_*) = F(t_*)$

(35)

$$\Rightarrow N^{\tilde{\beta}} + \lambda(t_* - t_0)$$

$$\leq N^{\tilde{\beta}} + C \left( 1 + N^{\tilde{\beta}(1 + c(\log N)^{-1})} \right)$$

$$\times \int_{t_0}^{t_*} N^{\lambda(1 + c(\log N)^{-1})(t - t_0)} dt$$

$$\times A \log N$$

$$\leq N^{\tilde{\beta}} + \lambda(t_* - t_0)$$

$$\times \left\{ N^{\frac{1}{\tilde{\beta} - \beta}} N^{\lambda(t_* - t_0)} + C \frac{A \log N}{N^{\tilde{\beta} + \lambda(t_* - t_0)}} \right\}$$

$$+ C A (\cancel{\log N}) N^{\tilde{\beta} c (\log N)^{-1}} \times \frac{1}{N^{\lambda(t_* - t_0)}}$$

$$\times \int_{t_0}^{t_*} N^{\lambda(1 + c(\log N)^{-1})(t - t_0)} dt$$

$$= \frac{N^{\lambda(1 + c(\log N)^{-1})(t_* - t_0)}}{\lambda(1 + c(\log N)^{-1}) \cancel{\log N}}$$

- 1

drop

$$\Rightarrow 1 \leq \left( \frac{1}{N^{\tilde{\beta}-\beta} N^{\lambda(t_*)-t_0}} + \frac{c A \log N}{N^{\tilde{\beta}} N^{\lambda(t_*)-t_0}} \right) \quad (36)$$

$\leq \frac{1}{2} \quad \text{if } A \gtrsim N^{\tilde{\beta}}$

$$+ c A \frac{N^{\tilde{\beta}} c (\log N)^{-1} N^{\lambda c (\log N)^{-1} (t_*) - t_0}}{\lambda (1 + c (\log N)^{-1})}$$

$(N^{c (\log N)^{-1}} = e^c)$

$$= c A e^{c \tilde{\beta} + c \lambda (t_*) - t_0}$$

$$\frac{c A e^{c \lambda}}{\lambda (1 + c (\log N)^{-1})}$$

$$t_* - t_0 \leq \frac{\lambda - \tilde{\beta}}{\lambda}$$

$$\Rightarrow \frac{1}{2} \leq \frac{c A e^{c \lambda}}{\lambda (1 + c (\log N)^{-1})}$$

$$\leq c' \frac{A}{\lambda} e^{c \lambda} \Rightarrow \textcircled{X} \quad \text{by choosing } \frac{A}{\lambda} = \kappa \ll 1.$$

$$\Rightarrow H(t) \leq F(t) \leq N^\lambda, \quad \forall t - t_0 \leq \frac{\lambda - \tilde{\beta}}{\lambda}$$

$$= \frac{I}{A}.$$

□

Prop 13: (almost a.s. GWP): Let  $s > 4/5$ . (37)

Given  $T, \varepsilon > 0$  (unrelated!!),  $\exists \Omega_{T, \varepsilon} \subset \Omega$

s.t. ①  $P(\Omega_{T, \varepsilon}^c) < \varepsilon$

②  $\forall \omega \in \Omega_{T, \varepsilon}$ ,

$\exists!$  soln  $u$  to the defocusing cubic rSNLW  
on  $[0, T]$  of the form  $u = v + \Phi$

satisfying  $\|v\|_{L_T^\infty H_x^\alpha} \leq C(s, T, \varepsilon)$ .

Rmk: a.a.s GWP  $\Rightarrow$  a.s. GWP (i.e. GWP as an SPDE)  
Borel-Cantelli

Pf: Fix  $N_0 = N_0(s)$

$$N_{k+1} = N_k^\sigma, \quad \sigma > 1$$

s.t.

$$\textcircled{+} \quad N_{k+1}^{2(1-s)} N_k^\alpha + N_k^{2\alpha} \ll N_{k+1}^\beta$$

$$\Rightarrow \text{Need } \beta > 2(1-s)$$

$$2(1-s) < \beta < \alpha \leq 1 - 3(1-s)$$

$$\Rightarrow s > \underline{\underline{4/5}}$$

Suppose  $H(I_{N_k} v)(t) \leq N_k^\alpha$

(38)

Then,

$$H(I_{N_{k+1}} v)(t) \lesssim \|I_{N_{k+1}} \vec{v}\|_{H^2}^2 + \|I_{N_{k+1}} v\|_{L^4}^4$$

$$\vec{v} = (v, \partial_t v)$$

$$\lesssim N_{k+1}^{2(1-s)} \| \vec{v} \|_{H^s}^2 + \| I_{N_{k+1}} v \|_{H^{\frac{1}{2}}}^4$$

$$( \| f \|_{H^0} \lesssim \| f \|_{H^1} ) \underbrace{\quad}_{\leq \| v \|_{H^{\frac{1}{2}}}^4} \lesssim \| I_{N_k} v \|_{H^{\frac{1}{2}}}^4$$

$$\lesssim N_{k+1}^{2(1-s)} H(I_{N_k} v) + H^2(I_{N_k} v)$$

$$\lesssim N_{k+1}^{2(1-s)} N_k^\alpha + N_k^{2d} \stackrel{+}{<} N_{k+1}^\beta$$

(+)

$$\Rightarrow H(I_{N_{k+1}} v)(t) \leq N_{k+1}^\beta$$

$$\text{Let } \Omega_\Lambda(N) = \{A(N) \subseteq \Lambda\}$$

$\Rightarrow$  By Lemma 10,

$$P(\Omega_\Lambda(N)^c) \leq \frac{C \log N}{N^{\log N}} T^{\frac{\log N}{2} + 1} \underset{N \rightarrow \infty}{\sim} N^{\log C_{T,s} - \log \Lambda}$$

$$\text{Define } \Omega_\Lambda = \bigcap_{k \in \mathbb{Z}_{\geq 0}} \Omega_\Lambda(N_k)$$

by choosing  
 $\Lambda \gg 1$

$$\Rightarrow P(\Omega_\Lambda^c) \leq \sum_k N_k^{\log C_{T,s} - \log \Lambda} \underset{k \rightarrow \infty}{\sim} N_0^{\log C_{T,s} - \log \Lambda}$$

$\rightarrow 0$  as  $\Lambda \rightarrow \infty$ .

choose  $M(T, \varepsilon), \lambda(T, \varepsilon) \gg 1$  s.t.

(39)

$$P(\Omega_{\lambda}^c) + P(\Omega_{M, \varepsilon, \theta}^c) < \varepsilon$$

Choose  $k_0 \gg 1$  s.t.

$$\lambda < N_{k_0}^{\alpha}$$

$$H(I_{N_{k_0}} v)(0) \leq N_{k_0}^{\beta}.$$

Prop 12

$$H(I_{N_{k_0}} v)(t) \leq N_{k_0}^{\alpha}, \quad \forall t < \frac{T}{A(N_{k_0})}$$

$$\text{With } t = \frac{T}{\lambda}.$$

(Recall  $A(N_k) \leq \lambda$   
on  $\Omega_\lambda$ )

⊕

$$\Rightarrow H(I_{N_{k_0+1}} v)\left(\frac{T}{\lambda}\right) \leq N_{k_0+1}^{\beta}.$$

Prop 12

$$\Rightarrow H(I_{N_{k_0+1}} v)\left(2 \frac{T}{\lambda}\right) \leq N_{k_0+1}^{\alpha}$$

⊕

$$\Rightarrow H(I_{N_{k_0+2}} v)\left(2 \frac{T}{\lambda}\right) \leq N_{k_0+2}^{\beta}$$

Prop 12

...

Iterate the argument  $\lceil \frac{\lambda T}{\varepsilon} \rceil + 1$  times.

(Note: increment is indep of  $N$ .)

$\Rightarrow$  Given  $t \in \left[ \frac{(j-1)\tau}{\Lambda}, \frac{j\tau}{\Lambda} \right]$ , we have (40)

$$\begin{aligned} \|\vec{v}(t)\|_{\mathcal{H}^s} &\lesssim H(I_{N_{k_0+j-1}} v)^{\frac{1}{k}}(t) \\ &\leq N_{k_0+j-1}^{d/2} \quad \left( \leq N_{k_0+\left[\frac{\Lambda T}{\tau}\right]-1}^{d/2} \right) \end{aligned}$$

$k_0, d, \Lambda, \tau$  depend only on  $s, T$ , and  $\varepsilon$ .



Rmk: A standard application of the I-method:

Given  $T \gg 1$ , choose  $N = N(T) \gg 1$ .

but in our argument, we needed to choose

an increasing seq  $N_{k_0+j}$ ,  $j = 0, 1, \dots, \left[\frac{\Lambda T}{\tau}\right]$

$\Leftarrow$  much more subtle.

J. Faraco on BBM '18.

- GWP of the defocusing cubic SNLW on  $\mathbb{R}^2$   
L. Tolomeo '18.

## Chap 2: quadratic SNLW in 3-d

(41)

$$(\partial_t^2 + 1 - \Delta) u + u^2 = \tilde{\gamma} \quad \text{on } \mathbb{R}_+ \times \mathbb{T}^3.$$
$$(u, \partial_t u)|_{t=0} = (u_0, u_1)$$

$d=3 \Rightarrow \tilde{\gamma}$  has spatial regularity  $-\frac{3}{2} - \varepsilon$ .

### • Parabolic case:

$$(\text{SNLH}) \quad (\partial_t + 1 - \Delta) u + u^2 = \tilde{\gamma}$$

Da Prato - Debussche:  $u = v + \varphi$

$$\varphi = \mathcal{I}(\tilde{\gamma})$$

vertex  $\bullet = \tilde{\gamma}$

edge = Duhamel integral operator  $\mathcal{I}$ .  
 $= (\partial_t + 1 - \Delta)^{-1}$

• The heat Duhamel integral operator  $\mathcal{I}$  gains  
2 derivatives (with  $dt'$ ) but  
with  $dW(t')$ , only 1 derivative  
( $\Leftarrow$  2 derivatives in the second moment.)

$\Rightarrow \varphi \sim -\frac{1}{2} -$  We only keep track of  
spatial regularities.

Rules

- A product of functions of reg.  $s_1$  and  $s_2$  is defined if  $s_1 + s_2 > 0$ . When  $s_1 > 0$  and  $s_1 \geq s_2$ , the resulting prod has regularity  $s_2$ .
- A product of stochastic objects (not depending on the unknown) is always well defined, possibly with a renormalization. The product of stoch. obj. of reg  $s_1$  and  $s_2$  has regularity

$$\min(s_1, s_2, s_1 + s_2)$$

$$(SNLH) \xrightarrow{D-D} (2t + 1 - \Delta) V = -(V + \varphi)^2$$

$\stackrel{""}{=} -V^2 - 2V\varphi - \varphi$

renorm  
 $\varphi^2 \rightsquigarrow V$

$\uparrow$   
 $-1-$

$$\Rightarrow \text{expect } V \sim 1- = (-1-) + 2.$$

$$V \quad \varphi$$

$1- \quad -\frac{1}{2}-$

$$(1-) + \left(-\frac{1}{2}-\right) > 0$$

$\Rightarrow$  The product  $V\varphi$  and hence all the terms on RHS are well defined.

$\Rightarrow$  run a contraction argument in  $C_T C_x^{1-}$

(43)

Back to the wave case:

$$\varphi = I(\xi) \sim -\frac{1}{2} -$$

$$\text{Here, } I = (\partial_t^2 + 1 - \Delta)^{-1}$$

(corresp to the forward fund. soln.)

$$\text{Wick power: } V_N = \varphi_N^2 - \sigma_N \text{ with } \varphi_N = P_{\leq N} \varphi$$

$$\cdot \sigma_N = \mathbb{E}[\varphi^2(t, x)] \sim tN.$$

$$\Rightarrow V_N \rightarrow V \text{ in } C_T W_x^{-1, \infty} \text{ a.s.}$$

2nd order stoch. obj:

$$Y = I(V) = \int_0^t \frac{\sin(t-t')\langle \nabla \rangle}{\langle \nabla \rangle} V(t') dt'$$

A naive "parabolic thinking" yields

$$0^- = 2\left(-\frac{1}{2} -\right) + 1.$$

$\Leftarrow$  BAD. We need to use the explicit product structure and exhibit multilinear dispersive smoothing

(44)

Prop 14:  $Y_N = I(V_N) \rightarrow Y$

in  $C_T W_x^{\frac{1}{2}-\varepsilon, \infty} \cap C_T^1 W_x^{-1-\varepsilon, \infty}$ , a.s.

i.e.  $Y \sim \frac{1}{2} -$   $\Leftarrow$  extra  $\frac{1}{2}$ -smoothing

Second order expansion:  $\Downarrow$  "at least as smooth as  $Y$ "

$$U = \vartheta - Y + V.$$

$$\begin{aligned} (\text{SNLW}) \Rightarrow (2t^2 + 1 - \Delta)V &= - (V + \vartheta - Y)^2 + V \\ &= -(V - Y)^2 - 2V\vartheta + \underbrace{2\vartheta Y}_{-\frac{1}{2} -} \end{aligned}$$

$\Rightarrow$  expect  $V \sim \frac{1}{2} -$

$$\Rightarrow V\vartheta : \left(\frac{1}{2} -\right) + \left(-\frac{1}{2} -\right) < 0$$

The product does not make sense.

So, the second order expansion is NOT enough.

- Littlewood-Paley projection:  $P_j$  onto  $\{|m| \sim 2^j\}$ .

$$f = \sum_{j=0}^{\infty} P_j f.$$

- Paraproduct decomposition (Bony '81).

$$fg = f \circledcirc g + f \circledast g + f \circledgt g$$

$$= \sum_{j < k-2} P_j f P_k g + \sum_{|j-k| \leq 2} P_j f P_k g + \sum_{k < j-2} P_j f P_k g.$$

- $f \circledcirc g =$  paraproduct of  $g$  by  $f$ . ( $g = \text{high freq}$ )

always well defined as a distribution  
of regularity  $\min(S_2, S_1 + S_2)$

- $f \circledast g =$  resonant product

In general, well defined only if  $S_1 + S_2 > 0$ .

$\Leftarrow$  The resonant prod is the source of difficulty  
in making sense of a nonlinearity.

- Set  $\circledast = \circledcirc + \circledast$ .

(46)

Further decomposition:

Paracontrolled ansatz:  $v = X + Y$

$$s_1, s_2, \quad 0 < s_1 < s_2$$

$$(\partial_t^2 + 1 - \Delta) X = -2(X + Y - Y) \circledleftarrow 1$$

(SNLW2)

$$(\partial_t^2 + 1 - \Delta) Y = -(X + Y - Y)^2 - 2(X + Y - Y) \circledrightarrow 1$$

• Why called a paracontrolled ansatz?

A distribution  $f$  is said to be paracontrolled  
(by a given reference distribution  $g$ ) if  $\exists f'$  s.t.

$$f = f' \circledleftarrow g + h$$

$\Leftarrow$  smoother

(SNLW2) says  $\square v = \square X + \square Y$  is paracontrolled.

Expect  $X \sim \frac{1}{2} - = (-\frac{1}{2}) + 1$ .

For now, ignore  $\circledrightarrow$  in the  $Y$ -eqn.

$$(X + Y - Y) \circledleftarrow 1 \sim 0 -$$

$$\Rightarrow Y \sim 1 -$$

$\Rightarrow Y \circledrightarrow 1$  makes sense as long as  $s_2 > \frac{1}{2}$ .

$$\mathbb{Y} = Y \odot \mathbf{1}$$

$$0 = (\frac{1}{2} -) + (-\frac{1}{2} -)$$

without renormalization

We did not expect any further renorm.  
since it was not needed for SNLH.

In order to prove this, we need to exploit dispersion at a multilinear level.  
(Otherwise, it looks as if there is a log divergence.)

Prop 15:  $\mathbb{Y}_N = Y_N \odot \mathbf{1}_N \rightarrow \mathbb{Y}$  in  $C_T W_x^{-\varepsilon, \alpha}$  a.s.

•  $X \odot \mathbf{1}$ ?

$(\frac{1}{2} -) + (-\frac{1}{2} -) < 0 \Rightarrow$  does not make sense  
as it is.

Idea: Use the structure of  $X$ , i.e.

$$\begin{aligned} X(t) &= \underbrace{\partial_t S(t) X_0 + S(t) X_1}_{= \vec{S}(t)(X_0, X_1)} - 2 \mathcal{I}((X + Y - Y) \odot \mathbf{1}) \end{aligned}$$

Lem 16:  $s_1 > 0$ .  $(X_0, X_1) \in \mathcal{H}^{s_1}(\mathbb{T}^3)$ .

$$Z_N = (\vec{S}(t)(X_0, X_1)) \ominus I_N$$



$$Z = (\vec{S}(t)(X_0, X_1)) \ominus I. \quad \leftarrow \text{Need to include } Z \text{ in an enhanced data set.}$$

in  $C_T H_x^{s_1 - \frac{1}{2} - \varepsilon}$ , a.s.

A straightforward computation with the Wiener chaos estimate and a direct application of Kolmogorov's continuity criterion.

Note: Set of prob 1 depends on the initial data  $(X_0, X_1)$ .

In the parabolic setting, at this point, one would introduce (smoother) commutators:

$$\begin{aligned} & [I((X+Y-Y) \odot I)] \ominus I \\ &= [(X+Y-Y) \odot I(I)] \ominus I + \text{com}_1 \ominus I \quad \text{BAD} \\ &= (X+Y-Y) \odot \underbrace{\{ I(I) \ominus I \}}_{\text{explicitly known}} + \text{com}_1 \ominus I + \text{com}_2 \end{aligned}$$

*stoch obj. of reg 0 -*

(49)

Bad News:  $\text{com}_1$  is NOT smooth in the dispersive setting!!

Main idea: directly study the following paracontrolled operator (and its res product with  $\mathbb{I}$ ).

Given  $w \in C(\mathbb{R}_+; H^{s_1}(\mathbb{T}^3))$  with  $0 < s_1 < \frac{1}{2}$ ,

define  $\boxed{g_{\odot}(w)(t) = \mathbb{I}(w \odot \mathbb{I})(t)}$

$$= \sum_{j < k-2} \mathbb{I}(P_j w \cdot P_k \mathbb{I})$$

$$= \sum_{n \in \mathbb{Z}^3} e_n \sum_{n = n_1 + n_2} \int_0^t \frac{\sin((t-t')\langle n \rangle)}{\langle n \rangle} \widehat{w}(t', m_1) \widehat{\mathbb{I}}(t', m_2) dt'$$

$|n_1| < |m_2|$

signifies the paraproduct  $\odot$ .

Goal: Make sense of  $g_{\odot}(w) \odot \mathbb{I}$

with  $w = X + Y - Y$ .

- Divide  $g_{\odot}$  into good and bad parts

Fix  $\theta > 0$  small

$\mathcal{F}_{\odot}^{(1)} = \text{restriction of } g_{\odot} \text{ onto } \{|m_1| \geq |m_2|^{\theta}\}$

$\mathcal{F}_{\odot}^{(2)} = g_{\odot} - \mathcal{F}_{\odot}^{(1)}$ .

Namely,

$$g_{\leftarrow}^{(1)}(w)(t) = \sum_{n \in \mathbb{Z}^3} e_m \sum_{n = n_1 + n_2} \int_0^t \frac{\sin((t-t')x_n)}{\langle m \rangle} \hat{w}(t', m_1) \hat{i}(t', m_2) dt'.$$

$|n_2|^\theta \lesssim |m_1| \ll |m_2|$ .

point:  $|m_1|$  is NOT too small.  $\rightarrow |m_1| \sim |m_2|$

$$\frac{\langle m \rangle^{\frac{1}{2}+2\varepsilon}}{\langle m \rangle} \lesssim \langle m_1 \rangle^{\frac{4\varepsilon}{\theta}} \langle m_2 \rangle^{-\frac{1}{2}-2\varepsilon}$$

$$\lesssim \langle m_1 \rangle^{s_1-\varepsilon} \langle m_2 \rangle^{-\frac{1}{2}-2\varepsilon}$$

by choosing  $\varepsilon = \varepsilon(s_1, \theta) > 0$  suff. small.

Lem 17:  $0 < s_1 < \frac{1}{2}$ . Given small  $\theta > 0$ ,  
 $\exists$  small  $\varepsilon = \varepsilon(s_1, \theta) > 0$  s.t. given any

$$B \in C(\mathbb{R}_+; W^{\frac{1}{2}-\varepsilon, \infty}(\mathbb{T}^3)),$$

the paracontrolled operator

$$g_{\leftarrow}^{(1), B} = g_{\leftarrow}^{(1)} \text{ with } i \text{ replaced by } B$$

belongs to  $\mathcal{L}_2 = \mathcal{L}\left(C_T H_x^{s_1}; C_T H_x^{\frac{1}{2}+2\varepsilon}\right)$



following the notation from GKO'18.

- When  $|m_1| \ll |m_2|^{\theta}$ , the positive neg of  $w$  does not help.

We directly study  $\mathcal{J}_{\Theta, \Xi}(w)(t)$ .

$$\mathcal{J}_{\Theta, \Xi}(w)(t) = \mathcal{J}_{\Theta}^{(2)}(w) \Xi(t)$$

$$= \sum_{n \in \mathbb{Z}^3} e_n \int_0^t \sum_{n_1, n_2 \in \mathbb{Z}^3} \hat{w}(t', m_1) A_{n, m_1}(t, t') dt'.$$

$$A_{n, m_1}(t, t') = \mathbf{1}_{[0, t]}(t') \sum_{n - n_1 = n_2 + n_3} \frac{\sin((t-t')\langle m_1, m_2 \rangle)}{\langle m_1 + m_2 \rangle} \hat{g}(t', m_2) \hat{g}(t, m_3)$$

$$|n_1| \ll |m_2|^{\theta}$$

$(m_1 + m_2) \sim |n_3|$  — resonant product

$$= \mathbf{1}_{[0, t]}(t) \sum_{n - n_1 = n_2 + n_3} \frac{\sin((t-t')\langle m_1 + m_2 \rangle)}{\langle m_1 + m_2 \rangle} (\hat{g}(t', m_2) \hat{g}(t, m_3) - \mathbf{1}_{m_2 + n_3 = 0} \Gamma_{n_2}(t, t'))$$

$$+ \mathbf{1}_{[0, t]}(t) \cdot \mathbf{1}_{m = n_1} \sum_{n \in \mathbb{Z}^3} \frac{\sin((t-t')\langle m + m_2 \rangle)}{\langle m + m_2 \rangle} \Gamma_2(t, t')$$

$$|n| \ll |m_2|^{\theta}$$

$$=: A_{m, m_1}^{(1)}(t, t') + A_{n, n_2}^{(2)}(t, t')$$

deterministic counter term

More difficult!!

Here,  $0 \leq t_2 \leq t_1$ ,

$$\Gamma_m(t_1, t_2) = \mathbb{E} [\hat{g}(t_1, m) \hat{g}(t_2, m)]$$

$$= \frac{\cos((t_1 - t_2)\langle m \rangle)}{2 \langle m \rangle^2} t_2 + \mathcal{O}\left(\frac{1}{\langle m \rangle^3}\right).$$

(52)

By exploiting dispersion (stationary phase) and symmetrization ( $m_2 \leftrightarrow -n_2$ )

i.e. the order of summation matters  
 $\Rightarrow$  only conditionally convergent.

Prop 18:  $s_2 < 1$ .  $\exists$  small  $\theta(s_2) > 0$  and  $\varepsilon > 0$  s.t.

$f_{\Theta, \Xi}$  belongs to the class

$$L_1 = L \left( C_T L_x^2 \cap C_T^1 H_x^{-\varepsilon}; C_T H_x^{s_2-1} \right).$$

Moreover,  $f_{\Theta, \Xi}^N$  (replacing  $\vartheta$  by  $\vartheta_N$ )

converges a.s. to  $f_{\Theta, \Xi}$  in  $L_1$ .

Final form:

(SNLW3)

$$\begin{aligned} (\partial_t^2 + I - \Delta) X &= -2(X + Y - Y^\circ) \otimes \vartheta \\ (\partial_t^2 + I - \Delta) Y &= -(X + Y - Y^\circ)^2 - 2(X + Y - Y^\circ) \otimes \vartheta \\ &\quad - 2Y \otimes \vartheta + 2Y^\circ - 2Z \\ &\quad + 4 f_{\Theta}^{(II)} (X + Y - Y^\circ) \otimes \vartheta \end{aligned}$$

$$+ 4 f_{\Theta, \Xi} (X + Y - Y^\circ)$$

$$(X, \partial_t X, Y, \partial_t Y) \Big|_{t=0} = (X_0, X_1, Y_0, Y_1).$$

Thm 19:  $\frac{1}{4} < s_1 < \frac{1}{2} < s_2 \leq s_1 + \frac{1}{4}$ .

$\exists \theta = \theta(s_2) > 0$  and  $\varepsilon = \varepsilon(s_1, s_2, \theta) > 0$  s.t.

if (i)  $\gamma, Y, \chi$  in  $C(\mathbb{R}_+; W_x^{s, \infty})$

$$s = -\frac{1}{2}-, \frac{1}{2}-, 0-$$

and

$$\gamma \in C^1(\mathbb{R}_+; W_x^{-1-\varepsilon})$$

$$(ii) \quad Z \in C(\mathbb{R}_+; H_x^{s, -\frac{1}{2}-\varepsilon}).$$

$$(iii) \quad f_{\otimes, \odot} \in L_1,$$

then the system (SNLW3) is locally well-posed.

More precisely, given  $(X_0, X_1, Y_0, Y_1) \in \mathcal{F}^{s_1} \times \mathcal{N}^{s_2}$ ,

$\exists T$  and unique soln  $(X, Y)$  in the class

$$Z_T^{s_1, s_2} = X_T^{s_1} \times Y_T^{s_2}$$

$$\subset C([0, T]; H^{s_1} \times H^{s_2}) \cap C^1([0, T]; H^{s_1-1} \times H^{s_2-2})$$

depending continuously on the enhanced data set.

$$\boxed{\square} = (X_0, X_1, Y_0, Y_1, \gamma, Y, \chi, Z, f_{\otimes, \odot})$$

in the class

$$\begin{aligned} \chi_T^{s_1, s_2, \varepsilon} &= \mathcal{F}^{s_1} \times \mathcal{N}^{s_2} \times C_T W_x^{-\frac{1}{2}-\varepsilon, \infty} \times (C_T W_x^{\frac{1}{2}-\varepsilon} \cap C_T^1 W_x^{+\varepsilon, \infty}) \\ &\times C_T W_x^{-\varepsilon, \infty} \times C_T H^{s, -\frac{1}{2}-\varepsilon} \times L_1 \end{aligned}$$

$$\bullet \quad u_N = P_N - Y_N + X_N + Y_N$$

↓      ↓      ↓  
 $u = P - Y + X + Y \text{ in } C_T H_x^{-\frac{1}{2} - \epsilon}$

Pf of Thm 19 : Duhamel formulation.

$$X(t) = \Phi_1(X, Y)(t)$$

$$= \vec{S}(t)(X_0, X_1) - 2 \int_0^t \vec{S}(t-t') [(X+Y-Y) \odot P](t') dt'.$$

$$Y(t) = \Phi_2(X, Y)(t)$$

$$= \vec{S}(t)(Y_0, Y_1) - \int_0^t \vec{S}(t-t') [ \dots ](t') dt'.$$

Strichartz estimates:  $0 \leq s \leq 1$ .

We say a pair  $(q, r)$  is  $s$ -admissible

$(\tilde{q}, \tilde{r})$  is dual  $s$ -admis.

(i.e.  $(\tilde{q}', \tilde{r}')$  is  $(1-s)$ -admis.)

if  $1 \leq \tilde{q}' < 2 < q \leq \infty$ ,  $1 < \tilde{r}' \leq 2 \leq r < \infty$ ,

$$\frac{1}{q} + \frac{d}{r} = \frac{d}{2} - s = \frac{1}{\tilde{q}'} + \frac{d-1}{\tilde{r}'} - z \quad \leftarrow \text{scaling condition}$$

$$\left. \begin{aligned} \frac{1}{q} + \frac{d-1}{2\tilde{r}'} &\leq \frac{d-1}{4} \\ \frac{2}{q} + \frac{d-1}{2\tilde{r}'} &\geq \frac{d+3}{4} \end{aligned} \right\} \text{admissibility cond.}$$

(55)

Then, a soln  $u$  to

$$\begin{cases} (\partial_t^2 + 1 - \Delta) u = f \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \end{cases}$$

satisfies

$$\|(\bar{u}, \partial_t \bar{u})\|_{C_T H_x^s} + \|u\|_{L_T^8 L_x^r}$$

$$\lesssim \|(\bar{u}_0, \bar{u}_1)\|_{H^s} + \|f\|_{L_T^{\frac{8}{3}} L_x^{\frac{8}{3}} + L_x^1 H_x^{s-1}}$$

for  $0 \leq T \leq 1$ .

$(q, r)$ ,  $s$ -admis

$(\tilde{q}, \tilde{r})$ , dual  $s$ -admis.

$\leftarrow$  follows from the Strichartz estimates on  $\mathbb{R}^d$  and finite speed of propagation.

• Back to the proof of Thm 19: Define the Strichartz space

$$X_T^{s_1} = C_T H_x^{s_1} \cap C_T^1 H_x^{s_1-1} \cap L_T^\delta W_x^{s_1 - \frac{4}{3}, \frac{8}{3}} \quad (\delta, \frac{8}{3}), \frac{1}{4}\text{-admis}$$

$$Y_T^{s_2} = C_T H_x^{s_2} \cap C_T^1 H_x^{s_2-1} \cap L_T^4 W_x^{s_2 - \frac{1}{2}, 4} \quad (4, 4), \frac{1}{2}\text{-admis}$$

$$Z_T^{s_1, s_2} = X_T^{s_1} \times Y_T^{s_2}$$

(56)

$$\cdot \| \Phi_1(X, Y) \|_{Y_T^{s_1}} \lesssim \| (X_0, X_1) \|_{N^{s_1}} + \| (X+Y-Y) \otimes I \|_{L_T^1 H_x^{s_1-1}}$$

(brace under the previous term)

$$\lesssim T \| X+Y-Y \|_{L_T^\infty L_x^2} \| I \|_{L_T^\infty W_x^{-\frac{1}{2}-\varepsilon, \infty}}$$

$$\lesssim \| (X_0, X_1) \|_{N^{s_1}} + T (1 + \| (X, Y) \|_{Z_T^{s_1, s_2}})$$

if  $s_1-1 < -\frac{1}{2}-\varepsilon$ .

Y-equation:

$$-2(X+Y-Y) \otimes I - 2Y \otimes I + 2 \circlearrowleft Y + Z$$

$$+ \oint_{\mathbb{S}}^{\prime\prime} (X+Y-Y) \otimes I$$

$$\Rightarrow \| \dots \|_{Y_T^{s_1}} \lesssim \| (Y_0, Y_1) \|_{N^{s_2}} + T (1 + \| (X, Y) \|_{Z_T^{s_1, s_2}})$$

If  $\frac{1}{2} < s_2 < \min(1, s_1 + \frac{1}{2})$

• Also

$$\| \int_0^t \delta(t-t') \oint_{\mathbb{S}} (X+Y-Y) dt' \|_{Y_T^{s_2}}$$

$$\lesssim \| \oint_{\mathbb{S}} (X+Y-Y) \|_{L_T^1 H_x^{s_2-1}}$$

$$\lesssim T \| X+Y-Y \|_{L_T^\infty L_x^2 \cap C_T^1 H_x^{-1-\varepsilon}}$$

$$\lesssim T (1 + \| (X, Y) \|_{Z_T^{s_1, s_2}}).$$

$$\left\| \int_0^t S(t-t') (X+Y-Y)^2(t') dt' \right\|_{Y_T^{S_2}}$$

strichartz

$$\lesssim \left\| \langle \nabla \rangle^{S_2 - \frac{1}{2}} (X+Y-Y)^2 \right\|_{L_{T,x}^{4/3}}$$

frac. leib.

$$\lesssim T^{\frac{1}{4}} \left( \left\| \langle \nabla \rangle^{S_2 - \frac{1}{2}} X \right\|_{L_T^{\infty} L_x^{S_2}}^2 + \left\| \langle \nabla \rangle^{S_2 - \frac{1}{2}} Y \right\|_{L_T^4 L_x^4}^2 \right)$$

Need  
 $\leq S_1 - \frac{1}{4}$

$$+ \left\| \langle \nabla \rangle^{S_2 - \frac{1}{2}} Y \right\|_{L_{T,x}^{\infty}}^2 \Big)$$

$$\lesssim T^{\frac{1}{4}} \left( 1 + \left\| (X, Y) \right\|_{\sum_T^{S_1, S_2}}^2 \right).$$

$$\text{if } S_2 \leq \min(1-\varepsilon, S_1 + \frac{1}{4}).$$

$$\Rightarrow \left\| \Phi(X, Y) \right\|_{\sum_T^{S_1, S_2}} \lesssim \left\| (X_0, X_1, Y_0, Y_1) \right\|_{N^{S_1} \times N^{S_2}} \\ + T^\theta \left( 1 + \left\| (X, Y) \right\|_{\sum_T^{S_1, S_2}}^2 \right).$$

for some  $\theta > 0$ .

Similarly,

$$\left\| \Phi(X, Y) - \Phi(\tilde{X}, \tilde{Y}) \right\|_{\sum_T^{S_1, S_2}}$$

$$\lesssim T^\theta \left( 1 + \left\| (X, Y) \right\| + \left\| (\tilde{X}, \tilde{Y}) \right\| \right) \left\| (X, Y) - (\tilde{X}, \tilde{Y}) \right\|_{\sum_T^{S_1, S_2}}$$

 $\Rightarrow$  By taking  $T > 0$  suff. small, $\Phi$  is a contraction on a ball  $B_R \subset \sum_T^{S_1, S_2}$ 

□

## Key point of the proof of Prop 14 on $\hat{Y}$

In using Lemma 2, we bound

$$\mathbb{E}[\hat{Y}(t, n)^2]$$

$$= 4 \sum_{\substack{n = n_1 + n_2 \\ n_1 \neq \pm n_2}} \int_0^t \frac{\sin(t - t_i) \langle n \rangle}{\langle n \rangle} \int_0^{t_i} \frac{\sin(t - t_2) \langle n \rangle}{\langle n \rangle} \Gamma_{n_1}(t_1, t_2) \times \Gamma_{n_2}(t_1, t_2) dt_2 dt_1$$

$$+ \dots \quad n_1 = n_2$$

$$\Gamma_n(t_1, t_2) = \frac{\cos(t_1 - t_2) \langle n \rangle}{2 \langle n \rangle^2} t_2 + O(\frac{1}{\langle n \rangle^3})$$

$\Rightarrow$  expand sines and cosines in complex exponentials

$$\sum_{\substack{n = n_1 + n_2 \\ n_1 \neq \pm n_2}} \frac{e^{i(\varepsilon_1 + \varepsilon_2)t \langle n \rangle}}{\langle n \rangle^2 \langle n_1 \rangle^2 \langle n_2 \rangle^2} \times \underbrace{\int_0^t e^{-it_1 K_1(\bar{n})} \int_0^{t_1} t_2 e^{it_2 K_2(\bar{n})} dt_2 dt_1}_{K_1(\bar{n}) = \varepsilon_1 \langle n \rangle - \varepsilon_3 \langle n_1 \rangle - \varepsilon_4 \langle n_2 \rangle}$$

$$K_2(\bar{n}) = \varepsilon_2 \langle n \rangle + \varepsilon_3 \langle n_1 \rangle + \varepsilon_4 \langle n_2 \rangle.$$

integrate in  $t_1$  first.

$$\left| \int_0^t t_2 e^{-it_2 K_2(\bar{n})} \frac{e^{-it_1 K_1(\bar{n})} - e^{-it_1 K_1(\bar{n})}}{-i K_1(\bar{n})} dt_2 \right| \leq \frac{C(T)}{(1 + K_1(\bar{n}))}$$

(59)


Need to bound

$$I = \sum_{n=n_1+n_2} \frac{1}{\langle m \rangle^2 \langle m_1 \rangle^2 \langle m_2 \rangle^2 (1 + |K_1(\bar{m})|)} \quad \left( \text{WTS } \lesssim \langle m \rangle^{-4+} \right)$$

BAD case  $(\varepsilon_1, \varepsilon_3, \varepsilon_4) = (\pm 1, \pm 1, \mp 1)$ .

Assume:  
 $|m_1| \gg 1$   
 $|n_1| \geq |n_2|$

$$|K_1(\bar{m})| = \langle m \rangle + \langle m_2 \rangle - \langle m_1 \rangle$$

Note:  $\langle m_1 \rangle \sim \langle m \rangle + \langle m_2 \rangle$ 

$$\begin{aligned} \text{law of cosines: } & |m|^2 + |m_2|^2 - |m_1|^2 \\ &= 2|m||m_2| \cos(\angle(m, m_2)). \end{aligned}$$

$$\begin{aligned} \Rightarrow |K_1(\bar{m})| &= \frac{(\langle m \rangle + \langle m_2 \rangle)^2 - \langle m_1 \rangle^2}{\langle m \rangle + \langle m_2 \rangle + \langle m_1 \rangle} \\ &= \frac{2\langle m \rangle \langle m_2 \rangle + |m|^2 + |m_2|^2 - |m_1|^2 + 1}{\langle m \rangle + \langle m_2 \rangle + \langle m_1 \rangle} \end{aligned}$$

$$\gtrsim \frac{|m| |m_2| (1 - \cos \theta)}{\langle m_1 \rangle} \quad \theta = \angle(m_2, -m)$$

Case 1 :  $1 - \cos \theta \gtrsim 1$ . (large angle)

(60)  
non-resonant

$$I \lesssim \sum_{n=n_1+n_2} \frac{1}{\langle m \rangle^3 \langle m_1 \rangle^3 \langle m_2 \rangle^3} \quad \langle m_1 \rangle \sim \max(\langle m \rangle, \langle m_2 \rangle)$$


$$\lesssim \langle m \rangle^{-4+}$$

Case 2 :  $1 - \cos \theta \ll 1$  (close to collinear) nearly resonant

$$\Rightarrow 0 \leq \theta \ll 1.$$

$$\Rightarrow 1 - \cos \theta \sim \theta^2 \ll 1.$$

Dyadically decompose  $|m_2| \sim N_2$ ,  $N_2 \geq 1$ , dyadic



$m_2$  is constrained to a cone  $C$   
 $|m_2| \sim N_2$  height  $\sim N_2 \sin \theta \sim N_2$   
base disc of radius  
 $\sim N_2 \sin \theta \sim N_2 \theta$ .

$$\Rightarrow \text{vol}(C) \sim N_2^3 \theta^2$$

$$\Rightarrow I \leq \sum_{N_2 \geq 1} \frac{1}{\langle m \rangle^3 \max(\langle m \rangle, N_2) N_2^3 \theta^2}$$

dyadic

$$\lesssim \langle m \rangle^{-4+}$$



(61)

• Prop 15 on  $\gamma = \gamma_{\oplus 1}$

- The proof proceeds analogously to that of Prop 14 but a bit more complicated

New difficult term.

$\boxed{\gamma > 0}$   
small

$$\sum_{\substack{n_2 \in \mathbb{Z}^3 \\ |n + n_2| \approx |n_2| \\ |n| \ll |n_2|^{\gamma}}} \frac{\sin(t - t')(\langle n + n_2 \rangle - \langle n_2 \rangle)}{\langle n \rangle^2 \langle n + n_2 \rangle \langle n_2 \rangle^2} \left( \begin{array}{l} \text{WTS} \\ \lesssim \langle n \rangle^{-3} \end{array} \right)$$

This condition allows us to rewrite the sum as

$$\sum_{\substack{n_2 \in \mathbb{Z}^3 \\ |n| \ll |n_2|^{\gamma}}} \quad \leftarrow \text{we can use symmetrization} \quad n_2 \leftrightarrow -n_2$$

$$\begin{aligned} \text{Let } \Theta^{\pm}(n, n_2) &= \langle n \pm n_2 \rangle - \langle n_2 \rangle \mp \frac{\langle n, n_2 \rangle}{\langle n_2 \rangle} \\ &= \Theta\left(\frac{\langle n \rangle^2}{\langle n_2 \rangle}\right) \end{aligned}$$

$$\begin{aligned} \text{sum} &= \sum_{\substack{n_2 \in \mathbb{Z}^3/2 \\ |n| \ll |n_2|^{\gamma}}} \frac{1}{\langle n \rangle^2 \langle n + n_2 \times n_2 \rangle^2} \\ &\quad \times \left[ \sin(t - t_i) \left( \frac{\langle n, n_2 \rangle}{\langle n_2 \rangle} + \Theta^+(n, n_2) \right) \right. \end{aligned}$$

$$\left. - \sin(t - t') \left( \frac{\langle n, n_2 \rangle}{\langle n_2 \rangle} - \Theta^-(n, n_2) \right) \right]$$

$$\begin{aligned} \text{MVT} &\lesssim \sum_{|n| \ll |n_2|^{\gamma}} \frac{1}{\langle n \rangle^2 \langle n_2 \rangle^3} \left( \frac{\langle n \rangle^2}{\langle n_2 \rangle} \right)^{\delta} \quad \text{for any } \delta \in [0, 1]. \end{aligned}$$

$$\lesssim \langle n \rangle^{-3+} \quad \square$$

On Prop 18 for  $\mathcal{J} \otimes \mathbb{E}$

(62)

The difficult part  $A^{(2)}$

- symmetrization:  $m_2 \leftrightarrow -m_2$
- integration by parts in time  
to handle  
 $\sin(t-t')(\langle m+m_2 \rangle + \langle m_2 \rangle)$ .