

施  
惟  
樞

## Differential Topology

Lectures by John Milnor, Princeton University, Fall term 1958

Notes by James Munkres

Differential topology may be defined as the study of those properties of differentiable manifolds which are invariant under diffeomorphism (differentiable homeomorphism).

Typical problems falling under this heading are the following:

- (1) Given two differentiable manifolds, under what conditions are they diffeomorphic?
- (2) Given a differentiable manifold, is it the boundary of some differentiable manifold-with-boundary?
- (3) Given a differentiable manifold, is it parallelizable?

All of these problems concern more than the topology of the manifold, yet they do not belong to differential geometry, which usually assumes additional structure (e.g., a connection or a metric).

The most powerful tools in this subject have been derived from the methods of algebraic topology. In particular, the theory of characteristic classes is crucial, whereby one passes from the manifold  $M$  to its tangent bundle, and thence to a cohomology class in  $M$  which depends on this bundle.

These notes are intended as an introduction to the subject; we will go as far as possible without bringing in algebraic topology. Our two main goals are Whitney's theorem that a differentiable  $n$ -manifold can be imbedded as a closed

subset of the euclidean space  $\mathbb{R}^{2n+1}$  (see §1.32); and Thom's theorem that the non-orientable cobordism group  $\pi_n$  is isomorphic to a certain stable homotopy group (see §3.15).

Chapter I is mainly concerned with approximation theorems. First the basic definitions are given and the inverse function theorem is exploited. (§1.1 - 1.12). Next two local approximation theorems are proved, showing that a given map can be approximated by one of maximal rank. (§1.13 - 1.21). Finally locally finite coverings are used to derive the corresponding global theorems: namely Whitney's imbedding theorem and Thom's transversality lemma (§1.35).

Chapter II is an introduction to the theory of vector space bundles, with emphasis on the tangent bundle of a manifold. Chapter III makes use of the preceding material in order to study the cobordism groups  $\pi_n$ .

Chapter I Imbeddings and Immersions of Manifolds.

Notation. If  $x$  is in the euclidean space  $R^n$ , the coordinates of  $x$  are denoted by  $(x^1, \dots, x^n)$ . Let  $\|x\| = \max |x^i|$ ; let  $C^n(r)$  denote the set of  $x$  such that  $\|x\| < r$ ; and  $C^n(x_0, r)$  the set of  $x$  such that  $\|x - x_0\| < r$ . The closure of a cube  $C$  is denoted by  $\bar{C}$ .

A real valued function  $f(x^1, \dots, x^n)$  is differentiable if the partials of  $f$  of all orders exist and are continuous (i.e., "differentiable" means  $C^\infty$ ). A map  $f: U \rightarrow R^p$  (where  $U$  is an open set, in  $R^n$ ) is differentiable if each of the coordinate functions  $f^1, \dots, f^p$  is differentiable.  $Df$  denotes the Jacobian matrix of  $f$ ; one verifies that  $D(gf) = Dg \cdot Df$ . The notation  $\partial(f^1, \dots, f^p) / \partial(x^1, \dots, x^n)$  is also used. If  $n = p$ ,  $|Df|$  denotes the determinant.

1.1. Definition. An  $n$ -manifold  $M^n$  is a Hausdorff space with a countable basis which is locally homeomorphic to  $R^n$ .

A differentiable structure  $\mathcal{D}$  on a manifold  $M^n$  is a collection of real-valued functions, each defined on an open subset of  $M$ , such that:

- 1) For every point  $p$  of  $M$  there is a neighborhood  $U$  of  $p$  and a homeomorphism  $h$  of  $U$  onto an open subset of  $R^n$  such that a function  $f$ , defined on the open subset  $W$  of  $U$ , is in  $\mathcal{D}$  if and only if  $fh^{-1}$  is differentiable.

2) If  $U_i$  are open sets contained in the domains of  $f$  and  $U = \cup U_i$ , then  $f|_U \in \mathcal{D}$  if and only if  $f|_{U_i}$  is in  $\mathcal{D}$ , for each  $i$ .

A differentiable manifold  $M^n$  is a manifold provided with a differentiable structure  $\mathcal{D}$ ; the elements of  $\mathcal{D}$  are called the differentiable functions on  $M$ . Any open set  $U$  and homeomorphism  $h$  which satisfy the requirements of 1) above are called a coordinate system on  $M$ . Notation. A coordinate system is sometimes denoted by the coordinate functions:  $h(p) = (u^1(p), \dots, u^n(p))$ .

1.2. Alternate definition. Let a collection  $(U_i, h_i)$  be given, where  $h_i$  is a homeomorphism of the open subset  $U_i$  of  $M^n$  onto an open subset of  $R^n$ , such that

- the  $U_i$  cover  $M$
- $h_j \circ h_i^{-1}$  is a differentiable map on  $h_i(U_i \cap U_j)$ , for all  $i, j$ .

Define a coordinate system as an open set  $U$  and homeomorphism  $h$  of  $U$  onto an open subset of  $R^n$  such that  $h|_U$  and  $h|_U^{-1}$  are differentiable on  $h(U \cap U_i)$  and  $h_i(U \cap U_i)$  respectively, for each  $i$ . Define a differentiable structure on  $M$  as the collection of all such coordinate systems. A function  $f$ , defined on the open set  $V$ , is differentiable if  $f \circ h^{-1}$  is differentiable on  $h(U \cap V)$ , for all coordinate systems  $(U, h)$ .

One shows readily that these two definitions are entirely equivalent.

1.3. Definition. Let  $M_1, M_2$  be differentiable manifolds. If  $U$  is an open subset of  $M_1$ ,  $f: U \rightarrow M_2$  is differentiable if for every differentiable function  $g$  on  $M_2$ ,  $gf$  is differentiable on  $M_1$ .

If  $A \subset M_1$ , a function  $f: A \rightarrow M_2$  is differentiable if it can be extended to a differentiable function defined on a neighborhood  $U$  of  $A$ .

$f: M_1 \rightarrow M_2$  is a diffeomorphism if  $f$  and  $f^{-1}$  are defined and differentiable.

(A coordinate system  $(U, h)$  on  $M^n$  is then an open set  $U$  in  $M$  and a diffeomorphism  $h$  of  $U$  onto an open set in  $R^n$ .)

If  $A \subset M$ , we have just defined the notion of differentiable function for subsets of  $A$ . Suppose that  $A$  is locally diffeomorphic to  $R^k$ : this collection is easily shown to be a differentiable structure on  $A$ . In this case,  $A$  is said to be a differentiable submanifold of  $M$ .

The following lemma is familiar from elementary calculus.

1.4. Lemma. Let  $f: C^n(r) \rightarrow R^n$  satisfy the condition

$$\left| \frac{\partial f^i}{\partial x^j} \right| \leq b, \text{ for all } i, j. \text{ Then } \|f(x) - f(y)\| \leq bn \|x - y\|,$$

for all  $x, y \in \bar{C}^n$ .

1.5. Theorem (inverse function theorem). Let  $U$  be an open subset of  $\mathbb{R}^n$ , let  $f: U \rightarrow \mathbb{R}^n$  be differentiable, and let  $Df$  be non-singular at  $x_0$ . Then  $f$  is a diffeomorphism of some neighborhood of  $x_0$  onto some neighborhood of  $f(x_0)$ .

Proof: We may assume  $x_0 = f(x_0) = 0$ , and that  $Df(x_0)$  is the identity matrix.

Let  $g(x) = f(x) - x$ , so that  $Dg(0)$  is the zero matrix. Choose  $r > 0$  so that  $x \in U$  and  $Df(x)$  is non-singular and  $|\partial g^i / \partial x_j| \leq 1/2n$ , for all  $x$  with  $\|x\| < r$ .

Assertion. If  $y \in C(r/2)$ , there is exactly one  $x \in C(r)$  such that  $f(x) = y$ :

(\*) By the previous lemma,  $\|g(x) - g(x_0)\| \leq 1/2 \|x - x_0\|$  on  $C(r)$ . Let us define  $x_0 = 0$ ,  $x_1 = y$ ,  $x_{n+1} = y - g(x_n)$ . This is defined, since  $x_n - x_{n-1} = g(x_{n-2}) - g(x_{n-1})$ , so that  $\|x_n - x_{n-1}\| \leq 1/2 \|x_{n-2} - x_{n-1}\|$ ; and thus  $\|x_n\| \leq 2\|y\|$  for each  $n$ . Hence the sequence  $x_n$  converges to a point  $x$  with  $\|x\| \leq 2\|y\|$ , so that  $x \in C(r)$ . Then  $x = y - g(x)$ , so that  $f(x) = y$ . This proves the existence of  $x$ . To show uniqueness, note that if  $f(x) = f(x_1) = y$ , then  $g(x_1) - g(x) = x - x_1$ , contradicting (\*). \*

Hence  $f^{-1}: C(r/2) \rightarrow C(r)$  exists. Note that  $\|f(x) - f(x_1)\| \geq \|x - x_1\| - \|g(x) - g(x_1)\| \geq 1/2 \|x - x_1\|$ , so that  $\|y - y_1\| \geq 1/2 \|f^{-1}(y) - f^{-1}(y_1)\|$ . Hence  $f^{-1}$  is continuous; the image of  $C(r/2)$  under  $f^{-1}$  is open because it equals  $C(r) \cap f^{-1}(C(r/2))$ , the intersection of two open sets. \*

To show that  $f^{-1}$  is differentiable, note that  $f(x) = f(x_1) + Df(x_1) \cdot (x - x_1) + h(x, x_1)$ , where  $(x - x_1)$  is written as a column matrix and the dot stands for matrix multiplication. Here  $h(x, x_1) / \|x - x_1\| \rightarrow 0$  as  $x \rightarrow x_1$ .

Let  $A$  be the inverse matrix of  $Df(x_1)$ . Then

$$A \cdot (f(x) - f(x_1)) = (x - x_1) + A \cdot h(x, x_1), \text{ or}$$

$$A \cdot (y - y_1) + A \cdot h_1(y, y_1) = f^{-1}(y) - f^{-1}(y_1),$$

where  $h_1(y, y_1) = -h(f^{-1}(y), f^{-1}(y_1))$ . Now

$$\frac{h_1(y, y_1)}{\|y - y_1\|} = - \frac{h(x, x_1)}{\|x - x_1\|} \frac{\|x - x_1\|}{\|y - y_1\|}.$$

Since  $\|x - x_1\| / \|y - y_1\| \leq 2$ ,  $h_1(y, y_1) / \|y - y_1\| \rightarrow 0$  as  $y \rightarrow y_1$ .

Hence  $D(f^{-1}) = A = (D(f))^{-1}$ .

This means that  $D(f^{-1})$  is obtained as the composition of the following maps:

$$C(r/2) \xrightarrow{f^{-1}} C(r) \xrightarrow{Df} GL(n) \xrightarrow{\text{matrix inversion}} GL(n);$$

where  $GL(n)$  denotes the set of non-singular  $n \times n$  matrices, considered as a subspace of  $n^2$ -dimensional euclidean space.

Since  $f^{-1}$  is continuous and  $Df$  and matrix inversion are  $C^\infty$ ,  $D(f^{-1})$  is continuous, i.e.,  $f^{-1}$  is  $C^1$ . In general, if  $f^{-1}$  is  $C^k$ , then by this argument  $D(f^{-1})$  is also, i.e.,  $f^{-1}$  is of class  $C^{k+1}$ . This completes the proof.

1.6. Lemma. Let  $U$  be an open subset of  $R^n$ , let  $f: U \rightarrow R^p$  ( $n \leq p$ ),  $f(0) = 0$ , and let  $Df(0)$  have rank  $n$ . Then there exists a diffeomorphism  $g$  of one neighborhood of the origin in  $R^p$  onto another so that  $g(0) = 0$  and  $gf(x^1, \dots, x^n) = (x^1, \dots, x^n, 0, \dots, 0)$ , in some neighborhood of the origin.

Proof: Since  $\partial(f^1, \dots, f^p)/\partial(x^1, \dots, x^n)$  has rank  $n$ , we may assume that  $\partial(f^1, \dots, f^n)/\partial(x^1, \dots, x^n)$  is the submatrix which is non-singular. Define  $F: U \times R^{p-n} \rightarrow R^p$  by the equation

$$F(x^1, \dots, x^p) = f(x^1, \dots, x^n) + (0, \dots, 0, x^{n+1}, \dots, x^p).$$

$F$  is an extension of  $f$ , since  $F(x^1, \dots, x^n, 0, \dots, 0) = f(x^1, \dots, x^n)$ .

$DF$  is non-singular at the origin, since its determinant everywhere equals  $|\partial(f^1, \dots, f^n)/\partial(x^1, \dots, x^n)|$ . Hence  $F$  has a local inverse  $g$ , so that  $g$  maps one neighborhood of the origin in  $R^p$  onto another, and

$$gF(x^1, \dots, x^p) = (x^1, \dots, x^p).$$

Hence

$$gf(x^1, \dots, x^n) = (x^1, \dots, x^n, 0, \dots, 0).$$



1.7. Corollary. Let  $A^k$  be a differentiable submanifold of  $M^n$ . Given  $x \in A$ , there is a coordinate system  $(U, h)$  on  $M$  about  $x$ , such that  $h(U \cap A) = h(U) \cap R^k$  (where  $R^k$  is considered as the subspace  $R^k \times 0$  of  $R^k \times R^{n-k} = R^n$ ).

Proof: Let  $(U_1, h_1)$  be a coordinate system on  $M$  about  $x$ ; by hypothesis, there is a differentiable map  $f$  of a neighborhood  $V$  of  $x$  in  $M$  into  $R^k$  such that  $f|_{V \cap A} = f_1$  is a diffeomorphism whose range is an open set  $W$  in  $R^k$ . We may assume  $U_1 = V$ , and  $h_1(x) = f(x) = 0$ .

Now  $f h_1^{-1} h_1 f_1^{-1}$  is the identity on  $W$ , so that its Jacobian, which equals  $D(fh_1^{-1}) \cdot D(h_1 f_1^{-1})$  is non-singular. Hence  $D(h_1 f_1^{-1})$  has rank  $k$ , so that by the previous lemma, there is a diffeomorphism  $g$  of some neighborhood  $V_1 \subset h_1(U_1)$  of  $0$  onto another such that  $g(0) = 0$  and  $g h_1 f_1^{-1}(x^1, \dots, x^k) = (x^1, \dots, x^k, 0, \dots, 0)$ .

Then  $U = h_1^{-1}(V_1)$  and  $h = g h_1$  will satisfy the requirements of the lemma.

1.8. Lemma. Let  $U$  be an open subset of  $R^n$ , let  $f: U \rightarrow R^p$ ,  $f(0) = 0$ , ( $n \geq p$ ), and let  $Df(0)$  have rank  $p$ . Then there is a diffeomorphism  $h$  of some neighborhood of the origin in  $R^n$  onto another such that  $h(0) = 0$  and  $fh(x^1, \dots, x^n) = (x^1, \dots, x^p)$ .

Proof: We may assume  $\frac{\partial(f^1, \dots, f^p)}{\partial(x^1, \dots, x^p)}$  is non-

singular at 0, since  $Df(0)$  has rank  $p$ . Define

$F: U \rightarrow \mathbb{R}^n$  by the equation

$F(x^1, \dots, x^n) = (f^1(x), \dots, f^p(x), x^{p+1}, \dots, x^n)$ . Then  $DF(0)$

is non-singular; let  $h$  be the local inverse of  $F$ . Let  $g$

project  $\mathbb{R}^n$  onto the subspace  $\mathbb{R}^p$ ;  $f = gF$ . Then

$fh(x^1, \dots, x^n) = gF h(x^1, \dots, x^n) = g(x^1, \dots, x^n) = (x^1, \dots, x^p)$ .

1.9. Exercise. Let  $U$  be an open subset of  $\mathbb{R}^n$ ,  $f: U \rightarrow \mathbb{R}^p$ ,  $f(0) = 0$ ; and let  $Df(x)$  have rank  $k$  for all  $x$  in  $U$ . Then there are local diffeomorphisms  $h$  and  $g$  of  $\mathbb{R}^n$  and  $\mathbb{R}^p$  respectively such that

$$g f h(x^1, \dots, x^n) = (x^1, \dots, x^k, 0, \dots, 0).$$

1.10. Definition. If  $f: M_1 \rightarrow M_2$ , the rank of  $f$  at  $x$  is the rank of  $D(h_2 f h_1^{-1})$  at  $h_1(x)$ , where  $(U_1, h_1)$  and  $(U_2, h_2)$  are coordinate systems about  $x$  and  $f(x)$ , respectively. The differentiable map  $f: M^n \rightarrow M^p$  is an immersion if  $\text{rank } f = n$  everywhere ( $n \leq p$ ). It is an imbedding if it is also a homeomorphism into.

If  $f: M^n \rightarrow M^p$ , then  $y \in M^p$  is a regular value of  $f$  if  $\text{rank } f = p$  on the entire set  $f^{-1}(y)$ . Otherwise,  $y$  is a critical value. (If  $y \notin f(M^n)$ ,  $y$  is thus a regular value of  $f$ .)

1.11. Exercise. If  $A$  is a differentiable submanifold of  $M$ , the inclusion  $A \rightarrow M$  is an imbedding and conversely if  $f: M_1 \rightarrow M$  is an imbedding then  $f(M_1)$  is a differentiable submanifold.

1.12. Exercise. If  $y$  is a regular value of  $f: M^n \rightarrow M^p$ , then  $f^{-1}(y)$  is a differentiable submanifold of  $M^n$  of dimension  $n - p$  (or is empty).

1.13. Definition. A subset  $A$  of  $R^n$  has measure zero if it may be covered by a countable collection of cubes  $C(x,r)$  having arbitrarily small total volume. In such a case,  $R^n - A$  is everywhere dense (i.e., it intersects every open set).

1.14. Lemma. Let  $U$  be an open subset of  $R^n$ ; let  $f: U \rightarrow R^n$  be differentiable. If  $A \subset U$  has measure 0, so does  $f(A)$ .

Proof: Let  $C$  be any cube with  $\bar{C} \subset U$ . Let  $b$  denote the maximum of  $|\partial f^i / \partial x^j|$  on  $\bar{C}$  for all  $i, j$ . By 1.4,  $\|f(x) - f(y)\| \leq b n \|x - y\|$  for  $x, y \in \bar{C}$ .

Now  $A \cap C$  has measure zero; let us cover  $A \cap C$  by cubes  $C(x_i, r_i)$  contained in  $C$ , such that  $\sum_{i=1}^{\infty} r_i^n < \epsilon$ .

Then  $f(C(x_i, r_i)) \subset C(f(x_i), b n r_i)$ , so that  $f(A \cap C)$  is covered by cubes of total volume  $b^n n^n \sum r_i^n < b^n n^n \epsilon$ .

Hence  $f(A \cap C)$  has measure zero.

Since  $A$  can be covered by countably many such cubes  $C$ ,  $f(A)$  has measure zero.

1.15. Corollary. If  $f: U \rightarrow \mathbb{R}^p$  is differentiable, where  $U$  is an open subset of  $\mathbb{R}^n$  and  $n < p$ , then  $f(U)$  has measure 0.

Proof: Project  $U \times \mathbb{R}^{p-n}$  onto  $U$  and apply  $f$ . Since  $U \times 0$  has measure 0 in  $\mathbb{R}^p$ , so does  $f(U)$ .

1.16. Definition. If  $A \subset M$ ,  $A$  has measure 0 if  $h(A \cap U)$  has measure 0 for every coordinate system  $(U, h)$ .

1.17. Corollary. If  $f: M^n \rightarrow M^p$  is differentiable and  $n < p$ , then  $f(M^n)$  has measure zero.

1.18. Definition. Let  $M(p, n)$  denote the space of  $p \times n$  matrices, with the differentiable structure of the euclidean space  $\mathbb{R}^{pn}$ . Let  $M(p, n; k)$  denote the subspace consisting of matrices of rank  $k$ . Thus  $M(p, n; n)$  is an open subset of  $M(p, n)$  if  $p \geq n$ ; the determinantal criterion for rank proves this. More generally, we have:

1.19. Lemma.  $M(p, n; k)$  is a differentiable submanifold of  $M(p, n)$  of dimension  $k(p+n-k)$ , where  $k \leq \min(p, n)$ .

Proof: Let  $E_0 \in M(p, n; k)$ ; we may assume that  $E_0$  is of the form  $\begin{pmatrix} A_0 & B_0 \\ C_0 & D_0 \end{pmatrix}$ , where  $A_0$  is a non-singular

$k \times k$  matrix. There is an  $\varepsilon > 0$  such that if all the entries of  $A - A_0$  are less than  $\varepsilon$ ,  $A$  must also be non-singular. Let  $U$  consist of all matrices in  $M(p, n)$  of the form  $E = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ , with all the entries of  $A - A_0$  less than  $\varepsilon$ .

Then  $E$  is in  $M(p, n; k)$  if and only if  $D = CA^{-1}B$ ;

For the matrix

$$\begin{pmatrix} I_k & 0 \\ X & I_{p-k} \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} A & B \\ XA + C & XB + D \end{pmatrix}$$

has the same rank as  $E$ . If  $X = -CA^{-1}$ , this matrix is

$$\begin{pmatrix} A & B \\ 0 & -CA^{-1}B + D \end{pmatrix}. \quad \text{If } D = CA^{-1}B, \text{ this matrix has rank}$$

$k$ . The converse also holds, for if any element of  $-CA^{-1}B + D$  is different from zero, this matrix has rank  $> k$ .

Let  $W$  be the open set in euclidean space of dimension  $(pn - (p-k)(n-k)) = k(p+n-k)$  consisting of matrices

$$\begin{pmatrix} A & B \\ C & 0 \end{pmatrix}, \quad \text{with all the entries of } A - A_0 \text{ less than } \varepsilon.$$

The map  $\begin{pmatrix} A & B \\ C & 0 \end{pmatrix} \rightarrow \begin{pmatrix} A & B \\ C & CA^{-1}B \end{pmatrix}$  is then a diffeomorphism

of  $W$  onto the neighborhood  $U \cap M(p, n; k)$  of  $E_0$ .

1.20. Theorem. Let  $U$  be an open set in  $\mathbb{R}^n$ , and let  $f: U \rightarrow \mathbb{R}^p$  be differentiable, where  $p \geq 2n$ . Given  $\varepsilon > 0$ , there is a  $p \times n$  matrix  $A = (a_j^i)$  with each  $|a_j^i| < \varepsilon$ , such that  $g(x) = f(x) + A \cdot x$  is an immersion. ( $x$  written as a column matrix.)

Proof:  $Dg(x) = Df(x) + A$ ; we would like to choose  $A$  in such a way that  $Dg(x)$  has rank  $n$  for all  $x$ . I.e.,  $A$  should be of the form  $Q - Df$ , where  $Q$  has rank  $n$ .

We define  $F_k: M(p, n; k) \times U \rightarrow M(p, n)$  by the equation

$$F_k(Q, x) = Q - Df(x).$$

Now  $F_k$  is a differentiable map, and the domain of  $F_k$  has dimension  $k(p+n-k) + n$ . As long as  $k < n$ , this expression is monotonic in  $k$  (its partial with respect to  $k$  is  $p + n - 2k$ ). Hence the domain of  $F_k$  has dimension not greater than  $(n-1)(p+n-(n-1)) + n = (2n-p) + pn - 1$  for  $k < n$ . Since  $p \geq 2n$ , this dimension is strictly less than  $pn = \dim M(p, n)$ .

Hence the image of  $F_k$  has measure zero in  $M(p, n)$ , so that there is an element  $A$  of  $M(p, n)$ , arbitrarily close to the zero matrix, which is not in the image of  $F_k$  for  $k = 0, \dots, n-1$ . Then  $A + Df(x) = Dg(x)$  has rank  $n$ , for each  $x$ .

1.21. Theorem. Let  $U$  be an open subset of  $\mathbb{R}^n$ ; and let  $f: U \rightarrow \mathbb{R}^p$  be differentiable. Given  $\varepsilon > 0$ , there are matrices  $A$  ( $p \times n$ ) and  $B$  ( $p \times 1$ ) with entries less than  $\varepsilon$  in absolute value, such that

$$g(x) = f(x) + A \cdot x + B$$

has the origin as a regular value.

Remark. The following much more delicate result has been proved by A. Sard: The set of critical values of any differentiable map has measure zero.

Proof of 1.21. Note that the theorem is trivial if  $p > n$ , since then  $f(U)$  has measure zero, and we may choose  $A = 0$  and  $B$  small in such a way that  $0$  is not in the image of  $g$ .

Assume  $p \leq n$ . We wish  $Dg(x_0) = Df(x_0) + A$  to have rank  $p$ , where  $x_0$  ranges over all points such that

$$g(x_0) = 0 = f(x_0) + A \cdot x_0 + B.$$

Hence  $A$  is of the form  $Q - Df(x)$ , and  $B$  is of the form  $-f(x) - A \cdot x$ , where  $Q$  is to have rank  $p$ .

We define  $F_k: M(p, n; k) \times U \rightarrow M(p, n) \times \mathbb{R}^p$  by the equation

$$F_k(Q, x) = (Q - Df(x), -f(x) - (Q - Df(x)) \cdot x).$$

Then  $F_k$  is differentiable. If  $k < p$ , the dimension of

its domain is not greater than  $(p-1)(p+n-(p-1)) + n = p + pn - 1$ . Hence the image of  $F_k$ ,  $k = 0, \dots, p-1$  has measure zero; so that there is a point  $(A, B)$  arbitrarily close to the origin which is not in any such image set. This completes the proof.

1.22. Definition. A covering of  $X$  is locally-finite if every point has a neighborhood which intersects only finitely many elements of the covering. A refinement of a covering of  $X$  is a second covering each element of which is contained in an element of the first covering. A Hausdorff space is paracompact if every open covering has a locally-finite open refinement.

If  $X$  is paracompact, and  $U_\alpha$  is an open covering, there is a locally-finite open covering  $V_\alpha$  with  $V_\alpha \subset U_\alpha$  for each  $\alpha$ . For let  $W_\beta$  be a locally-finite refinement of  $U_\alpha$ ; choose  $\alpha(\beta)$  so that  $W_\beta \subset U_{\alpha(\beta)}$  for each  $\beta$ . Set  $V_{\alpha_0} = U_{\alpha(\beta)=\alpha_0} W_\beta$ . Given a neighborhood intersecting only finitely many  $W_\beta$ , it intersects only finitely many  $V_\alpha$  as well.

1.23. Theorem. If  $X$  is locally compact and Hausdorff, having a countable basis,  $X$  is paracompact.

Proof: Let  $U_1, U_2, \dots$  be a basis for  $X$  with  $\bar{U}_i$  compact for each  $i$ . There exists a sequence  $A_1, A_2, \dots$  of compact sets whose union is  $X$ , such that  $A_i \subset \text{Int } A_{i+1}$ :



Set  $A_1 = \bar{U}_1$ . Given  $A_i$  compact, let  $k$  be the smallest integer such that  $A_i$  is contained in  $U_1 \cup \dots \cup U_k$ ; let  $A_{i+1}$  equal the closure of this set union  $\bar{U}_{i+1}$ .

Let  $O$  be an open covering of  $X$ . Cover the compact set  $A_{i+1} - \text{Int } A_i$  by a finite number of open sets  $V_1, \dots, V_n$  where each  $V_i$  is contained in some element of  $O$ , and in the open set  $\text{Int } A_{i+2} - A_{i-1}$ . Let  $P_i$  denote the collection  $\{V_1, \dots, V_n\}$ , and let  $P = P_0 \cup P_1 \cup \dots$ .  $P$  refines  $O$ , and since any compact closed neighborhood  $C$  is contained in some  $A_i$ ,  $C$  can intersect only finitely many elements of  $P$ .

1.24. Exercise. Prove: A paracompact space is normal. (First prove that it is regular.)

1.25. Theorem. Let  $M^n$  be a differentiable manifold,  $\{U_\alpha\}$  an open covering of  $M$ . There is a collection  $(V_j, h_j)$  of coordinate systems on  $M$  such that

- 1)  $\{V_j\}$  is a locally-finite refinement of  $\{U_\alpha\}$ .
- 2)  $h_j(V_j) = C^n(3)$
- 3) If  $W_j = h_j^{-1}(C(1))$ , then  $\{W_j\}$  covers  $M$ .

$C^n(1) *$

Proof: The proof proceeds along lines similar to the previous one. The only difference is that one chooses the  $V_j$  to satisfy 2), and makes sure that the sets  $h_j^{-1}(C(1))$  also cover  $A_{i+1} - \text{Int } A_i$ .

1.26. We wish to construct a  $C^\infty$  function  $\varphi(x_1, \dots, x_n)$  such that  $\varphi = 1$  on  $\bar{C}(1)$ ,  $0 < \varphi < 1$  on  $C(2) - \bar{C}(1)$ ,  $\varphi = 0$  on  $R^n - C(2)$ .

This function may be defined by the equation

$$\varphi(x_1, \dots, x_n) = \prod_1^n \psi(x_i), \text{ where}$$

$$\psi(x) = \frac{\lambda(2+x) \cdot \lambda(2-x)}{\lambda(2+x) \cdot \lambda(2-x) + \lambda(x-1) + \lambda(-x-1)}$$

and

$$\begin{aligned} \lambda(x) &= e^{-1/x} & \text{if } x > 0 \\ &= 0 & \text{if } x \leq 0. \end{aligned}$$

Note that the denominator in the expression for  $\psi$  is always positive, and that  $\psi(x) = 1$  for  $|x| \leq 1$

$$0 < \psi(x) < 1 \text{ if } 1 < |x| < 2$$

$$\psi(x) = 0 \text{ if } |x| \geq 2.$$

1.27. Definition. Let  $f, g: X \rightarrow Y$ , where  $Y$  is metrizable, and let  $\delta(x)$  be a positive continuous function defined on  $X$ . Then  $g$  is a  $\delta$ -approximation to  $f$  if  $d(f(x), g(x)) < \delta(x)$  for all  $x$ . [If one takes the  $\delta$ -approximations to  $f$  to be a neighborhood of  $f$  in the function space  $F(X, Y)$ , this imposes a topology on the function space, independent of the metric on  $Y$  ( $X, Y$  paracompact).]

1.28. Theorem. Given a differentiable map  $f: M^n \rightarrow R^p$  where  $p \geq 2n$ , and a continuous positive function  $\delta$  on  $M^n$ , there exists an immersion  $g: M^n \rightarrow R^p$

which is a  $\delta$ -approximation to  $f$ . If  $\text{rank } f = n$  on the closed set  $N$ , we may choose  $g|_{N=f|N}$ .

Proof: Rank  $f = n$  on a neighborhood  $U$  of  $N$ . Cover  $M^n$  by  $U$  and  $M^n - N$ . Let  $(V_i, h_i)$  be a refinement of this covering, constructed as in 1.25. As before,  $h_i(W_i) = C(1)$  and  $h_i(V_i) = C(3)$ . Let  $h_i(U_i) = C(2)$ . Let the  $V_i$  be so indexed with positive and negative integers that those  $V_i$  with non-positive indices are the ones contained in  $U$ . Let  $\varepsilon_i = \min$  of  $\delta(x)$  on the compact set  $\bar{U}_i$ .

Set  $f_0 = f$ . Given  $f_{k-1}: M^n \rightarrow R^p$ , having rank  $n$  on  $N_{k-1} = \bigcup_{j < k} \bar{W}_j$ , consider  $f_{k-1} h_k^{-1}: C(3) \rightarrow R^p$ . Let  $A$  be a  $p \times n$  matrix; let  $F_A: C(3) \rightarrow R^p$  be defined by the equation

$$F_A(x) = f_{k-1} h_k^{-1}(x) + \varphi(x) A \cdot (x),$$

where  $(x)$  is written (as usual) as a column matrix ( $n \times 1$ );  $A$  is yet to be chosen; and  $\varphi(x)$  is the function defined in 1.26.

First, we want  $F_A(x)$  to have rank  $n$  on the set  $K = h_k(N_{k-1} \cap \bar{U}_k)$ ; we are given that  $f_{k-1} h_k^{-1}$  has rank  $n$  on  $K$ . Now

$$D(F_A(x)) = D(f_{k-1} h_k^{-1}(x)) + A \cdot (x) \cdot D\varphi(x) + \varphi(x)A.$$

( $D\varphi$  is a  $1 \times n$  matrix.) The map of  $K \times M(p, n)$  into

$M(p,n)$  which carries  $(x,A)$  into  $D(F_A(x))$  is continuous. It carries  $K \times (0)$  into the open subset  $M(p,n;n)$  of  $M(p,n)$ . Hence if  $A$  is sufficiently small, this map will carry  $K \times A$  into  $M(p,n;n)$ ; our first requirement is that  $A$  be this small.

Secondly, we require  $A$  to be small enough that  $\|A \cdot (x)\| < \varepsilon_k / 2^k$  for all  $x \in C(3)$ .

Finally, by 1.20,  $A$  may be chosen arbitrarily small so that  $f_{k-1} h_k^{-1}(x) + A \cdot (x)$  has rank  $n$  on  $C(2)$ . Let  $A$  be chosen to satisfy this requirement.

We then define  $f_k: M^n \rightarrow R^D$  by the equation:

$$\begin{aligned} f_k(y) &= f_{k-1}(y) + \varphi(h_k(y)) A \cdot (h_k(y)) \quad \text{for } y \in V_k \\ &= f_{k-1}(y) \quad \text{for } y \in M - \bar{U}_k. \end{aligned}$$

These definitions agree on the overlapping domains, so that  $f_k$  is differentiable. By the first condition on  $A$ , it has rank  $n$  on  $N_{k-1}$ ; by the third condition it has rank  $n$  on  $\bar{W}_k$ . By the second condition,  $f_k$  is a  $\delta/2^k$  approximation to  $f_{k-1}$ .

We define  $g(x) = \lim_{k \rightarrow \infty} f_k(x)$ . Since the covering  $V_i$  is locally finite, all the  $f_k$  agree on a given compact set for  $k$  sufficiently large; it follows that  $g$  is differentiable and has rank  $n$  everywhere. It is also a  $\delta$ -approximation to  $f$ .

1.29. Lemma. If  $p > 2n$ , any immersion  $f: M^n \rightarrow R^p$  can be  $\delta$ -approximated by a 1-1 immersion  $g$ . If  $f$  is 1-1 in a neighborhood  $U$  of the closed set  $N$ , we may choose  $g|_{N=f|N}$ .

Proof: Choose a covering  $\{U_\alpha\}$  of  $M$  such that  $f|_{U_\alpha}$  is an imbedding (possible by 1.6). Let  $(V_1, h_1)$  be the locally finite refinement constructed in 1.25; let  $\varphi(x)$  be the function constructed in 1.26. Let  $\varphi_1(y) = \varphi(h_1(y))$  for  $y \in V_1$ ;  $= 0$  for other  $y$ . Then  $\varphi_1$  is differentiable. As before, we assume  $(V_1, h_1)$  refines the covering  $(U, M-N)$  and that those  $V_i$  with non-positive indices are the ones contained in  $U$ . Let  $f_0 = f$ . Given the immersion  $f_{k-1}: M^n \rightarrow R^p$ , we define  $f_k$  by the equation

$$f_k(y) = f_{k-1}(y) + \varphi_k(y)b_k, \text{ where } b_k \text{ is}$$

a point of  $R^p$  yet to be chosen. By the argument of the previous theorem, if  $b_k$  is chosen sufficiently small,  $f_k$  will have rank  $n$  everywhere. The first requirement is that  $b_k$  be this small; the second requirement is that  $b_k$  be small enough that  $f_k$  be a  $\delta/2^k$  approximation to  $f_{k-1}$ .

Finally, let  $N^{2n}$  be the open subset of  $M^n \times M^n$  consisting of pairs  $(y, y_0)$ , with  $\varphi_k(y) \neq \varphi_k(y_0)$ . Consider the differentiable map of  $N^{2n}$  into  $R^p$  which carries  $(y, y_0)$  into  $-(f_{k-1}(y) - f_{k-1}(y_0))/[\varphi_k(y) - \varphi_k(y_0)]$ . Since  $2n < p$ , the image of  $N^{2n}$  has measure 0, so that  $b_k$  may be chosen arbitrarily small and not in this image.

It follows that  $f_k(y) - f_k(y_0) = 0$  if and only if  $\varphi_k(y) - \varphi_k(y_0) = 0$  and  $f_{k-1}(y) - f_{k-1}(y_0) = 0$  ( $k > 0$ ).

Define  $g(y) = \lim_{k \rightarrow \infty} f_k(y)$ . If  $g(y) = g(y_0)$  and  $y \neq y_0$ , it would follow that  $f_{k-1}(y) = f_{k-1}(y_0)$  and  $\varphi_k(y) = \varphi_k(y_0)$  for all  $k > 0$ . The former condition implies that  $f(y) = f(y_0)$ , so that  $y$  and  $y_0$  cannot belong to any one set  $U_i$ . Because of the latter condition, this means that neither is in any set  $U_i$  for  $i > 0$ . Hence, they lie in  $U$ , contradicting the fact that  $f$  is 1-1 on  $U$ .

1.30. Definition. Let  $f: M^n \rightarrow R^p$ . The limit set  $L(f)$  is the set of  $y \in R^p$  such that  $y = \lim f(x_n)$  for some sequence  $\{x_1, x_2, \dots\}$  which has no limit point on  $M^n$ .

Exercise. Show the following:

1)  $f(M)$  is a closed subset of  $R^p$  if and only if

$$L(f) \subset f(M)$$

2)  $f$  is a topological imbedding if and only if  $f$  is 1-1 and  $L(f) \cap f(M)$  is vacuous.

1.31. Lemma. There exists a differentiable map  $f: M^n \rightarrow R$  with  $L(f)$  empty.

Proof: Let  $(V_i, h_i)$  and  $\varphi$  be chosen as in 1.25 and 1.26 with  $i$  ranging over positive integers; let  $\varphi_i(y) = \varphi h_i(y)$  if  $y \in V_i$ ;  $= 0$  otherwise.

Define  $f(y) = \sum_j (j \varphi_j(y))$ . This sum is finite, since  $V_1$  is a locally finite covering. If  $\{x_i\}$  is a set of points of  $M$  having no limit point, only finitely many lie in any compact subset of  $M$ . Given  $m$ , there is an integer  $i$  such that  $x_i$  is not in  $\bar{W}_1 \cup \dots \cup \bar{W}_m$ . Hence  $x_i \in \bar{W}_j$  for some  $j > m$ , whence  $f(x_i) > m$ . Thus the sequence  $f(x_m)$  cannot converge.

1.32. Corollary. Every  $M^n$  can be differentiably imbedded in  $R^{2n+1}$  as a closed subset.

Proof: Let  $f: M^n \rightarrow R \subset R^{2n+1}$  differentiably, with  $L(f) = 0$ . Set  $\delta(x) \equiv 1$ , and let  $g$  be a 1-1 immersion which is a  $\delta$ -approximation to  $f$ . Then  $L(g)$  is empty, so that  $g$  is a homeomorphism.

1.33. Definition. Let  $f: M^n \rightarrow N^p$  differentiably. Let  $N_1^{p-q}$  be a differentiable submanifold of  $N$ . Let  $f(x) \in N_1$ . Let  $(u^1, \dots, u^n)$  be a coordinate system about  $x$ ; and let  $(v^1, \dots, v^p)$  be a coordinate system about  $f(x)$  such that on  $N_1$ ,  $v^1 = \dots = v^q = 0$  (see 1.6). Consider the condition that

$$\frac{\partial v^i}{\partial u^j} \quad \begin{array}{l} i = 1, \dots, q \\ j = 1, \dots, n \end{array}$$

have rank  $q$  at  $x$ . This is the transverse regularity condition for  $f$  and  $N_1$  at  $x$ . [Exercise: Show that this condition is independent of coordinate system.] Note that

the set of points on which the transverse regularity condition is satisfied is an open subset of  $f^{-1}(N_1)$ .  $f$  is said to be transverse regular on  $N_1$  if the condition is satisfied for each  $x$  in  $f^{-1}(N_1)$ .

1.34. Lemma. If  $f: M^n \rightarrow N^p$  is transverse regular on  $N_1^{p-q}$  then  $f^{-1}(N_1)$  is a differentiable submanifold of dimension  $n - q$  (or is empty).

Proof: Let  $\pi$  project  $R^p$  onto its first  $q$  components;  $\pi: R^p \rightarrow R^q$ . If  $(V, h) = (v^1, \dots, v^p)$  is the coordinate system hypothesized in 1.33, then  $N_1 \cap V = h^{-1}\pi^{-1}(0)$  (here  $0$  denotes the origin in  $R^q$ ); and  $f^{-1}(N_1 \cap V) = (\pi h f)^{-1}(0)$ . Since  $\pi h f$  has rank  $q$  at  $x \in f^{-1}(N_1 \cap V)$ , the origin is a regular value of  $\pi h f$ . Hence  $(\pi h f)^{-1}(0)$  is a differentiable submanifold of  $M$  of  $\dim n - q$  (see 1.12).

1.35. Theorem. Let  $f: M^n \rightarrow N^p$  be differentiable; let  $N_1^{p-q}$  be a closed differentiable submanifold of  $N$ . Let  $A$  be a closed subset of  $M$  such that the transverse regularity condition for  $f$  and  $N_1$  holds at each  $x$  in  $A \cap f^{-1}(N_1)$ . Let  $\delta$  be a positive continuous function on  $M$ . There exists a differentiable map  $g: M^n \rightarrow N^p$  such that

- (1)  $g$  is a  $\delta$ -approximation to  $f$ ,
- (2)  $g$  is transverse regular on  $N_1$ , and
- (3)  $g|_A = f|_A$ .



Proof: There is a neighborhood  $U$  of  $A$  in  $M$  such that  $f$  satisfies the transverse regularity condition on  $U \cap f^{-1}(N_1)$ . Cover  $N$  by  $N - N_1 = Y_0$  and coordinate system  $(Y_i, k_i)$  for  $i > 0$ ; with coordinate functions  $(v^1, \dots, v^q)$  such that  $v^1 = \dots = v^q = 0$  on  $N_1$ . Now the open sets  $f^{-1}(Y_i)$  cover  $M$ , as do the open sets  $U$ ,  $M - A$ . Let  $(V_j, h_j)$  be a refinement of both coverings, constructed as in 1.25. Recall that  $h_j(V_j) = C(3)$ ,  $h_j(U_j) = C(2)$ ,  $h_j(W_j) = C(1)$ , and the  $W_j$  cover  $M$ . The  $V_j$  are to be indexed with positive and negative integers so that those  $V_j$  which are contained in  $U$  are the ones with non-positive indices.

Let  $\phi$  be as in 1.26, and define  $\phi_i(x) = \phi(h_i(x))$  for  $x \in V_i$  and  $\phi_i(x) = 0$  elsewhere. For each  $j$  choose  $i(j) \geq 0$  so that  $f(V_j)$  is contained in  $Y_{i(j)}$ .

Set  $f_0 = f$ .

Suppose  $f_{k-1}$  is defined and satisfies the transverse regularity condition for  $N_1$  at each point of the intersection of  $f_{k-1}^{-1}(N_1)$  with  $\cup_{j < k} \bar{W}_j$ . Furthermore suppose that  $f_{k-1}(\bar{U}_j) \subset Y_{i(j)}$  for each  $j$ . Setting  $i = i(k)$ , it follows in particular that  $f_{k-1}(\bar{U}_k) \subset Y_i$ .

Consider  $\pi_{k_i} f_{k-1} h_k^{-1} : C(2) \rightarrow R^q$ ; by 1.21. there is an arbitrarily small affine function  $L(x) = A \cdot (x) + B$  such that when added to the previous function, the resulting map has the origin as a regular value. Consider  $R^q$  as the

first  $q$  coordinates in  $R^D$ , and define

$$f_k(x) = k_1^{-1}(k_1 f_{k-1}(x) + L(h_k(x)) \varphi_k(x)) \text{ for } x \text{ in a neighborhood of } \bar{U}_k$$

$$= f_{k-1}(x) \text{ for } x \text{ in } M - U_k.$$

Here  $L$  is yet to be chosen. Of course, we must choose  $L$  small enough that  $k_1 f_{k-1}^{(*)} + L \varphi_k^{(*)}$  lies in  $C(1)$  for  $x \in \bar{U}_k$ , in order that  $k_1^{-1}$  may be applied to it. This is the first requirement on  $L$ . Secondly, we choose  $L$  small enough that  $f_k$  is a  $\delta/2^k$  approximation to  $f_{k-1}$ . Thirdly choose  $L$  small enough so that  $f_k(\bar{U}_j)$  is contained in  $Y_{i(j)}$  for each  $j$ . This is possible since only a finite number of the sets  $\bar{U}_j$  can intersect  $\bar{U}_k$ .

Now  $f_k$  by definition satisfies the transverse regularity condition for  $N_1$  at each point of  $f_k^{-1}(N_1) \cap \bar{W}_k$ . We want to choose  $L$  small enough that the condition is satisfied at each point of the intersection of  $f_k^{-1}(N_1)$  with  $\cup_{j < k} \bar{W}_j$ . It is sufficient to consider the intersection of this set with  $\bar{U}_k$ ; let this intersection be denoted by  $K$ . Consider the function which maps the pair  $(x, L)$  ( $x \in K$ ) into  $(f_k(x), D(\pi k_1 f_k h_k^{-1})(h_k(x)))$  in  $N \times M(q, n)$ . This function is continuous and carries  $K \times (0)$  into the set  $\{(x, L) \mid (x, L) \in [(N - N_1) \times M(q, n)] \cup [N_1 \times M(q, n; q)]\}$ , which is open in  $N \times M(q, n)$ . Hence for  $L$  sufficiently small,  $(K, L)$  is

$$= N \times M(q, n) - N_1 \times (M(q, n; q))$$

closed

$k_1^{-1}(k_1 f_{k-1}(x) + L \varphi_k(x))$


\*

\*

\*

carried into this set, so that  $f_k$  satisfies the transverse regularity condition for  $N_1$  at each point of  $f_k^{-1}(N_1) \cap (U_{j \leq k} \bar{W}_j)$ .

We define  $g(x) \doteq \lim_{k \rightarrow \infty} f_k(x)$ , as usual.



## Chapter II

Vector Space Bundles

2.1 Definition. An  $n$ -dimensional real vector space bundle  $\xi$  is a triple  $(\pi, a, s)$ .  $\pi$  is a continuous map of  $E$  onto  $B$  ( $E, B$  topological spaces;  $B$  Hausdorff). Let  $F_b = \pi^{-1}(b)$ ; it is called a fibre.  $s$  maps  $R \times E \rightarrow E$  and carries  $R \times F_b$  into  $F_b$ ;  $a$  is defined on  $U_b(F_b \times F_b) \subset E \times E$  and carries  $F_b \times F_b$  into  $F_b$ .

The following must be satisfied:

- (1)  $F_b$  is an  $n$ -dimensional real vector space with  $s$  and  $a$  as scalar product and vector addition, respectively.
- (2) (Local triviality) For each  $b$  in  $B$ , there is a neighborhood  $U$  of  $b$  and a homeomorphism  $\varphi : U \times R^n \rightarrow \pi^{-1}(U)$  such that  $\varphi$  is a vector space isomorphism of  $b' \times R^n$  onto  $F_{b'}$ , for each  $b'$  in  $U$ .

If in (2) the neighborhood  $U$  may be taken as all of  $B$ , the bundle is said to be the trivial bundle.

If  $\xi, \eta$  are  $n$ -dimensional and  $p$ -dimensional vector space bundles, respectively, we define the product bundle  $\xi \times \eta$  as follows:

$$E(\xi \times \eta) = E(\xi) \times E(\eta)$$

$$B(\xi \times \eta) = B(\xi) \times B(\eta)$$

$$(\pi \times \lambda)(x, y) = (\pi(x), \lambda(y))$$

where  $\pi, \lambda$  are the projections in  $\xi, \eta$  respectively and  $F_b(\xi \times \eta)$  has the usual product structure for vector spaces.

If  $U$  is a subset of  $B(\xi)$ , then  $\xi|U$  denotes the bundle  $\pi: \pi^{-1}(U) \rightarrow U$ . It is called the restriction of the bundle to  $U$ .

**2.2 Definition.** Let  $M^n$  be a differentiable manifold and let  $x_0$  be in  $M$ . A tangent vector at  $x_0$  is an operation  $X$  which assigns to each differentiable function  $f$  defined in a neighborhood of  $x_0$ , a real number. The following conditions must be satisfied:

- 1) If  $g$  is a restriction of  $f$ ,  $X(g) = X(f)$ .
- 2)  $X(cf + dg) = cX(f) + dX(g)$  ( $c, d$  real numbers)
- 3)  $X(f \cdot g) = X(f) \cdot g(x_0) + f(x_0) \cdot X(g)$ , where the dot means ordinary real multiplication.

Then  $X(1) = X(1 \cdot 1) = X(1) + X(1)$ , by 3). Hence  $X(1) = 0$ ; and  $X(c)$  also  $= 0$ , by 2).

If one thinks of a tangent vector as being the velocity vector of a curveling in the manifold, then  $X(f)$  is merely the derivative of  $f$  with respect to the parameter of the curve. This is made more precise below.

**2.3 Lemma.** Let  $(u^1, \dots, u^n)$  be a coordinate system about  $x$ . Let  $X$  be a tangent vector at  $x$ . Then  $X$  may be written uniquely as a linear combination of the operators  $\frac{\partial}{\partial u^i}$ :

$$X = \sum \alpha^i \frac{\partial}{\partial u^i}$$

Proof: We assume  $u(x)$  is the origin. Given any  $f(u^1, \dots, u^n)$  define

$$\begin{aligned} g_1(u^1, \dots, u^n) &= \frac{f(u^1, \dots, u^n) - f(0, u^2, \dots, u^n)}{u^1} \quad \text{if } u^1 \neq 0 \\ &= \partial f / \partial u^1 (0, u^2, \dots, u^n) \quad \text{if } u^1 = 0. \end{aligned}$$

To see that  $g_1$  is differentiable, note that

$$g_1(s, u^2, \dots, u^n) = \int_0^1 \frac{\partial f}{\partial u^1}(s t, u^2, \dots, u^n) dt.$$

(Then  $f(u^1, \dots, u^n) = u^1 g_1(u^1, \dots, u^n) + f(0, u^2, \dots, u^n)$ .)

Similarly,  $f(0, u^2, \dots, u^n) = u^2 g_2(u^2, \dots, u^n) + f(0, 0, u^3, \dots, u^n)$ ,

where  $g_2(0) = \partial f / \partial u^2(0)$ . Finally, we have

$$f(u^1, \dots, u^n) = \sum u^i g_i + f(0), \quad \text{where } g_i(0) = \frac{\partial f}{\partial u^i}(0).$$

Thus  $X(f) = \sum X(u^i) g_i(0) + 0 \cdot X(g_i)$

$$= \sum \alpha^i \frac{\partial f}{\partial u^i}(0), \quad \text{where } \alpha^i = X(u^i).$$

Remark. If  $(v^1, \dots, v^n)$  is another coordinate system about  $x$ , and  $X = \sum \beta^j \frac{\partial}{\partial v^j}$ , then  $\alpha^i = X(u^i) = \sum_j \beta^j \frac{\partial u^i}{\partial v^j}$ .

The  $\alpha^i$  are called the components of the vector  $X$  with respect to the coordinate system  $(u^1, \dots, u^n)$ .

2.4 Alternate definition. A tangent vector at  $x$  is an assignment to every coordinate system  $(u^1, \dots, u^n)$  about  $x$  of an element  $(\alpha^1, \dots, \alpha^n)$  of  $\mathbb{R}^n$ , with the requirement that if  $(\beta^j)$  is assigned to the system  $(v^1, \dots, v^n)$ , then

$$\alpha^i = \sum_j \frac{\partial u^i}{\partial v^j} \beta_j .$$

The derivation operator  $X$  is then defined

as  $\sum \alpha^i \frac{\partial}{\partial u^i}$ . One checks readily that

- a)  $X(f)$  is independent of the coordinate system used, and
- b)  $X(f)$  satisfies requirements 1), 2), and 3) for a tangent vector.

2.5. Definition. For each  $x$  in  $M$ , the tangents at  $x$  form an  $n$ -dimensional vector space (the operations  $\partial/\partial u^i$  form a basis, by 2.3). Let the totality of these be denoted  $E(\tau)$ ; define  $\pi: E(\tau) \rightarrow M$  as mapping the tangent vector  $X$  at  $x_0$  into  $x_0$ . The local product structure is given by  $\varphi_u: U \times \mathbb{R}^n \rightarrow E$ , where  $(U, h) = (u^1, \dots, u^n)$  is a coordinate system on  $M$ , and  $\varphi$  is defined as follows:

$$\varphi(x_0, a^1, \dots, a^n) = \text{the tangent vector } X = \sum a^i \frac{\partial}{\partial u^i} \text{ at } x_0 .$$

Since  $\varphi$  is to be a homeomorphism, this structure imposes a topology on  $E$ ; since  $\varphi_v^{-1} \varphi_u$  is a homeomorphism on  $(U \cap V) \times \mathbb{R}^n$

this topology is unambiguously determined. One checks immediately that  $\varphi$  gives us a vector space isomorphism for each fibre.

Indeed,  $\varphi_v^{-1} \varphi_u$  is a  $C^\infty$  map on  $(U \cap V) \times \mathbb{R}^n$ , so that  $E$  is a differentiable manifold of dimension  $2n$  (using definition 1.2 of a differentiable manifold). The map  $\pi$  is differentiable of rank  $n$ .

This bundle  $\tau$  is called the tangent bundle of  $M$ .

**2.6 Definition.** If  $f: M_1 \rightarrow M_2$ , there is an induced map  $df: E(\tau_1) \rightarrow E(\tau_2)$  defined as follows:  $df(X) = Y$ , where  $Y(g) = X(gf)$ . If  $X$  is a vector at  $x_0$ ,  $Y$  is a vector at  $f(x_0)$ . This is clearly linear on each fibre; it is called the derivative map.

If  $(U, h)$  and  $(V, k)$  are coordinate systems about  $x_0, f(x_0)$  respectively, and  $(\alpha^i), (\beta^j)$  are the respective components of  $X$  and  $Y$  with respect to these coordinate systems, then  $(\beta^j) = D(kfh^{-1}) \cdot (\alpha^i)$  where the vector components are written as column matrices, as usual.

**2.7 Definition.** Let  $\xi, \eta$  be two  $n$ -dimensional vector space bundles.

A bundle map  $f: \xi \rightarrow \eta$  is a continuous map of  $E(\xi)$  into  $E(\eta)$  which carries each fibre isomorphically



onto a fibre. The induced map  $f_B: B(\xi) \rightarrow B(\eta)$  is automatically continuous.

If  $B(\xi) = B(\eta)$  and the induced map is the identity,  $f$  is said to be an equivalence. Note that if  $f$  is an equivalence, it is a homeomorphism: Locally  $f$  is just a map  $U \times \mathbb{R}^n \rightarrow V \times \mathbb{R}^n$ . The projection of  $f^{-1}$  into the factor  $U$  is continuous, because  $f_B^{-1}$  is the identity. But  $f$  may be given by a non-singular matrix function of  $x \in U$ ;  $f^{-1}$  is the inverse of this matrix, so that the projection of  $f^{-1}$  into the factor  $\mathbb{R}^n$  is continuous. Hence  $f^{-1}$  is continuous.

If there is an equivalence of  $\xi$  onto  $\eta$ , we write  $\xi \simeq \eta$ .

**2.8 Lemma.** Given a bundle  $\eta$  with projection map  $\lambda: E(\eta) \rightarrow B(\eta)$ , and a map  $f: B_1 \rightarrow B(\eta)$ , there is a bundle  $\pi: E_1 \rightarrow B_1$  and a bundle map  $g: E_1 \rightarrow E(\eta)$  such that  $\lambda g = f\pi$ . Furthermore,  $E_1$  is unique up to an equivalence.

[ $E_1$  is called induced bundle and is often denoted by  $f^*\eta$ .]

Proof: Let  $E_1$  be that subset of  $B_1 \times E(\eta)$  consisting of points  $(b, e)$  such that  $f(b) = \lambda(e)$ . Define  $\pi(b, e) = b$ ;  $g(b, e) = e$ . To show that  $E_1$  is a vector

space bundle, let  $\varphi: V \times \mathbb{R}^n \rightarrow E(\eta)$  be a product neighborhood in  $E(\eta)$ , and let  $f(U) \subset V$ . Then define  $\varphi: U \times \mathbb{R}^n \rightarrow E_1$  by  $\varphi_1(b, x) = (b, \varphi(f(b), x))$ . This is continuous and 1-1; its image equals  $\pi^{-1}(U)$ . Its inverse carries  $(b, e)$  into  $(b, p \varphi^{-1}(e))$  (where  $p$  projects  $V \times \mathbb{R}^n$  onto  $\mathbb{R}^n$ ), so that it is continuous. The map  $g$  is an isomorphism on each fibre.

Now suppose  $g': E' \rightarrow E(\eta)$  is a bundle map, where  $\pi': E' \rightarrow B_1$  is a bundle and  $\lambda g' = f \pi'$ . We map  $E' \rightarrow E_1$  by mapping  $e' \rightarrow (\pi'(e'), g'(e'))$  in  $E_1$ . Because  $g'$  is an isomorphism on each fibre, so is this map; and it induces the identity on the base space. Hence it is an equivalence.

2.9 Definition. Let  $\xi, \eta$  be two bundles over  $B$ .

The Whitney sum  $\xi \oplus \eta$  is a bundle defined as follows:

Consider the product bundle  $E(\xi) \times E(\eta) \rightarrow B \times B$ ; let  $d$  be the diagonal map  $B \rightarrow B \times B$ . The induced bundle  $d^*(\xi \times \eta)$  is defined as the Whitney sum  $\xi \oplus \eta$ .

Note that the fibre over  $b$  in  $\xi \oplus \eta$  is merely  $F_b(\xi) \times F_b(\eta)$ , so that  $\dim(\xi \oplus \eta) = \dim \xi + \dim \eta$ .

Note also the commutativity and associativity of  $\oplus$ . I.e.,  $\xi \oplus \eta \simeq \eta \oplus \xi$  and  $(\xi \oplus \eta) \oplus \zeta \simeq \xi \oplus (\eta \oplus \zeta)$ . The proof is left as an exercise.

**2.10 Definition.** If  $\xi, \eta$  are bundles over  $B$ , then  $g: E(\xi) \rightarrow E(\eta)$  is a homomorphism if

- 1) it maps each fibre linearly into a fibre, and
- 2) the induced map on  $B$  is the identity.

Note that an equivalence is both a bundle map and a homomorphism. An irbedding of bundles is a 1 - 1 homomorphism.

**2.11 Theorem.** If  $f: E(\xi) \rightarrow E(\eta)$  maps each fibre linearly into a fibre, then  $f$  may be factored into a homomorphism followed by a bundle map.

Proof: Let  $\pi_1, \pi_2$  be the projections in  $\xi, \eta$ , respectively.

Let  $f_B: B(\xi) \rightarrow B(\eta)$  be the map induced by  $f$ . Let  $E_1 = f_B^* \eta$  be the bundle induced by  $f_B$ ; let  $g$  be the bundle map  $E_1 \rightarrow E(\eta)$  and  $\pi$  the projection  $E_1 \rightarrow B(\xi)$ .

Define  $h: E(\xi) \rightarrow B(\xi) \times E(\eta)$  by the equation  $h(e) = (\pi_1(e), f(e))$ . The image of  $h$  actually lies in that subset of  $B(\xi) \times E(\eta)$  which is  $E_1$ ; then  $h$  is a homomorphism. From the definition,  $f = gh$ .

$$\begin{array}{ccccc} E(\xi) & \xrightarrow{h} & E_1 & \xrightarrow{g} & E(\eta) \\ \downarrow \pi_1 & & \downarrow \pi & & \downarrow \pi_2 \\ B(\xi) & \xrightarrow{1} & B(\xi) & \xrightarrow{f_B} & B(\eta) \end{array}$$

note

$$g(\pi_1(e)) = \pi_2(f(e))$$

2.12 Lemma. Let  $\xi, \eta$  be bundles over  $B$  of dimensions  $n, p$ , respectively; let  $g: \xi \rightarrow \eta$  be a homomorphism. If  $g$  is onto, then kernel  $g$  is a bundle. If  $g$  is 1-1, then cokernel  $g$  (i.e., the quotient,  $\eta/\text{image } g$ ) is a bundle.

Proof: Suppose  $g$  is 1-1 (i.e., has rank  $n$  when restricted to each fibre). In  $E(\eta)$ , we define  $e \sim e'$  if  $e - e'$  exists and is in the image of  $g$ . We identify the elements of these equivalence classes; the resulting identification space is defined to be  $E(\eta/g(\xi))$ . It is a bundle over  $B$  with projection naturally defined and each fibre is a vector space of  $\dim p - n$ . We need only to show the existence of a local product structure.

Let  $U$  be an open set in  $B$ , with  $\xi|U$  equivalent to  $U \times \mathbb{R}^n$  and  $\eta|U$  equivalent to  $U \times \mathbb{R}^p$ . Let  $g_0$  denote the homomorphism of  $U \times \mathbb{R}^n \rightarrow U \times \mathbb{R}^p$  induced by  $g$ . Now  $(\eta/g(\xi))|U$  is equivalent to the quotient  $U \times \mathbb{R}^p / g_0(U \times \mathbb{R}^n)$ , so that it suffices to show that this latter quotient is locally a product.

$g_0$  is given by a matrix  $M(b) \in M(p, n)$  which depends continuously on the point  $b \in U$ . Given  $b_0$ , we may assume that in a neighborhood  $U_0$  of  $b_0$ , the first  $n$  rows are independent. We define  $h: U_0 \times \mathbb{R}^n \times \mathbb{R}^{p-n} \rightarrow U_0 \times \mathbb{R}^p$  as the linear function on  $\mathbb{R}^p$  whose matrix (non-singular) is

$$\left( M(b) \mid \begin{array}{c} 0 \\ I_{p-n} \end{array} \right)$$

The image of  $U_0 \times R^n \times 0$  under  $h$  is just  $g_0(U_0 \times R^n)$ ; since  $h$  is an equivalence, it induces an equivalence of

$$U_0 \times R^{p-n} \simeq \frac{U_0 \times R^n \times R^{p-n}}{U_0 \times R^n \times 0} \text{ onto } \frac{U_0 \times R^p}{g_0(U_0 \times R^n)}.$$

*not zero*

Secondly, suppose  $g$  is onto (i.e., it has rank  $p$  on each fibre).  $E(g^{-1}(0))$  is defined as that subset of  $E(\xi)$  consisting of points  $e$  with  $g(e) = 0$ . Again, we need to show the existence of a local product structure. Let  $U$ ,  $g_0$ , and  $M(b)$  be as above. Given  $b_0$ , we may assume that the first  $p$  columns of  $M(b)$  are independent in the neighborhood  $U_0$  of  $b_0$ . We define  $h: U_0 \times R^n \rightarrow U_0 \times R^p \times R^{n-p}$  by the matrix function

$$\left( \begin{array}{c} M(b) \\ 0 \mid I_{n-p} \end{array} \right)$$

Now  $h$  followed by the natural projection of  $U_0 \times R^p \times R^{n-p}$  onto  $U_0 \times R^p$  equals  $g_0|_{U_0}$ . Hence  $h^{-1}$  maps  $U_0 \times 0 \times R^{n-p}$  onto  $g_0^{-1}(U_0 \times 0)$ ; since  $h$  is an equivalence, the restriction of  $h^{-1}$  to  $U_0 \times 0 \times R^{n-p}$  is also.

Remark. If  $g$  is onto,  $\xi/g^{-1}(0)$  is a bundle, being the quotient of the inclusion homomorphism  $g^{-1}(0) \rightarrow \xi$ . If  $g$  is 1-1,  $g(\xi)$  is a bundle, being the kernel of the projection homomorphism  $\eta \rightarrow \eta/g(\xi)$ .

2.13 Definition. If  $\varphi$  is a non-negative function on  $B$ , the carrier of  $\varphi$  is the closure of the set of  $x$  with  $\varphi(x) > 0$ . A partition of unity is a collection  $\varphi_\alpha$  of continuous non-negative functions on  $B$ , such that the sets  $C_\alpha = \text{carrier } \varphi_\alpha$  form a locally-finite covering of  $B$ , and  $\sum \varphi_\alpha(x) = 1$  (this is a finite sum for each  $x$ ).

2.14 Lemma. Let  $B$  be a normal space;  $U_\alpha$  a locally-finite open covering of  $B$ . Then there is a partition of unity  $\varphi_\alpha$  with carrier  $\varphi_\alpha \subset U_\alpha$  for each  $\alpha$ .

Proof: First, we show that there is an open covering  $V_\alpha$  of  $B$  with  $\bar{V}_\alpha \subset U_\alpha$  for each  $\alpha$ . Assume the  $U_\alpha$  indexed by a set of ordinals (well ordering theorem). Let  $V_\alpha$  be defined for all  $\alpha < \beta$  and assume that the sets  $V_\alpha$  along with the sets  $U_\alpha$  for  $\alpha \geq \beta$  cover  $B$ . Consider the set  $A(\beta) = B - U_{\alpha < \beta} \cup V_\alpha - U_{\alpha > \beta} \cup U_\alpha$ . Then  $A(\beta) \subset U_\beta$ . Let  $V_\beta$  be an open set containing the closed set  $A(\beta)$ , with  $\bar{V}_\beta \subset U_\beta$  (normality). This completes the construction of the  $V_\alpha$ .

Now let  $g_\alpha$  be a function which is positive on  $\bar{V}_\alpha$  and 0 outside  $U_\alpha$  (normality again). Define  $\varphi_\alpha(x) = g_\alpha(x) / \sum g_\alpha(x)$ . Since  $U_\alpha$  is locally-finite, the sum in the denominator is finite and positive, so  $\varphi_\alpha$  is well-defined.

Remark. If  $B$  is a differentiable manifold,  $\varphi_\alpha$  may be chosen to be differentiable: Cover  $B$  with coordinate systems  $(V_i, h_i)$  as in 1.25 refining the covering  $U_\alpha, B - \bar{V}_\alpha$ . Let  $\varphi_i(y) = \varphi(h_i(y))$  for  $y \in V_i$ , and  $= 0$  otherwise ( $\varphi$  as in 1.26). Let  $g_\alpha(y) = \sum \varphi_i(y)$ , where the sum extends over all  $i$  such that  $V_i \subset U_\alpha$ .

2.15 Lemma. Let  $B$  be paracompact and let  $0 \longrightarrow \xi \xrightarrow{i} \eta \xrightarrow{\varphi} \zeta \longrightarrow 0$  be an exact sequence of homomorphisms of bundles. Then there is equivalence  $f: \eta \longrightarrow \xi \oplus \zeta$ , with  $f|_i$  the natural inclusion and  $\varphi f^{-1}$  the natural projection.

Proof. Let  $\dim \xi = n$ ;  $\dim \zeta = p$ .

We first construct a Riemannian metric on  $\eta$  (i.e., a continuous inner product in  $E(\eta)$ ). Let  $U_\alpha$  be a locally-finite covering of  $B$  with  $\eta|_{U_\alpha}$  trivial; let  $g_\alpha$  be the corresponding projection of  $\eta|_{U_\alpha}$  onto  $\mathbb{R}^{n+p}$ . Let  $\varphi_\alpha$  be a partition of unity with carrier  $\varphi_\alpha \subset U_\alpha$ .

If  $e, e'$  are in  $E(\eta)$  and  $\pi(e) = \pi(e')$ , define  $e \cdot e' = \sum_{\alpha} \varphi_{\alpha}(\pi(e)) g_{\alpha}(e) \cdot g_{\alpha}(e')$ , where the dot on the right hand side is the ordinary scalar product in  $R^{n+p}$ . This is a finite sum; it satisfies the axioms for a scalar product.

The way we use the Riemannian metric is to break  $\eta$  up into  $iE(\xi)$  and its orthogonal complement. Let  $\xi'$  be the image of  $\xi$  in  $\eta$  and let  $E(\xi')$  be defined as that subset of  $E(\eta)$  consisting of elements which are orthogonal to  $i(E(\xi))$ . In order to show that  $\xi'$  has a local product structure, consider the homomorphism

$$h: \eta \rightarrow \xi'$$

which sends each vector into its orthogonal projection in  $\xi'$ .

[Verification that  $h$  is continuous. Over any coordinate neighborhood  $U$  we can choose a basis  $a_1, \dots, a_n$  for the fibre of  $\xi'$ . Then the function  $h$  carries  $v \in E(\eta)$  into  $\sum t_j a_j \in E(\xi') \subset E(\eta)$ , where  $t_j = \sum B_{jk}(v \cdot a_k)$  and where  $(B_{jk})$  denotes the inverse matrix to  $(a_j \cdot a_k)$ .] Since  $h$  is onto, its kernel  $\xi'$  is again a vector space bundle.

Now the bundle  $i(\xi) = \xi'$  is equivalent to  $\xi$ . It remains to show that  $\xi'$  is equivalent to  $\xi$  and that  $\eta$  is equivalent to  $\xi' \oplus \xi'$ . The former follows immediately from the fact that  $\varphi|_{\xi'}$  is a homomorphism; from rank considerations it must be 1 - 1 and onto as well. The latter follows



by noting that  $E(\xi' \oplus \zeta')$  is defined as the subset of  $E(\xi') \times E(\zeta')$  consisting of points  $(e_1, e_2)$  such that  $\pi(e_1) = \pi(e_2)$ . Consider the map  $f$  of  $E(\xi' \oplus \zeta')$  into  $E(\eta)$  obtained by taking  $(e_1, e_2)$  into their sum in  $E(\eta)$  (this sum exists because  $e_1$  and  $e_2$  lie in the same fibre). This is clearly a homomorphism; from rank considerations, it must be 1 - 1 and onto.

2.16 Definition. Let  $M_1, M_2$  be differentiable manifolds.

Let  $f$  be an immersion  $M_1 \rightarrow M_2$ . The normal bundle  $\nu_f$  is defined as follows:

Let  $\tau_1, \tau_2$  be the tangent bundles of  $M_1, M_2$  respectively. By 2.11, the map  $df: E(\tau_1) \rightarrow E(\tau_2)$  may be factored into a homomorphism  $h$  of  $E(\tau_1)$  into  $E(f^*\tau_2)$  followed by a bundle map  $g$ . Now  $h$  is a 1 - 1 homomorphism because  $f$  is an immersion; hence by 2.12,  $f^*\tau_2/\text{image } g^h$  is a bundle over  $M_1$ . It is called the normal bundle  $\nu_f$ .

Then  $0 \rightarrow \tau_1 \rightarrow f^*\tau_2 \rightarrow \nu_f \rightarrow 0$  is an exact sequence of homomorphisms, so that by 2.15,  $f^*\tau_2$  is equivalent to  $\tau_1 \oplus \nu_f$ . Indeed, given a Riemannian metric on  $f^*\tau_2$ ,  $\nu_f$  is equivalent to the orthogonal complement of the image of  $\tau_1$ .

Let us consider the case  $M_2 = R^{n+p}$ , where  $\dim M_1 = n$ . Then  $\tau_2$  is the trivial bundle, so that  $f^* \tau_2$  is as well. (Proof: If  $f: B \rightarrow B(\eta)$  and  $\eta$  is trivial, so is  $f^* \eta$ .)

We have

$$\begin{array}{c} B \times R^n \\ \downarrow \pi \\ f: B_1 \rightarrow B \end{array}$$

$E(f^* \eta)$  is defined as that subset of  $B_1 \times (B \times R^n)$  consisting of points  $(b_1, b, x)$  such that  $f(b_1) = \pi(b, x)$ ; i.e., of all points  $(b_1, f(b_1), x)$ . If we map this into  $(b_1, x)$ , we obtain an equivalence of  $f^* \eta$  with the bundle  $B_1 \times R^n \rightarrow B_1$ .)

Thus  $\tau_1 \oplus \nu_f$  is equivalent to a trivial bundle. In what follows, we investigate the following question: Given  $\xi$ , does there exist an  $\eta$  with  $\xi \oplus \eta$  trivial? Using 1.28, this is always the case for  $\xi$  the tangent bundle of an  $n$ -manifold, and indeed  $\eta$  may be chosen also to have dimension  $n$ . A more general answer appears in 2.19.

**2.17 Definition.** Let  $f: M_1 \rightarrow M_2$ ; let  $\dim M_1 = n$ ,  $\dim M_2 = p$ . If  $f$  has rank  $p$  at every point of  $M_1$ , it is said to be regular. If  $f$  is regular, the homomorphism  $h: \tau_1 \rightarrow f^* \tau_2$  given by 2.11 is an onto map. By 2.12, the kernel of  $h$  is a bundle  $\alpha_f$ . It is called the bundle along the fibre.

Note that  $f^{-1}(y)$  is a submanifold of  $M_1$  of dim  $n-p$  (by 1.12 or 1.34). The inclusion  $i_y$  of  $f^{-1}(y)$  into  $M_1$  induces an inclusion  $di_y$  of its tangent bundle into  $\tau_1$ . The kernel of  $h$  consists precisely of the vectors which are in the image of some  $di_y$ , i.e., the vectors tangent to the submanifolds  $f^{-1}(y)$  are the ones carried into 0 by  $h$ .

One has the exact sequence  $0 \rightarrow \alpha_f \rightarrow \tau_1 \xrightarrow{g} f^* \tau_2 \rightarrow 0$ , so that by 2.15,  $\tau_1$  is equivalent to  $\alpha_f \oplus f^* \tau_2$ .

2.18 Definition. A bundle  $\xi$  is of finite type if  $B$  is normal and may be covered by a finite number of neighborhoods  $U_1, \dots, U_k$  such that  $\xi|_{U_i}$  is trivial for each  $i$ .

2.19 Lemma.  $\xi$  is of finite type if  $B$  is compact, or paracompact finite dimensional.

The former statement is clear; let us consider the latter. By definition, the dimension of  $B$  is not greater than  $n$  if every open covering has an open refinement such that (\*) no point of  $B$  is contained in more than  $n+1$  elements of the refinement. It is a standard theorem of topology that an  $n$ -manifold has dimension  $n$  in this sense.

Cover  $B$  by open sets  $U$ , with  $\xi|_U$  trivial; let  $\{V_\alpha\}$  be an open refinement of this covering satisfying (\*).

By 1.22, we may assume that  $\{V_\alpha\}$  is locally-finite as well. Let  $\varphi_\alpha$  be a partition of unity with carrier  $\varphi_\alpha \subset V_\alpha$  for each  $\alpha$  (2.14).

Let  $A_i$  be the set of unordered  $i + 1$  tuples of distinct elements of the index set of  $\{\varphi_\alpha\}$ . Given  $a$  in  $A_i$ , where  $a = \{\alpha_0, \dots, \alpha_i\}$ , let  $W_{ia}$  be the set of all  $x$  such that  $\varphi_\alpha(x) < \min[\varphi_{\alpha_0}(x), \dots, \varphi_{\alpha_i}(x)]$  for all  $\alpha \neq \alpha_0, \dots, \alpha_i$ . Each set  $W_{ia}$  is open, and  $W_{ia}$  and  $W_{ib}$  are disjoint if  $a \neq b$ . Also  $W_{ia}$  is contained in the intersection of the carriers of  $\varphi_{\alpha_0}, \dots, \varphi_{\alpha_i}$ , and hence in some set  $V_\alpha$ . If we set  $X_i$  equal to the union of all sets  $W_{ia}$ , for fixed  $i$ , the result is that  $\xi|X_i$  is trivial. (For  $\xi|W_{ia}$  is trivial, and the  $W_{ia}$  are disjoint.)

Finally, the sets  $X_0, \dots, X_n$  cover  $B$ . Given  $x$  in  $B$ ,  $x$  is contained in at most  $n + 1$  of the sets  $V_\alpha$ , so that at most  $n+1$  of the functions  $\varphi_\alpha$  are positive at  $x$ . Since some  $\varphi_\alpha$  is positive at  $x$ ,  $x$  is contained in one of the sets  $W_{ia}$  for  $0 \leq i \leq n$ .

[The intuitive idea of the proof is as follows: Consider an  $n$ -dimensional simplicial complex, with  $\varphi_\alpha$  the barycentric coordinate of  $x$  with respect to the vertex  $\alpha$ . The sets  $W_{0a}$  will be disjoint neighborhoods of the vertices, the sets  $W_{1a}$  disjoint neighborhoods of the open 1-simplices, and so on.]

2.20 Theorem. If  $\xi$  is of finite type, there is a bundle  $\eta$  such that  $\xi \oplus \eta$  is trivial.

Proof: We proceed by showing that  $\xi$  may be imbedded in a trivial bundle  $B \times \mathbb{R}^m$ , so that we have the exact sequence  $0 \rightarrow \xi \xrightarrow{i} B \times \mathbb{R}^m \rightarrow B \times \mathbb{R}^m / i(\xi) \rightarrow 0$  by 2.12. The theorem then follows from 2.15. (Paracompactness is not needed since the trivial bundle clearly has a Riemannian metric.)

Cover  $B$  by finitely many neighborhoods  $U_1, \dots, U_k$  with  $\xi|_{U_i}$  trivial for each  $i$ . Let  $\varphi_1, \dots, \varphi_k$  be a partition of unity with carrier  $\varphi_i \subset U_i$  for each  $i$  (2.14). Let  $f_i$  denote the equivalence of  $E(\xi|_{U_i})$  onto  $U_i \times \mathbb{R}^n$ ; let  $f_i^1, \dots, f_i^n$  denote the coordinate functions of its projection into  $\mathbb{R}^n$ .

We define  $h: E(\xi) \rightarrow B \times \mathbb{R}^{nk}$  as follows:

$$h(e) = (\pi(e), \varphi_i(\pi(e)) \cdot f_i^j(e)) \quad \begin{array}{l} i = 1, \dots, k \\ j = 1, \dots, n \end{array}$$

(no summation is indicated). This is well-defined, since  $\varphi_i(\pi(e)) = 0$  unless  $e \in E(\xi|_{U_i})$ . It is clearly a homomorphism, since each  $f_i^j$  is linear on  $E(\xi|_{U_i})$ . To show that it is 1-1, let  $e \neq 0$ . Then for some  $i$ ,  $\varphi_i(\pi(e)) > 0$ . Since  $f_i$  is an equivalence,  $f_i^j(e) \neq 0$  for some  $j$ . Hence  $h(e) \neq (\pi(e), 0)$ , as desired.

2.21 Definition. The bundle  $\xi$  is s-equivalent to  $\eta$  if there are trivial bundles  $o^p, o^n$  such that  $\xi \oplus o^p \simeq \eta \oplus o^n$ .

Here  $o^p = B \times R^p$ . Symmetry and reflexivity are clear. To show transitivity, assume  $\xi \oplus o^p \simeq \eta \oplus o^q$  and  $\eta \oplus o^r \simeq \zeta \oplus o^s$ . Then  $\xi \oplus o^p \oplus o^r \simeq \zeta \oplus o^s \oplus o^q$ .

Note that s-equivalence differs from equivalence. E.g., consider the two-sphere  $S^2$  in  $R^3$ . Then  $\tau^2 \oplus \nu^1 = o^3$ . The normal bundle  $\nu^1$  is easily seen to be trivial; but it is a classical theorem of topology that  $\tau^2$  is not (it does not admit a non-zero cross-section). Hence  $\tau^2$  is s-trivial, but not trivial.

2.22 Theorem. The set of s-equivalence classes of vector space bundles of finite type over  $B$  forms an abelian group under  $\oplus$ .

Proof. To avoid logical difficulties, we consider only subbundles of  $B \times R^m$ , for all  $m$ . This suffices, since any bundle  $\xi$  of finite type may be imbedded in some  $B \times R^m$ , by 2.20.

The class of trivial bundles  $o^p$  is the identity element. The existence of inverses is the substance of 2.20.

2.23 Corollary. Given two immersions of the differentiable manifold  $M$  in euclidean space, their normal bundles are  $s$ -equivalent.

2.24 Definition.  $M^n$  is a  $\pi$ -manifold if  $M$  may be immersed in some  $R^{n+p}$  so that its normal bundle is trivial.

This is equivalent to the requirement that  $\tau^n$  be  $s$ -trivial: Let  $\tau^n$  be  $s$ -trivial. If we take some immersion of  $M$  into  $R^{n+p}$ , then  $\tau^n \oplus \nu^p$  is trivial by 2.16, so that  $\nu^p$  is  $s$ -trivial, i.e.  $\nu^p \oplus o^q = o^{p+q}$  for some  $q$ . Consider the composite immersion  $M \rightarrow R^{n+p} \subset R^{n+p+q}$ . The normal bundle of  $M$  in  $R^{n+p+q}$  is just  $\nu^p \oplus o^q$ , which is trivial.

Conversely, if  $\nu^p$  is trivial for some immersion, then  $\tau^n$  is  $s$ -trivial because  $\tau^n \oplus \nu^p$  is trivial.

2.25 Definition. Let  $G_{p,n}$  denote the set of all  $n$ -dimensional vector subspaces of  $R^{n+p}$  (i.e., all  $n$ -dim hyperplanes through the origin). It is called the Grassman manifold of  $n$ -planes in  $n+p$  space.

Its topology is obtained as follows: Consider  $M(n, n+p; n)$ ; we identify two elements of this set if the hyperplanes spanned by their row vectors are the same.  $G_{p,n}$  is in 1-1 correspondence with this identification space, and is

given the identification topology. Let  $\rho$  be the projection of  $M(n, n+p; n) \rightarrow G_{p, n}$ .

Now  $\rho(A) = \rho(B)$  if and only if  $A = CB$  for some non-singular  $n \times n$  matrix  $C$ : The hyperplane  $\rho(A)$  consists of all points  $(x', \dots, x^{n+p}) \in R^{n+p}$  which equal  $(c', \dots, c^n) \cdot A$  for some choice of constants  $c^i$ . If  $\rho(A) = \rho(B)$ , then

$$(1, 0, \dots, 0) \cdot A = (c_1^1, \dots, c_1^n) \cdot B$$

$$(0, 1, \dots, 0) \cdot A = (c_2^1, \dots, c_2^n) \cdot B, \text{ etc., for some choice of } c_i^j.$$

Then  $IA = CB$ , where  $C$  has rank  $n$  because  $A$  does. The converse is clear.

(a)  $G_{p, n}$  is locally euclidean. Let  $A \in M(n, n+p; n)$ ; after permuting the columns, we may assume  $A = (P, Q)$  where  $P$  is  $n \times n$  and non-singular. Let  $U$  be the set of all such  $A$ ; it is an open set in  $M(n, n+p; n)$ , being the inverse image of the non-zero reals under the continuous map  $(P, Q) \rightarrow \det P$ . If  $\rho(P, Q) = \rho(R, S)$ , where  $P$  is non-singular, then  $(P, Q) = (CR, CS)$  for some non-singular  $C$ . Hence  $R$  is necessarily non-singular; it follows that  $\rho^{-1}(\rho(U)) = U$ , so that  $\rho(U)$  is open in  $G_{p, n}$  (by definition of the identification topology).

We show  $\rho(U)$  homeomorphic with  $R^{p, n}$ . Define  $\varphi: U \rightarrow R^{p, n}$  by  $\varphi(P, Q) = P^{-1}Q$ . If  $\rho(P, Q) = \rho(R, S)$  then  $(P, Q) = (CR, CS)$ , so that  $P^{-1}Q = (CR)^{-1}(CS) = R^{-1}S$ . Hence



$\phi$  induces a continuous map  $\phi_0 : \rho(U) \rightarrow \mathbb{R}^{p \times n}$ . Define  $\psi : \mathbb{R}^{p \times n} \rightarrow \rho(U)$  by  $\psi(Q) = \rho(I, Q)$  where  $Q$  is an  $n \times p$  matrix. One checks immediately that  $\psi$  and  $\phi_0$  are inverses of each other.

$$\begin{array}{ccc}
 M(n, n+p; n) \supset U & & \\
 \downarrow \rho & \searrow \phi & \\
 G_{p \times n} \supset \rho(U) & \xrightarrow{\phi_0} & \mathbb{R}^{p \times n} \\
 & \xleftarrow{\psi} & 
 \end{array}$$

(b) To show that  $G_{p \times n}$  is Hausdorff, we show that  $\psi$  maps every compact set into a closed set (this will clearly suffice). Let  $K$  be a compact subset of  $\mathbb{R}^{p \times n}$ ; we show  $\psi^{-1}(K)$  is closed in  $M(n, n+p; n)$ .  $\psi^{-1}(K)$  consists of all matrices  $(P, Q)$  with  $P$  non-singular and  $P^{-1}Q \in K$ . Let  $(P, Q) \in M(n, n+p; n)$  be the limit of the sequence  $(P_i, Q_i)$  of elements of  $\psi^{-1}(K)$ . Since  $K$  is compact some subsequence of the sequence  $\phi(P_i, Q_i) = P_i^{-1}Q_i$  converges to a point  $R$  of  $K$ . The corresponding subsequence of the sequence  $Q_i$  converges to  $PR$ , so that  $(P, Q) = P(I, R)$ . Since  $(P, Q)$  has rank  $n$  it follows that  $P$  is non-singular, so that  $(P, Q) \in \psi^{-1}(K)$ , as desired.

Hence  $G_{p \times n}$  is a manifold of dimension  $p \times n$ .

(c)  $G_{p \ n}$  is a differentiable manifold and  $\rho$  is a differentiable map. A function  $f$  on the open set  $V$  in  $G_{p \ n}$  belongs to the differentiable structure  $\mathcal{D}$  if  $f \circ \rho$  is differentiable. To show that this satisfies the conditions for a differentiable structure, we show that  $(\rho(U), \varphi_0)$ , as defined in (a), is a coordinate system. Let  $f$  be defined on  $V \subset \rho(U)$ . Given  $Q \in \mathbb{R}^{p \ n}$ ,  $f \circ \varphi_0^{-1}(Q) = f \circ \rho(I, Q)$  so that  $f \circ \varphi_0^{-1}$  is differentiable if  $f \circ \rho$  is. Conversely, given  $(P, Q) \in V$ ,  $f \circ \rho(P, Q) = f \circ \varphi_0^{-1} \circ \varphi_0 \circ \rho(P, Q) = f \circ \varphi_0^{-1}(P^{-1}Q)$ , so that  $f \circ \rho$  is differentiable if  $f \circ \varphi_0^{-1}$  is.

(d)  $G_{p \ n}$  is compact. Let  $L$  be the subset of  $M(n, n+p; n)$  consisting of matrices whose rows are orthonormal vectors.  $L$  is a closed and bounded subset of  $\mathbb{R}^{n(n+p)}$ . Since  $\rho(L) = G_{p \ n}$  (the Gram-Schmidt orthogonalization process proves this),  $G_{p \ n}$  is compact.

(e)  $G_{p \ n}$  is diffeomorphic to  $G_{n \ p}$ . Geometrically, the homeomorphism  $h$  is defined as carrying each hyperplane into its orthogonal complement. It is clearly 1-1; to show it differentiable we use the coordinate system  $(\rho(U), \varphi_0)$  defined in (a). Let  $g$  map  $U$  into  $M(p, n+p; p)$  by carrying  $(P, Q)$  into  $(-(P^{-1}Q)^T, I_p)$ ; it is differentiable ( $\tau$  denotes transpose). The row space of  $(P, Q)$  is the same as

that of  $(I_n, P^{-1}Q)$ , while the row vectors of this matrix are orthogonal to those of  $(-(P^{-1}Q)^T, I_p)$  (multiply the one by the transpose of the other). Hence  $g$  induces  $h|_{\rho(U)}$ , so that the latter is differentiable.

2.26 Definition. Let  $E(\gamma_p^n)$  be defined as that subset of  $G_{p,n} \times R^{n+p}$  consisting of pairs  $(H, x)$  where  $x$  is a vector lying in the hyperplane  $H$ . It is called the universal bundle (for reasons we shall see). The projection  $\pi$  maps  $(H, x)$  into  $H$ ; the fibre is thus an  $n$ -dimensional subspace of  $R^{n+p}$ .

$\gamma_p^n$  is an  $n$ -dimensional vector space bundle over  $G_{p,n}$ . We need to show the existence of a local product structure. Let  $(\rho(U), \varphi_0)$  be a coordinate neighborhood on  $G_{p,n}$ , as in (a) above. We define  $h: \rho(U) \times R^n \rightarrow \pi^{-1}\rho(U)$  as carrying  $(H, (x^1, \dots, x^n))$  into  $(x^1, \dots, x^n) \circ (I_n, Q)$  where  $Q = \varphi_0(H)$ . This is a vector in the hyperplane  $H$ ;  $h$  is clearly an isomorphism on each fibre. Its inverse is continuous, since it sends  $(H, (y^1, \dots, y^{n+p}))$  in  $G_{p,n} \times R^{n+p}$  into  $(H, (y^1, \dots, y^n))$  in  $\rho(U) \times R^n$ .

2.27 Definition.  $\xi$  is a differentiable vector space bundle if  $E(\xi)$  and  $B(\xi)$  are differentiable manifolds, and if the homeomorphisms

$$U \times \mathbb{R}^n \longrightarrow \pi^{-1}(U)$$

which specify the local product structure can be chosen as diffeomorphisms.

It follows that  $\pi : E \longrightarrow B$  is differentiable of maximum rank. Note that  $B$  can be differentiably imbedded in  $E$  by mapping  $b$  into the 0-vector of  $F_b$ . The normal bundle of this imbedding is just  $\xi$ .

Examples of differentiable bundles include the tangent bundle of a manifold, the normal bundle of an immersed manifold, and the universal bundle  $\gamma_p^n$  above. In the latter case,  $E(\gamma_p^n)$  is imbedded differentiably in  $G_{p,n} \times \mathbb{R}^{n+p}$ .

2.28 Theorem. Let  $\xi^n$  be an  $n$ -dimensional vector space bundle. The following conditions are equivalent:

- (a)  $\xi$  is of finite type.
- (b) There is a bundle  $\eta^p$  such that  $\xi^n \oplus \eta^p$  is trivial.
- (c) There is a bundle map  $\xi^n \longrightarrow \gamma_p^n$  for some  $p$ . (Thus the terminology "universal bundle" for  $\gamma_p^n$ .)

Proof: We have already shown that (a) implies (b) (2.20); the bundle  $\eta^p$  there constructed has dimension  $n(k-1)$ , where  $k$  is the number of elements in the covering  $U_1, \dots, U_k$  of  $B(\xi) = B$  such that  $\xi|_{U_i}$  is trivial.

(b) implies (c): Condition (b) means that  $\xi^n$  may be imbedded in the trivial bundle  $B(\xi) \times R^{n+p}$ ; let  $f$  be this imbedding. We wish to define  $g$  and  $g_B$  in the following diagram:

$$\begin{array}{ccc} E(\xi) & \xrightarrow{g} & E(\gamma_p^n) \\ \pi \downarrow & & \downarrow \\ B(\xi) & \xrightarrow{g_B} & G_{pn} \end{array}$$

Since  $f$  is a 1-1 homomorphism,  $f(F_b)$  is the cartesian product of  $b$  and an  $n$ -dim hyperplane  $H^n$  in  $R^{n+p}$ ; let  $g_B(b)$  equal this hyperplane  $H^n$ . If  $e \in F_b$ , then  $f(e) = (b, x)$ , where  $x$  is a vector in the hyperplane  $H^n$ ; let  $g(e) = (H^n, x)$  in  $G_{pn} \times R^{n+p}$ . Then  $g(e)$  actually lies in the subset of  $G_{pn} \times R^{n+p}$  which constitutes  $E(\gamma_p^n)$ . From rank considerations,  $g$  is automatically an isomorphism on each fibre.

It remains to show that  $g$  is continuous. Locally,  $g$  just looks like a map  $U \times R^n \longrightarrow G_{pn} \times R^{n+p}$ . We factor it into a continuous map  $h: U \times R^n \longrightarrow M(n, n+p; n) \times R^{n+p}$  followed by the projection  $\rho \times 1$  into  $G_{pn} \times R^{n+p}$ . Locally,  $f$  looks like a map  $U \times R^n \longrightarrow B \times R^{n+p}$ . Let  $e_1, \dots, e_n$  be a basis for  $R^n$ ; we define  $h(b, x)$  as  $(A, p_2 f(b, x))$ . Here  $p_2$  projects  $B \times R^{n+p}$  onto its second factor and  $A$

is the matrix having  $p_2 f(b, e_1), \dots, p_2 f(b, e_n)$  as its rows. Then  $h$  is continuous, and  $(\rho \times 1)h$  equals  $g$ .

(Note: The converse assertion, (c) implies (b), can be proved by the same argument.)

(c) implies (a): Being compact,  $G_{pn}$  is covered by finitely many neighborhoods  $U_i$  with  $\gamma_p^n|_{U_i}$  trivial. (In fact  $(n+p)!/n!p!$  neighborhoods will suffice.) If  $f$  is a bundle map  $\xi^n \rightarrow \gamma_p^n$  then the sets  $f_B^{-1}(U_i) = V_i$  cover  $B$ , and  $\xi|_{V_i}$  is equivalent to the bundle induced by  $f_B: V_i \rightarrow G_{pn}$  (the uniqueness part of 2.8). Then  $\xi|_{V_i}$  is trivial (since it is induced from a trivial bundle).

## Chapter III

The Cobordism Theory of Thom

3.1 Definition. An n-manifold-with-boundary  $Q$  is a Hausdorff space with a countable basis which is locally homeomorphic with  $H^n$  (the subset of  $R^n$  such that  $x^1 \geq 0$ ). The boundary  $\partial Q$  is that subset of  $Q$  corresponding to  $R^{n-1}$  under the local homeomorphism ( $R^{n-1}$  being the subset of  $R^n$  with  $x^1 = 0$ ).  $\partial Q$  is well-defined, since the image of an open set in  $R^n$  under a homeomorphism of it into  $R^n$  must be open (Brouwer theorem on invariance of domain). It is clear that  $\partial Q$  is an  $(n-1)$ -manifold.

A differentiable structure  $\mathcal{D}$  on  $Q$  is a collection of real-valued functions  $f$  defined on open subsets of  $Q$  such that

1) every point of  $Q$  has an open neighborhood  $U$  and a homeomorphism  $h$  of  $U$  into an open subset of  $H^n$ , such that  $f$  is in  $\mathcal{D}$  if and only if  $fh^{-1}$  is differentiable. ( $f$  is defined on an open subset of  $U$ ;  $fh^{-1}$  differentiable means that it may be extended to a neighborhood of  $h(U)$  in  $R^n$  so as to be differentiable).

2) If  $U_i$  are open sets contained in the domain of  $f$  and  $U = \cup U_i$ , then  $f|_U \in \mathcal{D}$  if and only if  $f|_{U_i} \in \mathcal{D}$  for each  $i$ .

As before,  $(U, h)$  is called a coordinate system on  $Q$ , and one can define differentiable structures alternatively by means of coordinate systems.

We impose an additional condition on  $\mathcal{D}$  in 3.2.

**3.2 Definition.** Let  $M_1, M_2$  be compact differentiable  $n$ -manifolds. They are said to lie in the same cobordism class  $(M_1 \sim M_2)$  if there is a compact differentiable  $n+1$  manifold-with-boundary  $Q$  such that  $\partial Q$  is diffeomorphic with the disjoint union of  $M_1$  and  $M_2$  (denoted by  $M_1 + M_2$ ).

Symmetry and reflexivity of this relation are clear.

To show transitivity, we impose the additional condition on  $\mathcal{D}$  that there is a neighborhood  $U$  of  $\partial Q$  in  $Q$  which is diffeomorphic with  $\partial Q \times [0, 1)$ , the diffeomorphism being the identity on  $\partial Q \times 0$ . This is redundant, but we assume it to avoid proving it. Transitivity follows:

Let  $M_1 + M_2$  be diffeomorphic with  $\partial Q_1$  and  $M_2 + M_3$  diffeomorphic with  $\partial Q_2$ ; let  $h_1, h_2$  be the diffeomorphisms. We form a new space  $Q_3$  from  $Q_1 \cup Q_2$  by identifying each point of  $h_1(M_2)$  with its image under  $h_2 h_1^{-1}$ . There is then a homeomorphism of  $M_2 \times (-1, 1)$  into this space which equals  $h_1$  when restricted to  $M_2 \times 0$ , and is a diffeomorphism of  $M_2 \times [0, (-1)^1)$  into  $Q_1$  for  $i = 1, 2$ . (It is derived from the postulated "product neighborhoods"  $\partial Q_i \times [0, 1)$ .) If this



is taken to be a coordinate system on  $Q_3$ ,  $Q_3$  becomes a differentiable manifold-with-boundary, and  $M_1 + M_3$  is diffeomorphic with  $\partial Q_3$ .  $Q_1$  and  $Q_2$  are diffeomorphic with subsets of  $Q_3$ .

3.3 Definition. As usual, there are logical difficulties involved in considering these cobordism classes. One way of avoiding them is to consider only manifolds-with-boundary imbedded in some euclidean space  $R^p$ : If  $Q_1$  is a differentiable manifold-with-boundary and  $Q_2 = \partial Q_1 \times [0,1)$ , then the space  $Q_3$  constructed in the preceding paragraph is a differentiable manifold, so that it may be imbedded in some euclidean space. Hence  $Q_1$  may so be imbedded.

With these restrictions, the set of cobordism classes of  $n$ -manifolds forms an abelian group (denoted by  $\mathcal{C}^n$ ) under the operation  $+$  (disjoint union). If  $M_1 \sim M'_1$  and  $M_2 \sim M'_2$ , this means that  $M_1 + M'_1$  is diffeomorphic with  $\partial Q_1$ . Then  $(M_1 + M_2) + (M'_1 + M'_2)$  is diffeomorphic with  $\partial(Q_1 \cup Q_2)$ , so that  $M_1 + M_2 \sim M'_1 + M'_2$  and the operation  $+$  is well-defined on cobordism classes. The zero element is the vacuous manifold or the  $n$ -sphere (or  $\partial Q$ , where  $Q$  is any compact differentiable  $(n+1)$ -manifold-with-boundary). The remaining axioms are clear. Note that  $M + M$  is diffeomorphic with  $\partial(M \times [0,1])$ , so that every element is of order 2.

The groups  $\mathcal{U}^n$  are called the (non-orientable) cobordism groups. Let  $\mathcal{U}$  denote the direct sum  $\mathcal{U}^0 \oplus \mathcal{U}^1 \oplus \mathcal{U}^2 \oplus \dots$ . There is a bilinear symmetric pairing of  $\mathcal{U}^i, \mathcal{U}^j$  into  $\mathcal{U}^{i+j}$ , i.e. a homomorphism of  $\mathcal{U}^i \otimes \mathcal{U}^j$  into  $\mathcal{U}^{i+j}$  induced by the operation of cartesian product.

First,  $(M_1 + M_2) \times M_3 = (M_1 \times M_3) + (M_2 \times M_3)$  by definition of cartesian product. Second, if  $M_1 \sim 0$ , i.e.  $M_1 = \partial Q$ , then  $M_1 \times M_2$  is diffeomorphic with  $\partial(Q \times M_2)$ , so that  $M_1 \times M_2 \sim 0$ .

Since  $M_1 \times M_2 \sim M_2 \times M_1$ , and since  $M_1 \times p \sim M_1$  (where  $p$  is a point-manifold), this pairing makes  $\mathcal{U}$  into a (graded) commutative ring with unit. Indeed, it is a graded algebra over the field  $Z_2$ .

3.4 Remark. The general result of Thom is the following

Theorem.  $\mathcal{U}$  is a polynomial algebra over  $Z_2$  with one generator in each positive dimension except those of the form  $2^m - 1$ . If  $n$  is even, projective  $n$ -space is a generator.

This theorem means that there are compact manifolds  $M^2, M^4, M^5, \dots$  such that every compact manifold is in the cobordism class of a disjoint union of products of these manifolds, and that there are no relations among the generators (except commutativity and associativity of products).

Thom's procedure is to show that  $\mathcal{V}^n$  is isomorphic with the  $(n+k)$ <sup>th</sup> homotopy group of a certain space  $T_k$ ; and then to compute these homotopy groups. We shall consider only the first of these two problems in the present notes.

**3.5 Definition.** Let  $h$  be an imbedding of the differentiable manifold  $M^n$  in  $R^{n+k}$ ; consider the normal bundle of this imbedding. Using the standard Riemannian metric for the tangent bundle to  $R^{n+k}$ , this normal bundle is equivalent to the orthogonal complement of the image in the tangent bundle of  $R^{n+k}$  of the tangent bundle of  $M^n$  (2.16); this complement we denote by  $\nu^k$ . Define  $e$  as the canonical map of  $E(\nu^k)$  into  $R^{n+k}$  which maps the vector  $v$  normal to  $M^n$  at  $x$  into its end point. (Described differently, one maps the tangent bundle to  $R^{n+k}$  into itself canonically by mapping the vector  $v$ , based at  $x$ , into the point  $x + v$  of  $R^{n+k}$ . This map is differentiable; its restriction to  $E(\nu^k)$  is the map  $e$ .)

Consider  $M^n$  as the zero vectors of  $E(\nu^k)$ . Then we have the

**3.6 Theorem.** There is a neighborhood of  $M^n$  in  $E(\nu^k)$  which is mapped diffeomorphically onto a neighborhood of  $M^n$  in  $R^{n+k}$ .

Proof: Note that  $e$  is differentiable, and that it has rank  $n+k$  at points of  $M^n \subset E(v^k)$ . (This is easily checked by computing the derivative matrix of  $e$  with respect to a local coordinate system.) Hence  $e$  has rank  $n+k$  in some neighborhood of  $M^n$  in  $E(v^k)$ , so that it is a local homeomorphism at points of  $M^n$ : it maps a neighborhood of each  $x \in M^n$  homeomorphically onto a neighborhood of  $f(x)$ . We then appeal to the topological lemma:

If  $f: X \rightarrow Y$  is a local homeomorphism and the restriction of  $f$  to the closed subset  $A$  is a homeomorphism, then  $f$  is a homeomorphism on some neighborhood  $V$  of  $A$ . ( $X, Y$  are Hausdorff spaces with countable bases;  $X$  is locally compact.) This lemma is proved as follows:

(1) If  $A$  is compact, the lemma holds. For otherwise, there would be points  $x, y$  arbitrarily close to  $A$  such that  $f(x) = f(y)$ . Since  $A$  has a compact neighborhood, we may choose sequences  $x_n, y_n$  converging to  $x, y$ , respectively, in  $A$  such that  $x_n \neq y_n$  and  $f(x_n) = f(y_n)$ . Hence  $f(x) = f(y)$  so that  $x = y$ ,  $f$  being a homeomorphism on  $A$ . But then  $f$  is not a local homeomorphism at  $x$ .

(2) Let  $A_0$  be a compact subset of  $A$ . Then there is a neighborhood  $U_0$  of  $A_0$  such that  $\bar{U}_0$  is compact and  $f$  is a homeomorphism on  $\bar{U}_0 \cup A$ : It will suffice for  $f$  to be 1-1,

since  $f$  is a local homeomorphism. By (1), let  $V_0$  be a neighborhood of  $A_0$  so that  $f|_{\bar{V}_0}$  is 1-1. If no neighborhood of  $A_0$  in  $V_0$  satisfies the requirements for  $U_0$ , there is a sequence of points  $x_n$  of  $X - A$  converging to  $x \in A_0$  with  $f(x_n) \in f(A)$ . Choose  $y_n \in A$  with  $f(x_n) = f(y_n)$ . Since  $f$  is continuous,  $f(y_n)$  converges to  $f(x)$ ; since  $f$  is a homeomorphism on  $A$ ,  $y_n$  converges to  $x$ . Since  $x_n \neq y_n$ , this contradicts the fact that  $f$  is a local homeomorphism at  $x$ .

(3) Express  $A$  as the union of an ascending sequence of compact sets  $A_1 \subset A_2 \subset \dots$ . Let  $V_1$  be a neighborhood of  $A_1$  such that  $\bar{V}_1$  is compact and  $f$  is a homeomorphism on  $\bar{V}_1 \cup A$  (by (2)). Given  $V_1$  a neighborhood of  $A_1$  satisfying these conditions, consider the set  $\bar{V}_1 \cup A_{i+1}$ . It is a compact subset of  $\bar{V}_1 \cup A$ , and  $f$  is a homeomorphism on  $\bar{V}_1 \cup A$ . Hence by (2) there is a neighborhood  $V_{i+1}$  of  $\bar{V}_1 \cup A_{i+1}$  with  $\bar{V}_{i+1}$  compact, such that  $f$  is a homeomorphism on  $\bar{V}_{i+1} \cup A$ . We proceed by induction.  $f$  is 1-1 on  $V = \cup V_{i+1}$ , so that it is a homeomorphism on  $V$  (being a local homeomorphism -onto).

**3.7 Corollary.** Any differentiable submanifold of  $\mathbb{R}^{n+k}$  is a differentiable neighborhood retract.

The projection of  $E(v^k) \rightarrow M^n$  induces (under  $e$ ) a differentiable map of a neighborhood of  $M^n$  in  $R^{n+k}$  onto  $M^n$  which is the identity on  $M^n$ .

**3.8 Definition.** Let  $\xi$  be a vector space bundle with  $B(\xi)$  compact; let  $T(\xi)$  denote the 1-point compactification of  $E(\xi)$ . It is called the Thom space of  $\xi$ . Let  $\infty$  denote the added point.

Let  $\xi$  have a Riemannian metric. Let  $T_\epsilon(\xi)$  be obtained from  $E(\xi)$  by identifying all vectors of length greater than or equal to  $\epsilon$  to a point. Let  $\alpha(x)$  be a  $C^\infty$  function with  $\alpha'(x) \geq 0$  which equals 1 in a neighborhood of  $x = 0$  and  $\rightarrow \infty$  as  $x \rightarrow 1$ . The map of  $E(\xi)$  into  $T(\xi)$  which carries the vector  $e$  into the vector  $e \alpha(\|e\|/\epsilon)$  induces a homeomorphism of  $T_\epsilon(\xi)$  onto  $T(\xi)$  which is a diffeomorphism on the set  $E_\epsilon(\xi)$ , consisting of vectors of length less than  $\epsilon$ . The fact that  $B$  is compact is used here.

**3.9 Definition.** Let the compact manifold  $M^n$  be imbedded in  $R^{n+k}$ .  $v^k$  is given the Riemannian metric of  $R^{n+k}$ ; by 3.6 there is a neighborhood of  $M^n$  in  $R^{n+k}$  which is diffeomorphic to the subset  $E_{2\epsilon}(v^k)$  of  $E(v^k)$ . Such a neighborhood is called a tubular neighborhood of  $M^n$ .

By 3.8, we see that  $T(v^k)$  is homeomorphic with the space obtained from  $R^{n+k}$  by collapsing the exterior of the tubular  $\epsilon$ -neighborhood of  $M$  to a point.

We will need three lemmas concerning approximation by differentiable functions.

**3.10 Lemma.** Let  $A$  be a closed subset of the differentiable manifold  $M$ , let  $f: M \rightarrow R^m$  be differentiable on  $A$ . Let  $\delta$  be a positive continuous function on  $M$ . There exists  $g: M \rightarrow R^m$  such that

- (1)  $g$  is differentiable
- (2)  $g$  is a  $\delta$ -approximation to  $f$
- (3)  $g|_A = f|_A$ .

Proof: It suffices to prove this lemma in the case  $m = 1$ .

Given  $x \in A$ ,  $f|_A$  may be extended to a differentiable function  $f_x$  in a neighborhood  $N_x$  of  $x$ . Let  $N_x$  be chosen small enough that  $|f_x(y) - f(y)| < \delta(y)$  for all  $y \in N_x$ .

Given  $x \in M - A$ , choose a neighborhood  $N_x$  of  $x$  small enough that  $|f(y) - f(x)| < \delta(y)$  for all  $y \in N_x$ . Define  $f_x(y) \equiv f(x)$  for  $y \in N_x$ .

Let  $\varphi_\alpha$  be a differentiable partition of unity with carrier  $\varphi_\alpha$  contained in some  $N_x$ , say  $N_{x(\alpha)}$ , for each  $\alpha$ .

Define  $g(y) = \sum \varphi_\alpha(y) f_{x(\alpha)}(y)$ . One checks the conditions of the lemma easily.

More generally:

**3.11 Lemma.** Let  $f: M_1 \longrightarrow M_2$  be a continuous map of differentiable manifolds which is differentiable on the closed subset  $A$  of  $M_1$ . Let  $\varepsilon(x) > 0$  be given; and give  $M_2$  the metric determined by some imbedding  $M_2 \subset \mathbb{R}^p$ . Then there exists  $g: M_1 \longrightarrow M_2$  such that

- (1)  $g$  is differentiable
- (2)  $g$  is an  $\varepsilon$ -approximation to  $f$
- (3)  $g|_A = f|_A$ .

Proof: There is a neighborhood  $U$  of  $M_2$  in  $\mathbb{R}^p$  of which  $M_2$  is a differentiable retract (3.7). Let  $\rho$  be the differentiable retraction of  $U$  onto  $M_2$ . Let  $\delta(x)$  be a positive function on  $M_2$  so chosen that the cubical neighborhood of  $f(x)$  of radius  $\delta(x)$  lies in  $U$ , and so that its image under  $\rho$  has radius less than  $\varepsilon(x)$ . Let  $f_1: M_1 \longrightarrow \mathbb{R}^p$  be a differentiable map which is a  $\delta$ -approximation to  $f$ , such that  $f_1|_A = f|_A$  (by 3.10). Define  $g(x) = \rho(f_1(x))$ .



3.12 Lemma. Let  $f: M_1 \longrightarrow M_2$  be a continuous map of differentiable manifolds; let the metric on  $M_2$  be obtained by imbedding it in some euclidean space. Given  $\varepsilon(x)$ , there is a  $\delta(x)$  such that if  $g: M_1 \longrightarrow M_2$  is a  $\delta$ -approximation to  $f$ ,  $g$  is homotopic to  $f$  under a homotopy  $F(x,t)$  with

- (1)  $F(x,t) = f(x)$  for any  $x$  such that  $g(x) = f(x)$  and
- (2)  $F(x,t)$  is an  $\varepsilon$ -approximation to  $f$  for any  $t$ .

Proof: Let  $U, \rho$ , and  $\delta(x)$  be chosen as in 3.11.

Let  $g: M_1 \longrightarrow M_2$  be a  $\delta$ -approximation to  $f$ .

Then the line segment from  $g(x)$  to  $f(x)$  lies in  $U$ , so that  $F(x,t) = \rho(tg(x) + (1-t)f(x))$  is well-defined. Furthermore  $F(x,t)$  is an  $\varepsilon$ -approximation to  $f(x)$  for any  $t$ .

3.13 Definition. Let  $\xi^k$  be a differentiable vector space bundle with  $B(\xi)$  compact and  $m$ -dimensional; let  $E(\xi^k)$  be given a metric by imbedding it as a closed differentiable submanifold in some euclidean space (it is an  $(m+k)$ -manifold).

Given an element of  $\pi_{n+k}(T(\xi^k), \infty)$ , let it be represented by the map

$$f: (\bar{C}_{n+k}, \partial \bar{C}_{n+k}) \longrightarrow (T(\xi^k), \infty),$$

where  $\bar{C}_{n+k}$  is the closed cube  $[0,1]^{n+k}$  and  $\partial \bar{C}_{n+k}$  is its boundary.

Let  $U$  denote the open subset  $f^{-1}(E(\xi^k))$  of  $C_{n+k}$ . Let  $g: U \rightarrow E(\xi^k)$  be a differentiable  $\delta$ -approximation to  $f|U$ , where  $\delta$  is so chosen that  $\delta < 1$  and  $g$  is homotopic to  $f$ , the homotopy  $F$  also being a 1-approximation to  $f$ . (This ensures that  $F$  will be continuous if we define  $F(x,t) \equiv \infty$  for  $x \in \bar{C}_{n+k} - U$ .)

Now  $g$  may be approximated in turn by a differentiable map  $h: U \rightarrow E(\xi)$  which is transverse regular on the submanifold  $B(\xi)$  of  $E(\xi)$ . We choose the approximation close enough that  $g$  is homotopic to  $h$ , the homotopy  $H$  being a 1-approximation to  $g$  for each  $t$ . Extend  $h$  to  $\bar{C}_{n+k}$  by defining  $h(x) = \infty$  for  $x \in \bar{C}_{n+k} - U$ . Then  $h$  is in the homotopy class of  $f$ .

$h^{-1}(B(\xi))$  is a differentiable submanifold  $M^n$  of  $U$  which is closed in  $\bar{C}_{n+k}$ , and thus compact.

Definition: Let  $\lambda$  assign to the homotopy class of  $h$  in  $\pi_{n+k}(T(\xi^k), \infty)$  the cobordism class of  $M^n$  in  $\mathcal{U}^n$ .

3.14 Theorem.  $\lambda$  is a well-defined homomorphism.

Proof: Let  $H: (\bar{C}_{n+k} \times I, \partial \bar{C}_{n+k} \times I) \rightarrow (T(\xi^k), \infty)$   
 be a homotopy between  $h_0 = H(x, 0)$  and  $h_1 = H(x, 1)$ . Let  
 $\checkmark h_0, h_1$  satisfy the conditions

- (1)  $h_1$  is differentiable on  $h_1^{-1}(E(\xi))$
- (2)  $h_1$  is transverse regular on  $B(\xi)$ . ( $i=1, 2$ .)

We wish to show that  $\checkmark h_0^{-1}(B)$  and  $h_1^{-1}(B)$  belong to the same  
 cobordism class.

We may assume that  $H(x, t) = H(x, 0)$  for  $t \leq 1/3$ , and  
 $H(x, t) = H(x, 1)$  for  $t \geq 2/3$ . Let  
 $U = H^{-1}(E(\xi)) \cap [\bar{C}_{n+k} \times (0, 1)]$ ; then  $U$  is an open subset of  
 $R^{n+k+1}$ . Let  $G: U \rightarrow E(\xi)$  be a differentiable 1-approximation  
 to  $H$  which equals  $H$  on the closed subset  $A$ , where  
 $A = U \cap [\bar{C}_{n+k} \times ((0, 1/4] \cup [3/4, 1))]$ . (see 3.11.  $H$  is dif-  
 ferentiable on  $A$ .)

Now  $G$  satisfies the transverse regularity condition  
 for  $B(\xi)$  at points in  $A$  (since  $h_0$  and  $h_1$  are transverse  
 regular on  $B$ ) so that by 1.35 there is a differentiable map  
 $F: U \rightarrow E(\xi)$  which equals  $G$  on  $A$ , is transverse regular  
 on  $B(\xi)$ , and is a 1-approximation to  $G$ . Because  $F$  is a  
 2-approximation to  $H$ , it remains continuous if we define  
 $F(x, t) = \infty$  for  $(x, t) \in (\bar{C}_{n+k} \times (0, 1)) - U$ . Because  $F$   
 equals  $H$  on  $A$ , it remains continuous if we define

$F(x,t) = H(x,t)$  for  $t = 0,1$ . Hence  $F^{-1}(B)$  is a compact subset of  $\bar{C}_{n+k}$ , being closed and bounded.

Because  $F|U$  is transverse regular on  $B$ ,  $(F|U)^{-1}(B)$  is a differentiable  $(n+1)$ -submanifold of  $\bar{C}_{n+k} \times (0,1)$ . Its intersection with  $\bar{C}_{n+k} \times t$  equals  $h_0^{-1}(B) \times t$  for  $t \leq 1/4$  and  $h_1^{-1}(B) \times t$  for  $t \geq 3/4$ . Hence  $F^{-1}(B)$  is a differentiable manifold-with-boundary whose boundary is  $h_0^{-1}(B) + h_1^{-1}(B)$ . Thus  $\lambda$  is well-defined.

It is trivial to show  $\lambda$  is a homomorphism, because the sum in  $\mathcal{U}^n$  is derived from disjoint unions of representative manifolds.

**3.15 Theorem.** If  $\xi^k$  is the universal bundle  $\gamma_m^k$  where  $k \geq n+1$ ,  $m \geq n$  then  $\lambda: \pi_{n+k}(\mathbb{T}(\gamma_m^k), \infty) \rightarrow \mathcal{U}^n$  is onto.

Proof: Let  $M^n$  be a compact  $n$ -manifold; let  $k \geq n+1$ . Let  $M^n$  be imbedded in  $C_{n+k}$  (1.32); let  $v^k$  be the normal bundle of this imbedding. The Riemannian metric on  $E(v^k)$  is that derived from the natural scalar product on the tangent bundle to  $R_{n+k}$ , in which  $v^k$  is contained.

By 3.6, for small  $\epsilon$  the subset of  $E_{2\epsilon}(v^k)$  of  $E(v^k)$  is diffeomorphic with a tubular neighborhood of  $M^n$  in  $C_{n+k}$ ; let  $U$  be the image of  $E_\epsilon(v^k)$ .

Let  $p_1$  project  $\bar{C}_{n+k}$  onto the space obtained from  $\bar{C}_{n+k}$  by identifying  $\bar{C}_{n+k} - U$  to a point (denoted by  $\bar{C}_{n+k}/\bar{C}_{n+k} - U$ ).

Let  $p_2$  be the diffeomorphism of  $U$  onto  $E_\epsilon(v^k)$ , followed by the map of  $E(v^k)$  into  $T_\epsilon(v^k)$  which identifies all vectors of length  $\geq \epsilon$  (3.8).  $p_2$  is then extended by mapping  $\bar{C}_{n+k} - U$  into  $\infty$ .

Let  $p_3$  be the homeomorphism of  $T_\epsilon(v^k)$  onto  $T(v^k)$  constructed in 3.8. The composite map  $p_3 p_2 p_1$  is a diffeomorphism of  $U$  onto  $E(v^k)$ .

Finally, let  $p_4$  be the bundle map of  $v^k$  into  $\gamma_m^k$  induced from the imbedding of  $M^n$  in  $R_{n+k} \subset R_{m+k}$ . Because both fibres have dimension  $k$ , this map satisfies the transverse regularity condition for  $G_{k m}$  at each point of  $M^n$ . Extend  $p_4$  in the obvious way to map  $T(v^k)$  into  $T(\gamma_m^k)$ .

Let  $g = p_4 p_3 p_2 p_1$ . Then  $g: \partial \bar{C} \rightarrow \infty$ . Let  $\mu(M^n)$  denote the homotopy class of  $g$  in  $\pi_{n+k}(T(\gamma_m^k), \infty)$ . Now  $g$  is transverse regular on  $G_{k m}$  and  $M^n = g^{-1}(G_{k m})$ . By definition, the cobordism class of  $M^n$  is the image of  $\mu(M^n)$  under  $\lambda$ , so that  $\lambda \mu(M^n) = [M^n]$ .

3.16 Theorem. If  $\xi^k$  is the universal bundle  $\gamma_m^k$  with  $k \geq n+2$ ,  $m > n$  then  $\lambda$  is one-to-one.

Proof: Given an element of  $\pi_{n+k}(\mathbb{T}(\gamma_m^k), \infty)$ , we may suppose it represented by a map

$$f: (\bar{C}_{n+k}, \partial \bar{C}_{n+k}) \longrightarrow (\mathbb{T}(\gamma_m^k), \infty)$$


which is differentiable on  $f^{-1}(E)$  and transverse regular on  $G_{m k}$  (by 3.13). Let  $M^n = f^{-1}(G_{m k})$ ; we wish to show that if  $M^n$  is the boundary of an  $(n+1)$ -manifold-with-boundary  $Q$ , then  $f$  is homotopic to the constant map.

$M^n$  is a submanifold of  $C_{n+k}$ ; let its normal bundle be  $\nu^k$ . Let  $\epsilon$  be chosen so that  $E_{2\epsilon}(\nu^k)$  is diffeomorphic with the  $2\epsilon$ -neighborhood of  $M^n$ ; let  $U_\epsilon$  be the image of the vectors of  $E_\epsilon(\nu^k)$ . Impose a Riemannian metric on  $\gamma_m^k$ ; let  $\delta$  be so chosen that  $\|x\| \geq \epsilon$  implies  $\|f(x)\| \geq \delta$  for  $x \in E(\nu^k)$ .

Step 1.  $f$  is homotopic to a map  $f_1$  such that

- (1)  $f_1$  is differentiable on  $f_1^{-1}(E)$  and transverse regular on  $G_{m k}$ .
- (2)  $f = f_1$  on  $M^n = f_1^{-1}(G_{m k})$ .
- (3)  $f_1$  carries everything outside  $U_\epsilon$  into  $\infty$ .

$\alpha = 1$  in nbh of 0



Define  $F: E(\gamma_m^k) \times I \rightarrow T(\gamma_m^k)$  by the equation

$F(e,t) = e \alpha(t\|e\|/\delta)$ , where  $\alpha$  is the function defined in

3.8. Let  $f_1(x) = F(f(x),1)$ .

Step 2. By the diffeomorphism of  $U_{2\epsilon}$  with  $E_{2\epsilon}$ ,  $f_1$  induces a map  $\bar{f}_1$  of  $\bar{E}_\epsilon(v^k)$  into  $T(\gamma_m^k)$  which carries  $\partial(E_\epsilon)$  into  $\infty$ . Any homotopy of  $\bar{f}_1$  which leaves  $\partial(E_\epsilon)$  at  $\infty$  induces a homotopy of  $f_1$ .

Now  $\bar{f}_1$  is homotopic to a map  $\bar{f}_2$  such that

- (1)  $\bar{f}_2$  is differentiable on  $\bar{f}_2^{-1}(E)$  and transverse regular on  $G_{mk}$ .
- (2)  $\bar{f}_2 = \bar{f}_1$  on  $M^n = \bar{f}_2^{-1}(G_{mk})$ .
- (3)  $\bar{f}_2$  is locally a bundle map in some neighborhood of  $M^n$ .

The homotopy leaves  $\partial(E_\epsilon)$  at  $\infty$ .

Consider  $G: \bar{E}_\epsilon(v^k) \times I \rightarrow T(\gamma_m^k)$  defined by the equation  $G(e,t) = \bar{f}_1(te)/t$ . As  $t \rightarrow 0$ ,  $G(e,t)$  approaches a limit which is non-zero if  $e \neq 0$  (since  $\bar{f}_1$  is differentiable and  $t$ -regular). It is easily seen to be a bundle map. It will not suffice for our purposes, since it does not carry  $\partial(E_\epsilon) \times I$  into  $\infty$ . Choose  $\delta > 0$  so that  $\|x\| \geq \epsilon$  implies  $\|G(x,t)\| \geq \delta$  for  $x \in E(v^k)$ ,  $t \in I$ , and define  $H(e,t) = [G(e,t)] \alpha(\|G(e,t)\|/\delta)$

If we set  $\bar{f}_2 = H(e, 0)$ , then  $\bar{f}_2$  is a bundle map for  $\|e\|$  small (since  $\alpha(x) \equiv 1$  for  $x$  small). The map  $H(e, 1) = \bar{f}_1(e) \alpha(\|\bar{f}_1(e)\|/\delta)$  does not equal  $\bar{f}_1$ , but it is homotopic to  $\bar{f}_1$ , the homotopy leaving  $\partial(E_\epsilon)$  at  $\infty$ . The homotopy is defined by the equation

$$K(e, t) = \bar{f}_1(e) \alpha(t\|\bar{f}_1(e)\|/\delta), \text{ as in step 1.}$$

Step 3. Let  $Q$  be the  $n+1$  manifold-with-boundary such that  $M^n = \partial Q$ . Let  $h$  be a diffeomorphism of  $M^n \times [0, 1]$  into  $Q$  which carries  $M^n \times 0$  onto  $\partial Q$ .

Define  $h_1: Q \rightarrow C_{n+k} \times I$  as follows:

If  $x = h(y, t)$  where  $y \in M^n$  and  $0 \leq t \leq 1/2$ , let  $h_1(x) = (y, t)$ .

If  $x \notin \text{image } h$ , let  $h_1(x) = p$ , where  $p$  is some fixed point interior to  $C_{n+k} \times I$ .

If  $x = h(y, t)$  where  $y \in M^n$  and  $1/2 \leq t \leq 1$ , let

$$h_1(x) = (1 - \beta(t)) h_1(y, 1/2) + \beta(t)p, \text{ where}$$

$\beta(t)$  is a  $C^\infty$  function with  $\beta'(t) \geq 0$ ,  $\beta(t) \equiv 0$  in a neighborhood of  $t = 1/2$  and  $\beta(t) \equiv 1$  in a neighborhood of  $t = 1$ .

$h_1$  is a differentiable map of  $\text{Int } Q$  into  $\text{Int}(C_{n+k} \times I)$ ; and  $h_1$  is a 1-1 immersion in a neighborhood of  $\partial Q$ .



Since  $\dim(C_{n+k} \times I) > 2(n+1)$ ,  $h_1$  may be approximated by a 1-1 immersion  $h_2$  which equals  $h_1$  in a neighborhood of  $\partial Q$  (by 1.29). It may be extended to an imbedding of  $Q$  into  $C_{n+k} \times I$ . (Since  $Q$  is compact, a 1-1 immersion is automatically an imbedding.) Let  $Q$  now be considered as this subset of  $C_{n+k} \times I$ .

Step 4. We have a map  $f_2$  of  $\bar{C}_{n+k} \times 0$  into  $T(\gamma_m^k)$  which is a bundle map when restricted to a small tubular neighborhood of  $M^n \times 0$  in  $C_{n+k} \times 0$ . We extend it to  $\bar{C}_{n+k} \times [0, b)$  for  $b$  small in the trivial way. Suppose there exists a map  $g$  of the  $\epsilon'$ -neighborhood  $N$  of  $Q$  in  $C_{n+k} \times I$  into  $T(\gamma_m^k)$  which equals  $f_2$  in some neighborhood of  $\partial Q$  in  $C_{n+k} \times I$  and maps each point of  $N - Q$  into a non-zero vector in  $E(\gamma_m^k)$ . Our theorem then follows: Let  $\delta$  be so chosen that, if the distance of  $x$  from  $Q$  is  $\geq \epsilon/2$ , then  $\|g(x)\| \geq \delta$ .

Define  $g_1: C_{n+k} \times I \longrightarrow T(\gamma_m^k)$  by the equation

$$g_1(x, s) = g(x, s) \alpha (\|g(x, s)\|/\delta) \quad \text{for } (x, s) \in N, \text{ and}$$

$$g_1(x, s) = \infty \quad \text{otherwise.}$$

The restriction of  $g_1$  to  $C_{n+k} \times 0$  does not equal the map  $f_2$ ,

but it is homotopic to  $f_2$ , by the same technique as used at the end of Step 2.  $g_1$  is the homotopy required for our theorem.

To show that the extension  $g$  exists, we refer to Steenrod, *Fibre Bundles* (Princeton Press, 1951). According to §19.4 and 19.7 of this book, the principal bundle associated with  $\gamma_m^k$  is an  $m$ -universal bundle. That is: given a vector space bundle  $\xi^k$  over a complex of dimension  $\leq m$ , any bundle map of  $\xi^k$ , restricted to a subcomplex, into  $\gamma_m^k$  can be extended throughout  $\xi^k$ . We will assume the well known result that  $Q$  can be triangulated. The dimension  $n+1$  of  $Q$  is  $\leq m$ . Hence any bundle map of the normal bundle  $\nu^k$  of  $Q$ , restricted to a polyhedral neighborhood of  $\partial Q$ , into  $\gamma_m^k$  can be extended throughout  $\nu^k$ .

Applying this result to the map  $f_2$ , this completes the proof of 3.16.

Letting  $T_k$  stand for the union of the Thom spaces  $T(\gamma_m^k) \subset T(\gamma_{m+1}^k) \subset \dots$ , in the weak topology, Theorems 3.15 and 3.16 imply the following.

**3.17 Theorem.** The cobordism group  $\mathcal{U}^n$  is canonically isomorphic to the stable homotopy group  $\pi_{n+k}(T_k)$ , for  $k \geq n + 2$ .

References

- H. Whitney, Differentiable manifolds, Ann. of Math. vol. 37 (1936) pp. 645-680.
- The self-intersections of a smooth  $n$ -manifold in  $2n$ -space, Ann. of Math. Vol. 45 (1944) pp. 220-246.
- The singularities of a smooth  $n$ -manifold in  $(2n-1)$ -space, Ann. of Math. vol. 45 (1944) pp. 247-293.
- A function not constant on a connected critical set of points, Duke Math. J. vol. 1 (1935) pp. 514-517.
- A. P. Morse, The behavior of a function on its critical set, Ann. of Math. vol. 40 (1939) pp. 62-70.
- A. Sard, The measure of the critical values of differentiable maps, Bull. Am. Math. Soc. vol. 48 (1942) pp. 883-890.
- R. Thom, Quelques propriétés globales des variétés différentiables, Comment. Math. Helv. vol. 28 (1954) pp. 17-86.