



**THE BOOK WAS  
DRENCHED**

UNIVERSAL  
LIBRARY

**OU\_166424**

UNIVERSAL  
LIBRARY







**OSMANIA UNIVERSITY LIBRARY**

Call No. 513.8/E 237

Accession No. 14282

Author *Edge. W. L.*

Title *Theory of ruled surfaces. 193*

This book should be returned on or before the date last marked below.





THE THEORY  
OF  
RULED SURFACES

Cambridge University Press  
Fetter Lane, London

*New York*

*Bombay, Calcutta, Madras*

*Toronto*

Macmillan

*Tokyo*

Maruzen Company Ltd.

All rights reserved

THE THEORY  
OF  
RULED SURFACES

BY

W. L. EDGE

FELLOW OF  
TRINITY COLLEGE  
CAMBRIDGE

CAMBRIDGE  
AT THE UNIVERSITY PRESS

1931

**PRINTED IN GREAT BRITAIN**

# CONTENTS

PREFACE . . . . .	page ix
-------------------	---------

## CHAPTER I: INTRODUCTORY

### *Section I*

PRELIMINARIES . . . . .	1
Birational transformation, 3. Normal curves, 5. Ruled surfaces, 7. Correspondence formulae, 9. The valency of a correspondence, 10. Brill's formula, 12. Zeuthen's formula, 14. The genus of a simple curve on a ruled surface, 15. The genus of a curve on a quadric, 16. The ruled surface generated by a correspondence between two curves, 17.	

### *Section II*

THE REPRESENTATION OF A RULED SURFACE IN THREE DIMENSIONS AS A CURVE ON A QUADRIC $\Omega$ IN FIVE DIMENSIONS . . . . .	22
The six coordinates of a line, 22. Line geometry in [3] considered as the point geometry of a quadric primal in [5], 24. The planes on $\Omega$ , 26. The quadric point-cone in [4], 26. The representation of a ruled surface, 27. Torsal generators, 28. The double curve and bitangent developable, 28. Triple points and tritangent planes, 30. Cubic ruled surfaces, 32.	

### *Section III*

THE PROJECTION OF RULED SURFACES FROM HIGHER SPACE . . . . .	33
--	----

## CHAPTER II: QUARTIC RULED SURFACES

INTRODUCTORY . . . . .	45
------------------------	----

### *Section I*

RATIONAL QUARTIC RULED SURFACES CONSIDERED AS CURVES ON $\Omega$	48
The representation of the double curve and bitangent developable . . . . .	52

### *Section II*

RATIONAL QUARTIC RULED SURFACES CONSIDERED AS PROJECTIONS OF NORMAL SURFACES IN HIGHER SPACE . . . . .	55
The normal surface in [5] with directrix conics . . . . .	55
The normal surface in [5] with a directrix line . . . . .	59

*Section III*

ELLIPTIC QUARTIC RULED SURFACES . . . . .	page 60
---	---------

*Section IV*

ALGEBRAICAL RESULTS CONNECTED WITH QUARTIC RULED SURFACES	62
---	----

## CHAPTER III: QUINTIC RULED SURFACES

*Section I*

RATIONAL QUINTIC RULED SURFACES CONSIDERED AS CURVES ON $\Omega$	72
The rational quintic curves on a quadric $\Omega$ in [5] . . . . .	73
The general surface in [3] . . . . .	75
Further consideration of the surfaces whose generators do not belong to a linear complex . . . . .	77
Surfaces whose generators belong to a linear complex which is not special . . . . .	82
Surfaces with a directrix line which is not a generator . . . . .	93
Surfaces with a directrix line which is also a generator . . . . .	100
Surfaces whose generators belong to a linear congruence . . . . .	102

*Section II*

RATIONAL QUINTIC RULED SURFACES CONSIDERED AS PROJECTIONS OF NORMAL SURFACES IN HIGHER SPACE . . . . .	104
The general surface in [6] . . . . .	104
The surfaces in [3] derived by projection from the general surface in [6] . . . . .	110
The surfaces in [3] derived by projection from the surface in [6] with a directrix line . . . . .	123

*Section III*

QUINTIC RULED SURFACES WHICH ARE NOT RATIONAL . . . . .	126
Elliptic quintic ruled surfaces considered as curves on $\Omega$ . . . . .	126
Elliptic quintic ruled surfaces in [3] considered as projections of normal surfaces in higher space . . . . .	130
The quintic ruled surface of genus 2 . . . . .	135

CHAPTER IV: SEXTIC RULED SURFACES

*Section I*

RATIONAL SEXTIC RULED SURFACES . . . . .	<i>page</i> 139
The rational sextic curves which lie on quadrics . . . . .	139
Surfaces without either a directrix line or a multiple genera- tor . . . . .	141
Surfaces with a multiple generator but without a directrix line . . . . .	154
Surfaces with a directrix line which is not a generator . . . . .	167
Surfaces with a directrix line which is also a generator . . . . .	184
Surfaces whose generators belong to a linear congruence . . . . .	196

*Section II*

ELLIPTIC SEXTIC RULED SURFACES . . . . .	199
Elliptic sextic curves which lie on quadrics . . . . .	200
Surfaces whose generators do not belong to a linear com- plex . . . . .	202
Surfaces whose generators belong to a linear complex which is not special . . . . .	206
Surfaces with a directrix line which is not a generator . . . . .	209
Surfaces with a directrix line which is also a generator . . . . .	215
Surfaces whose generators belong to a linear congruence <sup>ft</sup> . . . . .	217
The normal elliptic ruled surfaces of a given order $n$ . . . . .	217
The normal surfaces in [5] with directrix cubic curves . . . . .	224
The normal surface in [5] with a double line . . . . .	240

*Section III*

SEXTIC RULED SURFACES WHICH ARE NEITHER RATIONAL NOR ELLIPTIC . . . . .	248
The normal sextic curve of genus 2 . . . . .	248
Sextic curves of genus 2 which lie on quadrics . . . . .	249
The sextic ruled surfaces in [3] whose plane sections are curves of genus 2 . . . . .	250
A sextic ruled surface in [4] whose prime sections are curves of genus 2, and the ruled surfaces in [3] derived from it by projection . . . . .	253
The sextic ruled surfaces of genus 2 which are normal in [3] . . . . .	256
The sextic ruled surfaces whose plane sections are of genus greater than 2 . . . . .	258

## CHAPTER V: DEVELOPABLE SURFACES

Introduction . . . . .	<i>page</i> 260
The developable surface of the fourth order . . . . .	266
The developable surface of the fifth order . . . . .	267
The projections of the developable formed by the tangents of a normal rational quartic curve . . . . .	270
The sections of the locus formed by the osculating planes of a normal rational quartic curve . . . . .	274
The developables of the sixth order considered as curves on $\Omega$ . . . . .	278
The classification of the developable surfaces . . . . .	284

## CHAPTER VI: SEXTIC RULED SURFACES (CONTINUED)

Further types of rational sextic ruled surfaces . . . . .	286
Sextic ruled surfaces with a triple curve . . . . .	295

TABLES SHEWING THE DIFFERENT TYPES OF RULED SURFACES IN [3] UP TO AND INCLUDING THOSE OF THE SIXTH ORDER . . . . .	302
--	-----

NOTE. The intersections of two curves on a ruled surface . . . . .	311
--	-----



## PREFACE

In this volume all the ruled surfaces in ordinary space of orders up to and including the sixth are studied and classified. Tables shewing the different types of surfaces are given towards the end of the book, and the tables for the surfaces of the fifth and sixth orders are here obtained for the first time.

It seems that the results so obtained are of great importance; but the incidental purpose which, it is hoped, may be served by the book is perhaps of still greater importance. For there exists at present no work, easily accessible to English readers, which tests the application of the general ideas here employed in anything like the same detail. One might mention especially the use of higher space and the principle of correspondence, and these two ideas are vital and fundamental in all modern algebraic geometry. It is hoped therefore that the book may be of use to a wide circle of readers.

I wish to express here my thanks to the staff of the University Press for their unfailing accuracy in the printing and for the ready courtesy with which they have accepted my suggestions.

Notwithstanding the large number of surfaces which are herein investigated, the book would be incomplete were I not to make an acknowledgment of my obligations to Mr White, of St John's College, and Professor Baker. Even those who have only a slight knowledge of the multifariousness of Mr White's mathematical public services will be surprised to learn that he found time not only to read the proof sheets but also to read through the whole of the manuscript, and I am very grateful to him for his criticisms and suggestions.

My gratitude to Professor Baker is something more than that of a student to his teacher. He it was who first suggested that I should undertake this work, and his encouragement has been given unsparingly—and effectively—in times of difficulty. I have derived great benefit not only from my personal conversations with him but also from attending his courses of lectures. I thank him for many things; but especially for his interest, which has never flagged, and for his trust, which has never wavered.

W. L. E.



# CHAPTER I

## INTRODUCTORY

### SECTION I

#### PRELIMINARIES

1. The system of points on a line is determined by two of them, any third point of the line being derivable from these two; the same line is equally well determined by any two of its points. Similarly, if three points are taken which are not on the same line they determine a plane, the same plane being equally well determined by any three non-collinear points of it. Proceeding in this way we say that  $n + 1$  independent points determine a linear space of  $n$  dimensions, the points being independent when they are such that no one of them belongs to the space of less than  $n$  dimensions determined by the others; the same space of  $n$  dimensions is equally well determined by any  $n + 1$  independent points belonging to it.

We shall use the symbols  $[n]$  and  $S_n$  to denote a space of  $n$  dimensions. In  $[n]$  two spaces  $[m]$  and  $[n - m]$  of complementary dimensions have, in general, one point in common and no more. A space  $[p]$  and a space  $[q]$  have, in general, no common points if  $p + q < n$ , while if  $p + q > n$  they have, in general, a common  $[p + q - n]$ . If they have in common a space  $[r]$  where  $r > p + q - n$ , then they are contained in a space  $[p + q - r]$  or  $[n - s]$ , where  $s = r - p - q + n$ . For example: two lines in ordinary space do not intersect in general; if they do so they lie in a plane. If we call the intersection of  $[p]$  and  $[q]$  their *meet* and the space of lowest dimension which contains them both their *join*, then the sum of the dimensions of the meet and the join is  $p + q$ .

2. Just as we can project, in ordinary space, on to a plane so we can project, in  $[n]$ , on to  $[n - 1]$ ; if  $O$  is the centre of projection and  $P$  any point of  $[n]$  the line  $OP$  meets  $[n - 1]$  in a point  $P_1$  which is the projection of  $P$ . We can then project again from a point  $O_1$  of  $[n - 1]$  on to a space  $[n - 2]$  in  $[n - 1]$ , the line  $O_1P_1$  meeting  $[n - 2]$  in a point  $P_2$ . The passage from  $P$  to  $P_2$  can, however, be carried out in one step, simply by joining  $P$  to the line  $OO_1$  by a plane and taking  $P_2$  as the intersection of the plane with  $[n - 2]$ . We thus speak of projecting the points of  $[n]$  from a line on to  $[n - 2]$ . Similarly, we can project from a plane on to an  $[n - 3]$ , from a solid on to an  $[n - 4]$ , and so on; the sum of the dimensions of the space which is the centre of projection and of the space on to which we are projecting being always  $n - 1$ .

3. Just as the order\* of a plane curve is the number of points in which it is met by a line, and the order of a twisted curve is the number of points in which it is met by a plane, so the order of a curve in [4] is the number of points in which it is met by a solid† and so on, the order of a curve in [n] being the number of points in which it is met by a space [n - 1] of complementary dimension. The order of a surface in [n] is the number of points in which it is met by a space [n - 2], just as the order of a surface in ordinary space is the number of points in which it is met by a line. It is here implied that the space [n - 2] has a general position in regard to the surface, otherwise it might meet it in a curve; a line in ordinary space may itself lie on a surface. Similarly the order of a locus of r dimensions is equal to the (finite) number of points in which it is met by a space [n - r] of complementary dimension and of general position. A locus of dimension r and order m will be denoted by a symbol  $M_r^m$  or  $V_r^m$ , and if  $r = n - 1$  the locus will be spoken of as a "primal." A space [n - 1] lying in [n] is called a "prime" of [n];

4. If we have a curve in ordinary space its chords fill up the space; there is a finite number of them passing through a point of general position. But in [4] the chords of a curve do not fill up the space; they form a locus of three dimensions whose order is the number of points in which it meets a line. If we have a system of coordinates in [4], say five homogeneous or four non-homogeneous coordinates, the locus is given by an equation in these coordinates, and the order of the locus is the order of this equation. In [n] the chords of a curve form a three-dimensional locus whose order is equal to the number of points in which it meets an [n - 3]. The chords of a surface form a five-dimensional locus.

5. Suppose that we have a curve of order N in [n]; there may be a point of the curve such that any [n - 1] passing through it only meets the curve in N - 2 other points. Such a point is called a double point of the curve. In particular we have the double points of a plane curve; for example, the point  $x = y = 0$  is a double point on the cubic curve

$$x^3 + y^3 = 3xyz,$$

any line through it meeting the curve in only one further point. It is known that a plane curve of order N cannot possess more than

$$\frac{1}{2}(N - 1)(N - 2)$$

double points, a k-ple point (i.e. a point such that any line through it meets the curve only in N - k further points) counting as  $\frac{1}{2}k(k - 1)$  double

\* It is always to be understood that the curves and loci spoken of are *algebraical*.

† The word *solid* will always mean a three-dimensional space. We shall sometimes find it convenient to use the word *solid* as well as the symbols [3] and  $S_3$ .

points\*. If  $d$  is the actual number of double points possessed by the plane curve the number  $\frac{1}{2}(N-1)(N-2) - d$  was called by Cayley the *deficiency* of the curve. This number is in fact the same as the *genus* of the curve. The most fundamental property of the genus is that it is invariant for birational transformation of the curve; the genus of a curve in space of any number of dimensions can therefore be defined as the deficiency of a plane curve with which it is birationally equivalent.

The explanation of what is meant by *birational transformation* must be given here. Two curves are said to be *birationally equivalent* or to be in *birational correspondence* when the coordinates of a point on either curve are rational functions of the coordinates of a point on the other. In this way to a given point of either curve there will correspond one and only one point of the other; but multiple points will prove exceptions to this rule, to a multiple point on one of the curves there will correspond several points on the other. Thus we can say that there is a (1, 1) correspondence between the two curves, with certain reservations as to the multiple points. But it appears that we can always regard a multiple point as consisting of several points on different *branches* of a curve, and if we regard the multiple point in this way we can say that the correspondence is (1, 1) without exception. Thus a birational correspondence and a (1, 1) correspondence between two curves mean the same thing; and the fundamental property of the genus is that it is the same for two curves which are in (1, 1) correspondence.

If we are considering correspondences between the points of two curves, or between the points of a single curve, then a double point must be regarded as two distinct points on different branches of the curve. At a cusp, however, there is only a single branch.

In the quadratic transformation

$$x = \frac{1}{X}, \quad y = \frac{1}{Y}, \quad z = \frac{1}{Z},$$

the rational quartic

$$y^2z^2 + z^2x^2 + x^2y^2 = 0,$$

with nodes at the three vertices of the triangle of reference, is transformed into the conic

$$X^2 + Y^2 + Z^2 = 0,$$

and to each node of the quartic there correspond two distinct points of the conic. Corresponding to the node  $y = z = 0$  we have the two points in which the conic is met by the line  $X = 0$ ; and to either of these points on the conic corresponds the node  $y = z = 0$  on the quartic, the two points on the conic giving points on two distinct branches of the quartic.

\* It may be equivalent to more than this number of double points if the  $k$  tangents are not all distinct or are such that some of them meet the curve in more than  $k + 1$  (instead of exactly  $k + 1$ ) points at the multiple point.

On the other hand, the rational quartic

$$y^2z^2 + z^2x^2 + x^2y^2 = 2xyz(x + y + z)$$

has cusps at the three vertices of the triangle of reference, and is transformed by the same transformation into the conic

$$X^2 + Y^2 + Z^2 = 2(YZ + ZX + XY).$$

Then to each cusp of the quartic there corresponds only one point of the conic, e.g. to the cusp  $y = z = 0$  corresponds the point in which the conic is touched by its tangent  $X = 0$ .

6. When two curves are in (1, 1) correspondence it is of course not necessary that they should belong to spaces of the same number of dimensions; either of them can belong to a space of any number of dimensions. The genus therefore of a curve in  $[n]$  is simply the genus or deficiency of the projection of this curve from a space  $[n - 3]$  on to a plane; the correspondence between the curve and its projection will be (1, 1) if the  $[n - 3]$  is of general position. A curve has the same genus as any curve of which it is the projection.

For example, we may project the curve of intersection of two quadric surfaces in ordinary space on to a plane from a point  $O$ . If  $O$  is of general position in regard to the curve there are two and only two of its chords\* which pass through  $O$ ; the projection is a plane quartic with two double points and therefore of genus 1. Hence the curve of intersection of two quadrics is also of genus 1.

Of the  $\infty^3$  possible positions of  $O$  there are four (not on the curve) for which an infinity of chords of the curve pass through  $O$ , these being the vertices of the four quadric cones which belong to the pencil of quadrics containing the curve. The projection from one of these points does not give a (1, 1) correspondence but a (2, 1) correspondence, and the genus of the curve is altered by such a projection.

A curve of genus zero is said to be a *rational curve* because the coordinates of any point on it can be expressed as rational functions of a parameter, and this parameter can be so chosen as to be a rational function of the coordinates of a point of the curve†. Thus to each point of the curve corresponds one and only one value of the parameter and to each value of the parameter corresponds one and only one point of the curve. A rational curve is birationally equivalent to a straight line and all rational curves are birationally equivalent to one another.

A curve of genus 1 is said to be an *elliptic curve*; but it is not true that all elliptic curves are birationally equivalent to one another. There is belonging to an elliptic curve an invariant called its *modulus*; and in order

\* Salmon, *Geometry of Three Dimensions* (Dublin, 1914), vol. 1, pp. 355, 356.

† If we have expressed the coordinates of a point of a curve as rational functions of a parameter and this parameter is *not* a rational function of the coordinates, we can always find a second parameter which is; the second parameter is a rational function of the first and the coordinates are rational in terms of it. See Lüroth, *Math. Ann.* 9 (1875), 163.

that two elliptic curves should be birationally equivalent it is necessary and sufficient that they should have the same modulus.

A curve of genus 2 is said to be *hyperelliptic*, although not all hyperelliptic curves are of genus 2\*.

7. When we project a curve  $C$  of order  $N$  in  $[n]$  on to any lower space the order of the projected curve is also  $N$  provided that the centre of projection does not meet  $C$ . If the centre of projection is a space  $[r]$  the space on to which we are projecting is an  $[n - r - 1]$ ; an arbitrary  $[n - r - 2]$  in this space meets the projected curve in a number of points equal to its order, and this number is the same as the number of points in which the  $[n - 1]$  joining  $[n - r - 2]$  to  $[r]$  meets  $C$ . If  $C$  is met in  $M$  points by the centre of projection the projected curve is of order  $N - M$ . If we project on to a plane from an  $[n - 3]$  which does not meet  $C$  we know that we shall obtain a plane curve of order  $N$  with  $\frac{1}{2}(N - 1)(N - 2) - p$  double points, where  $p$  is the genus of  $C$ . But the space  $[n - 2]$  which joins  $[n - 3]$  to any one of these double points must, unless it contains a double point of  $C$  itself, contain two different points of  $C$ ; so that we have a chord of  $C$  meeting  $[n - 3]$ . Conversely, any chord of  $C$  which meets  $[n - 3]$  gives rise to a double point of the projected curve. Thus, if  $\delta$  is the number of actual double points of  $C$ , there must be  $\frac{1}{2}(N - 1)(N - 2) - p - \delta$  chords of  $C$  meeting an  $[n - 3]$  of general position; so that *the chords of  $C$  form a three-dimensional locus of order  $\frac{1}{2}(N - 1)(N - 2) - p - \delta$ .*

8. *Normal curves.* We now introduce the important concept of a *normal curve*†. A curve is said to be *normal* when it cannot be obtained by projection from a curve of the same order in space of higher dimension. It is clear that no curve can lie in a space of higher dimension than the order  $n$  of the curve, for taking any  $n + 1$  points of the curve we determine thereby a space of dimension  $n$  at most, which contains the curve since it meets it in a number of points greater than its order. For example: a curve of the second order always lies in a plane.

The coordinates of a point of a rational curve of order  $n$  in  $[m]$  can be expressed as rational functions of a parameter  $\theta$ . If the coordinates are homogeneous, and so  $m + 1$  in number, the coordinates of a point of the curve can be taken as polynomials in  $\theta$ . Further,  $\theta$  can be so chosen that it is a rational function of the coordinates (§ 6) so that to any given value of  $\theta$  there corresponds one and only one point of the curve. Then none of the  $m + 1$  polynomials can be of degree higher than  $n$ , for otherwise a prime  $S_{m-1}$ , which is given by a single linear equation in the

\* A curve of genus 2 is the simplest example of a class of curves which are said to be *hyperelliptic*. We can have hyperelliptic curves of any genus; but *all* curves of genus 2 are necessarily hyperelliptic. See e.g. Severi, *Trattato di Geometria Algebrica*, I, 1, 159 (Bologna, 1926).

† See Severi, *ibid.* 110–111.

coordinates, would meet the curve in more than  $n$  points; while one polynomial at least must actually be of degree  $n$ . Thus a rational curve of order  $n$  cannot lie in a space of dimension greater than  $n$ , since we cannot have more than  $n + 1$  linearly independent polynomials of order  $n$  in  $\theta$ .

On the other hand, a rational curve of order  $n$  can always be regarded as the projection of a rational curve of order  $n$  in  $[n]^*$ . If the curve is in  $[m]$  we can suppose the homogeneous coordinates  $x_0, x_1, \dots, x_m$  of any point on it to be linearly independent polynomials of order  $n$  in a parameter  $\theta$ . We can then choose  $n - m$  further polynomials of order  $n$  in  $\theta$  such that all the  $n + 1$  polynomials are linearly independent; we then take a curve in  $[n]$ , the homogeneous coordinates  $x_0, x_1, \dots, x_n$  of a point on it being proportional to these polynomials. The former curve can be regarded as lying in the  $[m]$  whose equations are  $x_{m+1} = x_{m+2} = \dots = x_n = 0$  and is the projection of the normal curve from the  $[n - m - 1]$  whose equations are  $x_0 = x_1 = \dots = x_m = 0$ .

We can, merely by means of a linear transformation of the coordinates, take the coordinates of a point on a rational normal curve of order  $n$  to be

$$x_0 = \theta^n, \quad x_1 = \theta^{n-1}, \quad \dots, \quad x_r = \theta^{n-r}, \quad \dots, \quad x_{n-1} = \theta, \quad x_n = 1.$$

The expressions  $(\theta^2, \theta, 1)$  for a point on a conic and  $(\theta^3, \theta^2, \theta, 1)$  for a point on a twisted cubic are well known.

The curve is given uniquely by the equations

$$\frac{x_0}{x_1} = \frac{x_1}{x_2} = \dots = \frac{x_r}{x_{r+1}} = \dots = \frac{x_{n-1}}{x_n},$$

or

$$\left\| \begin{array}{ccccccc} x_0 & x_1 & \dots & x_r & \dots & x_{n-1} & \\ x_1 & x_2 & \dots & x_{r+1} & \dots & x_n & \end{array} \right\| = 0.$$

Incidentally we have the equations of  $\frac{1}{2}n(n-1)$  quadric primals containing the curve; these are linearly independent and any other quadric primal containing the curve is in fact linearly dependent from these.

The chords of the curve form the three-dimensional locus given by

$$\left\| \begin{array}{ccccccc} x_0 & x_1 & \dots & x_r & \dots & x_{n-2} & \\ x_1 & x_2 & \dots & x_{r+1} & \dots & x_{n-1} & \\ x_2 & x_3 & \dots & x_{r+2} & \dots & x_n & \end{array} \right\| = 0,$$

which is of order  $\frac{1}{2}(n-1)(n-2)^\dagger$ .

\* Veronese, *Math. Ann.* 19 (1882), 208.

† The coordinates of a point on the three-dimensional locus of chords are of the form

$$(\theta^n + \lambda\phi^n, \quad \theta^{n-1} + \lambda\phi^{n-1}, \quad \dots, \quad \theta + \lambda\phi, \quad 1 + \lambda),$$

and depend on the three parameters  $\theta, \phi, \lambda$ .

For the order of the system of equations given by the vanishing of the determinants of a matrix see Salmon, *Higher Algebra* (Dublin, 1885), Lesson 19.



In particular the rational quartic curve is normal in [4] and can be given by

$$x_0 : x_1 : x_2 : x_3 : x_4 = \theta^4 : \theta^3 : \theta^2 : \theta : 1.$$

Its equations are

$$\frac{x_0}{x_1} = \frac{x_1}{x_2} = \frac{x_2}{x_3} = \frac{x_3}{x_4}$$

and it lies on six linearly independent quadric primals. It can be shewn that there is one quadric primal which contains not only the quartic curve but all its tangents; its equation is

$$x_0x_4 - 4x_1x_3 + 3x_2^2 = 0.$$

The chords of the curve form the cubic primal

$$\begin{vmatrix} x_0 & x_1 & x_2 \\ x_1 & x_2 & x_3 \\ x_2 & x_3 & x_4 \end{vmatrix} = 0.$$

We now state a fundamental result: *A curve of order  $n$  and genus  $p$  is normal in  $[n - p]$  if  $n > 2p - 2$ ; that is to say, every curve of genus  $p$  whose order  $n$  is greater than  $2p - 2$  can be obtained by projection from some curve of order  $n$  and genus  $p$  in  $[n - p]^*$ . In particular an elliptic curve of order  $n$  can always be obtained by projection from an elliptic curve of order  $n$  in  $[n - 1]$ .*

**9. Ruled surfaces.** A surface formed by a singly infinite system of straight lines is called a *ruled surface*; the lines are called the *generators* of the surface. If two different prime sections of the surface are taken it is clear that they are in (1, 1) correspondence, two points corresponding when they lie on the same generator; the sections are met each in one point by every generator. Hence all prime sections of the surface are of the same genus, so that we can speak of the genus of a ruled surface, meaning thereby the genus of its prime sections. We thus have rational ruled surfaces, elliptic ruled surfaces and so on.

Incidentally we can speak of the genus of any singly infinite set of elements, meaning thereby the genus of a curve whose points are in (1, 1) correspondence with the elements of the set.

**10.** Suppose now that we have a ruled surface of order  $n$  in [3]. It is clear that the tangent plane at any point contains the generator which passes through that point.

Consider the section of the surface by a plane passing through a generator  $g$ ; it consists of  $g$  and a curve  $C_{n-1}$  of order  $n - 1$ . We may

\* See Clifford, "On the classification of loci," *Phil. Trans.* 169 (1879), 663; and *Collected Papers* (London, 1882), 329. Clifford's result is obtained by the use of Abelian integrals and states that  $p \leq \frac{n}{2}$ . This same result is arrived at differently by Veronese, *Math. Ann.* 19 (1882), 213.

For the complete statement with  $n > 2p - 2$  see Segre, *Math. Ann.* 30 (1887), 207.

assume that, for a general position of this plane,  $C_{n-1}$  is irreducible and does not touch  $g$ . Then it will meet  $g$  in  $n - 1$  distinct points.  $C_{n-1}$  is simply the locus of the points in which the plane is met by the generators, and every generator other than  $g$  meets  $C_{n-1}$  in one point. Thus  $C_{n-1}$  has the same genus as the ruled surface; to each point of  $C_{n-1}$  there corresponds a generator of the surface passing through it and conversely, the point of  $C_{n-1}$  corresponding to  $g$  being one of their  $n - 1$  common points. Thus through the  $n - 2$  remaining intersections of  $g$  and  $C_{n-1}$  there will pass other generators, so that we conclude that *every generator is met by  $n - 2$  others*.

If now we take a point  $P$  on  $C_{n-1}$  the plane which contains the tangent of  $C_{n-1}$  at  $P$  and the generator which passes through  $P$  will be the tangent plane of the ruled surface at  $P$ . In particular, the tangent plane of the ruled surface at that intersection of  $C_{n-1}$  and  $g$  which is not an intersection of  $g$  with another generator is the plane of  $C_{n-1}$  itself.

If we consider any generator  $g$ , every point of  $g$  is the point of contact of a tangent plane passing through  $g$ , while every plane passing through  $g$  is a tangent plane touching the surface at some point of  $g$ . There is thus established a projectivity between the range of points on  $g$  and the pencil of planes through  $g$ ; the range of points of contact is related to the pencil of corresponding tangent planes. The particular case of this property of a quadric is familiar; the generators of either system are met by those of the other in related ranges of points.

11. We have seen that every generator of a ruled surface of order  $n$  in [3] is met by  $n - 2$  others. We thus have on the surface a *double curve* meeting every generator in  $n - 2$  points; this curve is the locus of intersections of pairs of generators and at any point of it there are two tangent planes to the surface, one containing each of the two generators which intersect there.

Similarly we have a *bitangent developable* formed by the planes containing pairs of intersecting generators; there are  $n - 2$  planes of this developable passing through each generator, and every plane of the developable touches the surface in two points—one on each of the generators lying in the plane.

The section of the ruled surface by any plane has double points at the intersections of the plane with the double curve. The tangent cone to the surface from any point has as double tangent planes those planes of the bitangent developable which pass through the point.

If the ruled surface is of order  $n$  and genus  $p$  a plane section is a curve of order  $n$  and genus  $p$ ; such a curve has  $\frac{1}{2}(n - 1)(n - 2) - p$  double points. Hence the plane must meet the double curve of the ruled surface in these  $\frac{1}{2}(n - 1)(n - 2) - p$  points, so that the order of the double curve is  $\frac{1}{2}(n - 1)(n - 2) - p$ .

Those planes through an arbitrary point which contain the generators of the surface form a singly infinite aggregate of genus  $p$ ; they meet an arbitrary plane in the tangents of a curve of class\*  $n$  and genus  $p$ . Such a curve has  $\frac{1}{2}(n-1)(n-2) - p$  double tangents; so that there are, passing through an arbitrary point,  $\frac{1}{2}(n-1)(n-2) - p$  planes each of which contains two generators of the surface. Hence there are  $\frac{1}{2}(n-1)(n-2) - p$  planes of the bitangent developable passing through an arbitrary point, so that the bitangent developable is of class  $\frac{1}{2}(n-1)(n-2) - p$ ; the class of a developable in [3] being defined as the number of its planes which pass through an arbitrary point.

We have thus shewn that, for a ruled surface of order  $n$  and genus  $p$ , the order of the double curve and the class of the bitangent developable are both equal to

$$\frac{1}{2}(n-1)(n-2) - p.$$

12. *The classification of ruled surfaces in three dimensions.* We shall classify ruled surfaces in three dimensions according to:

- (i) the order;
- (ii) the genus;
- (iii) the double curve;
- (iv) the bitangent developable.

13. *Correspondence formulae.* The position of a point on a straight line or on any rational curve is given by a single parameter, this being chosen as in § 6. Suppose then that we have a correspondence between the points of a rational curve; this means that there is an algebraic relation connecting the parameters  $\theta$  and  $\phi$  of corresponding points  $P$  and  $Q$ , the correspondence being given by equating some polynomial in  $\theta$  and  $\phi$  to zero. If the polynomial is of degree  $\alpha$  in  $\theta$  and of degree  $\beta$  in  $\phi$  we say that there is an  $(\alpha, \beta)$  correspondence between  $P$  and  $Q$ ; when  $P$  is given there are  $\beta$  corresponding positions of  $Q$ , and when  $Q$  is given there are  $\alpha$  corresponding positions of  $P$ . It is then evident that there are  $\alpha + \beta$  points which coincide with one of their corresponding points; their parameters are simply the roots of the equation of degree  $\alpha + \beta$  which is obtained by equating the polynomial to zero after putting  $\theta = \phi$ . We shall then say that on a rational curve there are  $\alpha + \beta$  coincidences in an  $(\alpha, \beta)$  correspondence. This is Chasles' principle of correspondence†.

There will be certain points  $P$  for which two of the  $\beta$  corresponding points  $Q$  coincide; the number of these is  $2\alpha(\beta - 1)$  since the condition that an equation of order  $\beta$  should have a double root is of order  $2\beta - 2$

\* The class of a plane curve is defined as the number of its tangents which pass through an arbitrary point.

† *Comptes Rendus*, 53 (1864), 1175.

in its coefficients\*. Here it is not necessary that the correspondence should be between the points of the same curve;  $\theta$  and  $\phi$  may be the parameters of points on two different rational curves. These points  $P$  for which two corresponding points  $Q$  coincide are called branch-points of the correspondence. Similarly, there are  $2\beta(\alpha - 1)$  branch-points  $Q$  for which two of the  $\alpha$  corresponding points  $P$  coincide.

We may have two correspondences between the points  $P$  and  $Q$  of a rational curve or between the points  $P$  of one rational curve and the points  $Q$  of another rational curve; suppose that we have an  $(\alpha, \beta)$  correspondence between  $P$  and  $Q$  and an  $(\alpha', \beta')$  correspondence between  $P'$  and  $Q'$ ;  $P'$  is on the same rational curve as  $P$  and  $Q'$  is on the same rational curve as  $Q$ . Then for any given position of  $P$  and  $P'$  the  $\beta$  positions of  $Q$  will, in general, be distinct from the  $\beta'$  positions of  $Q'$ ; there are, however,  $\alpha\beta' + \alpha'\beta$  positions such that if  $P$  and  $P'$  together take up one of them one of the  $\beta$  corresponding points  $Q$  coincides with one of the  $\beta'$  corresponding points  $Q'$ . The condition that two equations of orders  $\beta$  and  $\beta'$  should have a common root is of order  $\beta'$  in the coefficients of the first equation and of order  $\beta$  in the coefficients of the second equation†.

14. These results which we have obtained for rational curves can be generalised for curves of any genus  $p$ ; in order to do this we must introduce the idea of the *valency*‡ of a correspondence and the idea of a *linear series of sets of points* on a curve.

If we have a curve in  $[n]$  then the family of primals§

$$\lambda_0 f_0 + \lambda_1 f_1 + \dots + \lambda_r f_r = 0$$

cuts out on the curve a *linear series of sets of points*. The primals  $f = 0$  are all of the same order and the equation of any primal of the system depends linearly and homogeneously upon the  $r + 1$  parameters  $\lambda$ ; we have a *linear system* of primals. Thus on a plane curve a linear series of sets of points is cut out by a linear system of curves; on a twisted curve a linear series of sets of points is cut out by a linear system of surfaces, and so on. We will then consider for definiteness a linear series of sets of points on a plane curve.

The linear system of plane curves

$$\lambda_0 f_0 + \lambda_1 f_1 + \dots + \lambda_r f_r = 0$$

contains  $\infty^r$  curves and cuts out on a curve  $h = 0$  a linear series of sets of points. There may be an infinity of curves of the linear system passing

\* Salmon, *Higher Algebra* (Dublin, 1885), 99.

† Salmon, *ibid.* 69.

‡ Called *Werthigkeit* by Brill, *Math. Ann.* 7 (1874), 611.

§  $f$  is a homogeneous polynomial in the  $n + 1$  homogeneous coordinates

$$x_0, x_1, \dots, x_n.$$

It is assumed that these  $r + 1$  polynomials  $f$  are linearly independent.

through a set of the linear series; it can be shewn that this happens when and only when there are curves of the system containing the curve  $h = 0$ . We can then always work with a reduced linear system of curves\*, none of which contains the curve  $h = 0$  as a part. Then, supposing that we are working with such a system containing the  $r + 1$  parameters  $\lambda$ , the linear series of sets of points contains  $\infty^r$  sets, each set containing the same number  $n$  of points. Through  $r$  general points of the curve  $h = 0$  there will pass just one curve of the linear system, as we have just enough conditions to determine the ratios of the parameters  $\lambda$ . The linear series of sets of points is then said to be of *freedom*  $r$  and is denoted by  $g_n^r$ ;  $r$  general points of the curve  $h = 0$  determine just one set of  $g_n^r$ . The number  $n$  is called the *grade* of the linear series.

If now we have an  $(\alpha, \beta)$  correspondence between the points  $P$  and  $Q$  of a curve of genus  $p$ , and if a point  $P$  of the curve counted  $\gamma$  times, taken together with the  $\beta$  points  $Q$  which correspond to it, gives a set of points which varies in a linear series of sets of  $\beta + \gamma$  points as  $P$  varies on the curve, the correspondence is said to be of *valency*†  $\gamma$ . It is not necessary that every set of the linear series should be given by the variation of  $P$ , but merely that all the  $\infty^1$  sets of points so obtained should belong to some linear series of sets of  $\beta + \gamma$  points. When this is so it can be shewn that a point  $Q$  of the curve counted  $\gamma$  times, taken together, with the  $\alpha$  points  $P$  which correspond to it, gives a set of points which vary in a linear series of sets of  $\alpha + \gamma$  points as  $Q$  varies on the curve. On a general curve every correspondence has a valency; correspondences without a valency can only exist on curves which are special for their genus‡.

We can, for example, set up a correspondence between the points  $P$  and  $Q$  of a plane cubic without a double point; saying that the points  $P$  and  $Q$  correspond when the tangent at  $P$  passes through  $Q$ . To any point  $P$  corresponds one point  $Q$ , since the tangent at  $P$  only meets the cubic in one other point; to any point  $Q$  correspond four points  $P$  since four tangents can be drawn to the cubic from any point of itself; thus the correspondence is a  $(4, 1)$  correspondence. Also its valency is 2; the point  $P$  counted twice together with the point  $Q$  which corresponds to it form the complete intersection of the cubic with its tangent, and therefore vary, as  $P$  varies on the curve, in a linear series of sets of three points, viz. that cut out by the lines of the plane. The four points of contact of the tangents drawn to the curve from any point  $Q$  of itself lie on the first polar of  $Q$ , which is a conic touching the curve at  $Q$ ; hence the point  $Q$  counted twice together with the four points  $P$  which correspond to it form the complete intersection of the cubic with a conic and therefore vary in a linear series of

\* See Severi, *Trattato di Geometria Algebrica*, I, 1, 20 (Bologna, 1926).

† Severi, *ibid.* 198.

‡ Hurwitz, *Math. Ann.* 28 (1887), 565.

sets of six points as  $Q$  varies on the curve, viz. that cut out by the conics of the plane.

The analogues of the formulae for rational curves are as follows.

If we have on a curve of genus  $p$  an  $(\alpha, \beta)$  correspondence of valency  $\gamma$  there are  $\alpha + \beta + 2\gamma p$  coincidences\*. For example, there are nine coincidences in the  $(4, 1)$  correspondence just mentioned on the cubic curve of genus 1; these are simply the nine inflections of the curve. Again, the number of branch-points  $P$  for which two of the  $\beta$  corresponding points  $Q$  coincide† is  $2\alpha(\beta - 1) + 2(\alpha - \gamma^2)p$  and the number of branch-points  $Q$  for which two of the  $\alpha$  corresponding points  $P$  coincide‡ is  $2\beta(\alpha - 1) + 2(\beta - \gamma^2)p$ . Finally, if we have on the same curve of genus  $p$  an  $(\alpha, \beta)$  correspondence of valency  $\gamma$  and an  $(\alpha', \beta')$  correspondence of valency  $\gamma'$  the number of pairs of points common to the two correspondences‡ is  $\alpha\beta' + \alpha'\beta - 2\gamma\gamma'p$ .

As an example of this last result consider a plane quartic curve without double points ( $p = 3$ ). If  $P$  and  $Q$  correspond when the tangent at  $Q$  passes through  $P$  we have a  $(2, 10)$  correspondence of valency 2; if inversely  $P'$  and  $Q'$  correspond when the tangent at  $P'$  passes through  $Q'$  we have a  $(10, 2)$  correspondence of valency 2. Then the number of pairs of points common to these two correspondences is

$$2 \cdot 2 + 10 \cdot 10 - 2 \cdot 2 \cdot 2 \cdot 3 = 80 = 2 \times 28 + 24.$$

The interpretation of this result is clear since the curve has 28 bitangents and 24 inflections. If one of the points of contact of a bitangent is taken as a position of  $P$  and  $P'$  then the other point of contact is one of the corresponding positions of  $Q$  and also one of the corresponding positions of  $Q'$ . Since there are two points of contact of each bitangent we have in this way 56 pairs of points common to the two correspondences. Again, if the point of contact of an inflectional tangent is taken as a position of  $P$  and  $P'$  this same point is one of the corresponding positions of  $Q$  and also one of the corresponding positions of  $Q'$ .

We must also mention the fact that the valency of a correspondence can be negative; this is clear from other definitions of the valency but it also arises naturally from the idea of the equivalence of sets of points on a curve, two sets of points on a curve being *equivalent* when they belong to the same linear series.

As an example of a correspondence with negative valency let us consider the correspondence between the point  $P$  of a plane cubic without a double point and

\* See Cayley, *Comptes Rendus*, 62 (1866), 586 = *Papers*, 5, 542; *Phil. Trans.* 158 (1868), 146 = *Papers*, 6, 265. Brill, *Math. Ann.* 6 (1873), 33; 7 (1874), 607. Severi, *Memorie Torino* (2), 54 (1904), 11; *Trattato di Geometria Algebrica*, I, 1, 233 (Bologna, 1926).

† This formula has been given by Professor Baker in lectures. It can be deduced from Cayley's formula (*Papers*, 6, 267).  $P$  and  $Q$  are on the same curve.

‡ Brill, *Math. Ann.* 6 (1873), 42; 7 (1874), 611. Hurwitz, *ibid.* 28 (1887), 568. This result we shall in future call *Brill's formula*.

the point  $R$  which is the second tangential of  $P$ . If the tangent at  $P$  meets the curve again in  $Q$ ,  $Q$  is called the *tangential* of  $P$ ; if then the tangent at  $Q$  meets the curve again in  $R$ ,  $R$  is called the *second tangential* of  $P$ . Clearly to a given point  $P$  there corresponds one and only one point  $R$ .

Now it is known that if conics are drawn through four fixed points of a cubic curve the line joining their two remaining intersections with the curve meets the cubic again in a fixed point\*. In particular, the conics which have four-point contact with the curve at  $P$  meet it again in two points whose join passes through  $R$ .

Take then any point  $O$  on the cubic and join  $OR$ , meeting the cubic again in  $R'$ . The conic through  $O$  which has four-point contact with the curve at  $P$  passes through  $R'$ . Moreover, if  $P'$  is any point whose second tangential is  $R'$  we can draw a conic having four-point contact with the curve at  $P'$  and passing through  $O$  and  $R$ .

Now the conics passing through  $O$  cut out on the cubic a linear series of sets of five points; one set of this series consists of  $R'$  taken with  $P$  counted four times, while another consists of  $R$  taken with  $P'$  counted four times. Thus these two sets of points are equivalent, so that we write

$$4P + R' \equiv 4P' + R.$$

Writing this in the form

$$R - 4P \equiv R' - 4P',$$

we say that the correspondence from  $P$  to  $R$  has the valency  $-4$ .

This exemplifies a general theorem which states that the "product" of two correspondences of valencies  $\gamma_1$  and  $\gamma_2$  has the valency  $-\gamma_1\gamma_2$ . The correspondence from  $P$  to its second tangential is the "square" of the correspondence from  $P$  to its first tangential, and this latter correspondence we have already seen to have valency 2.

15. A correspondence has, of course, two senses; in the "forward" sense the  $\beta$  points  $Q_1, Q_2, \dots, Q_\beta$  correspond to the point  $P$ , while in the "backward" sense the  $\alpha$  points  $P_1, P_2, \dots, P_\alpha$  correspond to the point  $Q$ . In calculating the number of pairs of points common to two correspondences by Brill's formula we imply that we take both correspondences in the same sense: having chosen a point we proceed to the two sets which correspond to it, either both in the forward sense or both in the backward sense; the formula gives the number of positions of the chosen point for which a point of one of the two sets coincides with a point of the other.

There is a special kind of correspondence which we shall call a *symmetrical correspondence*; in such a correspondence there is only one sense, or there is only one way of passing from a given point to those points which correspond to it. The two indices of such a correspondence are equal; and if  $Q$  is one of the set of points corresponding to  $P$  then  $P$  is conversely one of the set of points corresponding to  $Q$ . As an example of such a correspondence let us consider, on a quadric  $S$  in ordinary space, the curve  $C$  which is the intersection of  $S$  with another quadric; and suppose that two points of  $C$  correspond when they lie on the same generator of  $S$ .

\* Salmon, *Higher Plane Curves* (Dublin, 1879), 134.

Every generator of  $S$  meets  $C$  in two points; so that to any point  $P$  of  $C$  there correspond the two other points  $Q_1, Q_2$  in which  $C$  is met by the generators of  $S$  through  $P$ . To  $Q_1$  correspond  $P$  and another point  $P_1$ , while to  $Q_2$  correspond  $P$  and another point  $P_2$ . We thus have a symmetrical (2, 2) correspondence.

The formulae giving coincidences and branch-points hold for symmetrical correspondences as for other correspondences; but it is important to remark that *the number of pairs of points common to two correspondences as given by Brill's formula must be halved when both the correspondences are symmetrical\**.

Still more special correspondences are those known as *involutions*; an involution is a symmetrical correspondence in which a point  $P$  and the  $m$  points  $Q_1, Q_2, \dots, Q_m$  corresponding to it form a closed set; to any one of them correspond the remaining  $m$ . A simple example of an involution is the correspondence between the points of a plane curve of order  $n$  which are collinear with a given point  $O$ . If  $P$  is any point of the curve there are  $n - 1$  points corresponding to  $P$ , namely, the remaining intersections of the curve with the line  $OP$ . We thus have a set of  $n$  points such that to any one point of the set there correspond the remaining  $n - 1$ ; or we have a symmetrical  $(n - 1, n - 1)$  correspondence which is an involution.

An involution, as consisting of  $\infty^1$  sets of points, will have a genus just as will a curve consisting of  $\infty^1$  points. In the example given the involution is rational, its sets being in (1, 1) correspondence with the lines of a plane pencil. We can, however, have involutions of any genus $\dagger$ .

**16. Zeuthen's formula.** There is a formula due to Zeuthen concerning a correspondence between the points of two curves which is of great importance. If we have an  $(\alpha, \alpha')$  correspondence between the points of two curves  $C$  and  $C'$  whose respective genera are  $p$  and  $p'$ , so that to any given point of  $C$  there correspond  $\alpha'$  points of  $C'$  and to any given point of  $C'$  there correspond  $\alpha$  points of  $C$ , and if the correspondence has  $\eta$  branch-points on  $C$  and  $\eta'$  branch-points on  $C'$ , then we have the relation

$$\eta - \eta' = 2\alpha(p' - 1) - 2\alpha'(p - 1).$$

We shall always refer to this as *Zeuthen's formula* $\ddagger$ . A geometrical interpretation of the formula has been given by Severi $\S$ .

If  $\alpha$  and  $\alpha'$  are both unity then  $\eta$  and  $\eta'$  are both zero; and so two curves in (1, 1) correspondence have the same genus.

\* In this case Brill's formula includes each pair twice (from each end it might be said).

$\dagger$  Severi, *Trattato di Geometria Algebrica*, I, 1, 52 (Bologna, 1926).

$\ddagger$  Zeuthen, *Math. Ann.* 3 (1871), 150.

$\S$  *Rendiconti del Reale Istituto Lombardo* (2), 36 (1903), 495.



The formula is at once verified when the curves are both rational; it becomes then  $\eta + 2\alpha = \eta' + 2\alpha'$ . But we have already seen\* that in such a correspondence  $\eta = 2\alpha (\alpha' - 1)$  and  $\eta' = 2\alpha' (\alpha - 1)$ , so that

$$\eta + 2\alpha = \eta' + 2\alpha' = 2\alpha\alpha'.$$

17. *The genus of a simple curve on a ruled surface.* One very important application of Zeuthen's formula is the calculation of the genus of a curve on a ruled surface. We suppose that the curve is a simple and not a multiple curve on the surface; so that through a general point of the curve there passes only one generator of the surface.

Suppose that we have in a space  $[r]$  a ruled surface of order  $n$  and genus  $p$ ; and suppose that there is on this ruled surface a curve of order  $\nu$  meeting each generator in  $k$  points. Suppose that this curve is of genus  $\pi$  and is touched by  $\eta$  generators. Then Zeuthen's formula gives

$$\eta = 2(\pi - 1) - 2k(p - 1),$$

there being a  $(1, k)$  correspondence between the points of a prime section of the ruled surface and the points of the curve, two points corresponding when they lie on the same generator.

If we take an arbitrary  $[r - 2]$  there is a pencil of primes passing through it; we consider the correspondence between primes of this pencil, two primes corresponding when they join  $[r - 2]$  to two points of the curve lying on the same generator. Then, since any one of the primes meets the curve in  $\nu$  points, there are  $\nu(k - 1)$  primes of the pencil which correspond to it; we have a symmetrical correspondence between the primes of the pencil in which both indices are  $\nu(k - 1)$ . Since Chasles' principle of correspondence can be applied to the primes of this pencil just as it can to the points of a line there will be  $2\nu(k - 1)$  coincidences of pairs of corresponding primes.

Now these coincidences can occur in two ways; either by the prime passing through one of the  $\eta$  points where the curve touches some generator or by the prime containing one of the generators which pass through the  $n$  intersections of  $[r - 2]$  with the surface. On each of these generators there are  $k$  points of the curve, and we count the prime joining  $[r - 2]$  to such a generator  $k(k - 1)$  times among the coincidences. Hence

$$2\nu(k - 1) = \eta + nk(k - 1),$$

so that, eliminating  $\eta$ ,

$$2\nu(k - 1) = 2(\pi - 1) - 2k(p - 1) + nk(k - 1),$$

whence †  $\pi = (\nu - 1)(k - 1) + pk - \frac{1}{2}nk(k - 1)$ ,

giving the genus  $\pi$  of the curve in terms of the other constants.

\* § 13.

† Segre, *Rom. Acc. Lincei Rendiconti* (4), 3<sup>2</sup> (1887), 3.

We have not mentioned the fact that the curve may have double points. If it has, the formula

$$\eta = 2(\pi - 1) - 2k(p - 1)$$

is not altered; but the prime joining  $[r - 2]$  to a double point  $P$  will count twice among the coincidences provided that  $P$  is not also a double point of the ruled surface. The double point must be regarded as two points  $P_1$  and  $P_2$  on different branches of the curve; these are on the same generator, the prime joining  $[r - 2]$  to  $P_1$  has the prime joining  $[r - 2]$  to  $P_2$  among its corresponding primes and the two coincide; and so with  $P_1$  and  $P_2$  interchanged. Thus

$$2\nu(k - 1) = \eta + nk(k - 1) + 2d,$$

or 
$$\pi = (\nu - 1)(k - 1) + pk - \frac{1}{2}nk(k - 1) - d,$$

where  $d$  is strictly the number of those points\* of the curve which count twice among the  $k$  intersections with a generator\*.

18. *The genus of a curve on a quadric.* Suppose, for example, that we have a curve on a quadric surface in [3]; it is clear that any two generators of the same system of this quadric are met by the curve in the same number of points, because planes can be drawn through them and any generator of the opposite system. Suppose then that the curve meets all generators of one system in  $\alpha$  points and is of order  $\alpha + \beta$ , meeting all generators of the other system in  $\beta$  points. Then its genus will be

$$\begin{aligned} \pi &= (\alpha + \beta - 1)(\alpha - 1) - \alpha(\alpha - 1) - d \\ &= (\alpha - 1)(\beta - 1) - d, \end{aligned}$$

where  $d$  is the number of double points of the curve. If the curve has no double points then  $\pi = (\alpha - 1)(\beta - 1)$ .

This result for the genus of a curve on a quadric surface is important and will be of use subsequently. It is obtained at once by projecting the curve from a point on the quadric into a plane curve; if the point of projection is not on the curve we obtain a plane curve of order  $\alpha + \beta$  with  $d$  double points, a point of multiplicity  $\alpha$  and a point of multiplicity  $\beta$ , and therefore of genus

$$\frac{1}{2}(\alpha + \beta - 1)(\alpha + \beta - 2) - d - \frac{1}{2}\alpha(\alpha - 1) - \frac{1}{2}\beta(\beta - 1) = (\alpha - 1)(\beta - 1) - d.$$

If the point of projection is on the curve we obtain a plane curve of order  $\alpha + \beta - 1$  with  $d$  double points, a point of multiplicity  $\alpha - 1$  and a point of multiplicity  $\beta - 1$  and therefore of genus

$$\begin{aligned} \frac{1}{2}(\alpha + \beta - 2)(\alpha + \beta - 3) - d - \frac{1}{2}(\alpha - 1)(\alpha - 2) - \frac{1}{2}(\beta - 1)(\beta - 2) \\ = (\alpha - 1)(\beta - 1) - d. \end{aligned}$$

Zeuthen's formula can also be used to find the genus of a curve on a quadric cone. More generally, if we have a cone of order  $n$  and genus  $p$

\* Segre, *Math. Ann.* 34 (1889), 3.

and on it a curve of order  $\nu$  meeting every generator in  $k$  points other than the vertex, the genus of the curve is given by

$$\pi = (\nu - 1)(k - 1) + pk - \frac{1}{2}nk(k - 1)$$

with possibly a reduction for double points\*.

19. *The ruled surface generated by a correspondence between two curves.* If in a space  $[n]$  we have a correspondence between the points of two curves the lines joining pairs of corresponding points are  $\infty^1$  in aggregate and form a ruled surface.

The order of this ruled surface can be calculated when the properties of the correspondence and of the two curves themselves are known; suppose that the correspondence is an  $(\alpha, \alpha')$  correspondence between two curves of orders  $m$  and  $m'$ . Let us then take a space  $[n - 2]$  of general position; the order of the ruled surface is the number of its intersections with  $[n - 2]$ , i.e. the number of its generators which meet  $[n - 2]$ , or the number of pairs of corresponding points of the two curves whose joins meet  $[n - 2]$ .

We can set up a correspondence among the primes passing through  $[n - 2]$ , two primes corresponding when they join  $[n - 2]$  to corresponding points of the two curves. It is clear that the indices of this correspondence are  $m'\alpha$  and  $m\alpha'$ , so that there are  $m'\alpha + m\alpha'$  coincidences of pairs of corresponding primes. There is clearly a coincidence when the join of two corresponding points meets  $[n - 2]$ ; and, in general, these will be the only coincidences. Hence, in general, the order of the ruled surface is  $m'\alpha + m\alpha'$ . Through every point of the curve of order  $m$  there pass  $\alpha'$  generators of the surface, or we shall say that this curve is a multiple curve on the ruled surface of multiplicity  $\alpha'$ . Similarly the curve of order  $m'$  is a multiple curve on the ruled surface of multiplicity  $\alpha$ .

In particular, the order of the ruled surface generated by joining pairs of corresponding points in a  $(1, 1)$  correspondence between two curves is, in general, the sum of the orders of the two curves.

The result which we have obtained for the order of the ruled surface is always true except when the  $m'\alpha + m\alpha'$  coincidences include primes which do not contain generators of the ruled surface; this can happen only when the curves have one or more intersections which are "united points" of the correspondence. If  $P$  is an intersection of the two curves which is a united point then the  $\alpha'$  points of the curve of order  $m'$ , which correspond to  $P$  regarded as a point of the curve of order  $m$ , include  $P$ ; and the  $\alpha$  points of the curve of order  $m$ , which correspond to  $P$  regarded as a point of the curve of order  $m'$ , also include  $P$ . Thus the prime joining  $[n - 2]$

\* The result for a curve on a cone was first obtained by Sturm by application of a coincidence formula due to Schubert, *Math. Ann.* 19 (1882), 487.

to  $P$  counts among the  $m'\alpha + m\alpha'$  coincidences; the generator of the surface joining the pair of points which are united is indeterminate.

It is then clear that the order of the ruled surface is always  $m'\alpha + m\alpha'$  when the curves do not intersect; and it also has this value when the curves do intersect unless there are united points. But if there are  $i$  united points the order of the ruled surface\* is  $m'\alpha + m\alpha' - i$ . Here again a still further adjustment may be necessary, as the prime joining  $[n - 2]$  to a united point may count more than once among the  $m'\alpha + m\alpha'$  coincidences; it may be that when  $P$  is regarded as a point of the curve of order  $m$  there are more than one of the  $\alpha'$  corresponding points coinciding with it, and inversely. This cannot happen however when we are dealing with a  $(1, 1)$  correspondence; and we can always say that *the order of the ruled surface determined by a  $(1, 1)$  correspondence with  $i$  united points between two curves of orders  $m$  and  $m'$  is  $m + m' - i$ .*

20. In order to exemplify this last result let us consider in [3] a conic and a line which intersect in a point  $X$ .

Suppose that there is a  $(1, 1)$  correspondence between them and suppose first that  $X$  is not a united point. Then to  $X$  regarded as a point of the line there will correspond a point  $Z$  of the conic and to  $X$  regarded as a point of the conic there will correspond a point  $T'$  of the line. The line itself will then be a generator of the resulting ruled surface as joining the points  $X$  and  $T'$ , and the line  $ZX$  will also be a generator. If the tangents to the conic at  $Z$  and  $X$  meet in  $Y$ , we can take  $XYZT'$  as the tetrahedron of reference; the equations to the conic are

$$xz - y^2 = t = 0,$$

and to the line

$$y = z = 0.$$

Then any point of the conic has coordinates  $(\theta^2, \theta, 1, 0)$ , while any point of the line has coordinates  $(\phi, 0, 0, 1)$ , and the  $(1, 1)$  correspondence will be determined by a bilinear relation

$$a\theta\phi + b\theta + c\phi + d = 0.$$

But we already know that to  $\phi = \infty$  must correspond  $\theta = 0$  and that to  $\theta = \infty$  must correspond  $\phi = 0$ ; hence  $b = c = 0$  and  $\theta\phi$  is constant. We can therefore take

$$\theta\phi = 1$$

\* There are always united points if we consider the ruled surface generated by a correspondence between the points of the same curve. If the curve is of genus  $p$  and the correspondence of valency  $\gamma$  it has  $\alpha + \alpha' + 2\gamma p$  united points, and the order of the ruled surface will be  $m(\alpha + \alpha') - (\alpha + \alpha' + 2\gamma p)$ , where  $m$  is the order of the curve. But if the correspondence is a symmetrical correspondence of index  $\alpha$  this number will have to be halved; the order of the ruled surface is then  $\alpha(m - 1) - \gamma p$ . Thus, in particular, the order of the ruled surface formed by the joins of pairs of an involution of pairs of points on a rational curve is  $m - 1$ , on an elliptic curve  $m - 2$ . A curve which is neither rational nor elliptic and which possesses an ordinary (rational) involution of pairs of points is always hyperelliptic; for such a curve the order of the ruled surface is  $m - p - 1$ .

as giving the correspondence, and a generator of the ruled surface joins the two points

$$(\theta^2, \theta, 1, 0) \text{ and } (1, 0, 0, \theta).$$

Hence the coordinates of any point of the ruled surface are of the form

$$(\theta^2 + \lambda, \theta, 1, \lambda\theta).$$

If  $\theta$  is constant we have the points of a generator of the surface, and if  $\lambda$  is constant we have the points of a conic on the surface lying in a plane through  $XZ$ .

The coordinates satisfy the homogeneous cubic relation

$$xyz = y^3 + tz^2,$$

so that we have a ruled surface of the third order.

But suppose now that the (1, 1) correspondence between the line and conic is such that  $X$  is a united point. Then to a definite point  $T$  of the line will correspond a definite point  $Z$  of the conic, and our coordinates are as before with the bilinear relation

$$a\theta\phi + b\theta + c\phi + d = 0.$$

Now to  $\theta = \infty$  corresponds  $\phi = \infty$  so that  $a = 0$ , and to  $\theta = 0$  corresponds  $\phi = 0$  so that  $d = 0$ ; hence  $\theta$  is a constant multiple of  $\phi$  and we can take without loss of generality

$$\theta = \phi.$$

Then a generator of the ruled surface joins the two points

$$(\theta^2, \theta, 1, 0) \text{ and } (\theta, 0, 0, 1),$$

and the coordinates of any point on the ruled surface are of the form

$$(\theta^2 + \lambda\theta, \theta, 1, \lambda).$$

These satisfy the homogeneous quadratic relation

$$xz = y^2 + ty,$$

so that when there is a united point the ruled surface is only of the second order.

**21.** Just as, given two curves of orders  $m$  and  $m'$  in  $(\alpha, \alpha')$  correspondence in [3], there are (subject to a deduction for united points)  $m'\alpha + m\alpha'$  joins of pairs of corresponding points which meet an arbitrary line; so, dually, given two developables of classes  $m$  and  $m'$  in  $(\alpha, \alpha')$  correspondence in [3], there are (subject to a deduction for united planes)  $m'\alpha + m\alpha'$  intersections of pairs of corresponding planes which meet an arbitrary line. Or we can say that the two developables give a ruled surface of order  $m'\alpha + m\alpha'$ .

In space of three dimensions we have three fundamental constructs; *curves* formed by singly infinite families of points, *ruled surfaces* formed by singly infinite families of lines, and *developables* formed by singly infinite families of planes. We have seen how a ruled surface can be generated either by joining pairs of corresponding points on two curves or by taking the intersections of pairs of corresponding planes of two developables, and how the order of such a ruled surface is given in terms of the orders of the two curves or the classes of the two developables and the constants of the correspondence.

22. It is also clear that if a correspondence is set up between the points of a curve and the generators of a ruled surface the planes so determined, as joining the points of the curve to those generators of the ruled surface which correspond to them, form a developable; and, dually, if a correspondence is set up between the planes of a developable and the generators of a ruled surface, the points of intersection of corresponding planes and lines form a curve.

Let us consider then a curve of order  $m$  and a ruled surface of order  $\mu$ ; suppose that to any point of the curve there correspond  $\alpha$  generators of the ruled surface and that to any generator of the ruled surface there correspond  $a$  points of the curve. Let us calculate the class of the developable so formed; i.e. the number of its planes passing through any given point  $O$ .

We set up a correspondence between the points  $P$  and  $Q$  of the curve; the points  $P$  and  $Q$  corresponding when the line  $OP$  meets a generator of the ruled surface which corresponds to  $Q$ . Then given a point  $P$ , the line  $OP$  meets the ruled surface in  $\mu$  points, and each of the generators through these points gives  $a$  points  $Q$ ; so that when  $P$  is given there are  $\mu a$  positions of  $Q$ . Given a point  $Q$ , we have  $\alpha$  generators of the ruled surface, each of which is joined to  $O$  by a plane meeting the curve in  $m$  points; so that when  $Q$  is given there are  $m\alpha$  positions of  $P$ . The correspondence from  $P$  to  $Q$  is therefore an  $(m\alpha, \mu a)$  correspondence. The correspondence is of valency zero; the points  $P$  corresponding to a given point  $Q$  form the complete intersection of the curve with a set of  $\alpha$  planes through  $O$ . Hence there are  $m\alpha + \mu a$  coincidences of points  $P$  and  $Q$ ; and such a coincidence means that the plane joining a point of the curve to a corresponding generator of the ruled surface passes through  $O$ . This will be a plane of the developable unless the point of the curve happens to lie on the corresponding generator of the ruled surface; when this happens the plane of the developable is indeterminate.

Hence, subject to a deduction for united elements of the correspondence, the class of the developable is  $m\alpha + \mu a$ .

In particular, if we have a  $(1, 1)$  correspondence between the points of a curve of order  $m$  and the generators of a ruled surface of order  $\mu$ , and if there are  $i$  points of the curve which lie on the corresponding generators of the ruled surface, the planes joining corresponding elements form a developable of class  $m + \mu - i$ .

If we have a  $(1, 1)$  correspondence between the points of a line and the lines of a regulus\* the planes joining corresponding elements give a develop-

\* The word *regulus* is used to denote either system of generators of a quadric surface. It is thus a ruled surface of the second order; the other system of generators of the quadric are not, strictly speaking, generators, but *directrices* of this surface.

able of the third class\*, supposing that neither of the points in which the line meets the regulus lies on the line which corresponds to it; if however one of the two lines of the regulus which meet the line does so in the point corresponding to it the developable is only of the second class; while if both lines of the regulus which meet the line do so in the points corresponding to them we have a developable of the first class. In the first case we have the osculating planes of a twisted cubic, in the second the tangent planes of a quadric cone and in the third a pencil of planes through a line.

Analytically, let the line meet the regulus in the points  $X$  and  $Z$ ; through  $X$  there passes a generator of the quadric surface on which the regulus lies meeting the line of the regulus through  $Z$  in  $T$ ; similarly,  $Y$  is the point in which the line of the regulus through  $X$  is met by the generator of the complementary regulus through  $Z$ .

Then any line of the regulus is given as the intersection of two planes

$$y = \theta z, \quad x = \theta t,$$

the line  $XY$  being given by  $\theta = \infty$  and the line  $ZT$  by  $\theta = 0$ .

Any point of the line  $XZ$  is given by  $(\phi, 0, 1, 0)$ , the point  $X$  being given by  $\phi = \infty$  and the point  $Z$  by  $\phi = 0$ .

We then set up a correspondence between  $\theta$  and  $\phi$  and join lines of the regulus to corresponding points of  $XZ$  by planes.

If  $\phi = \frac{\theta + \beta}{\theta + \gamma}$  the planes are given by

$$(\theta + \beta)(y - \theta z) + \theta(\theta + \gamma)(x - \theta t) = 0,$$

and involve the parameter  $\theta$  in the third degree.

If  $\phi = \theta + \alpha$  then  $\theta = \infty$  gives the point  $X$  of the line corresponding to the line  $XY$  of the regulus. The planes are given by

$$(\theta + \alpha)(y - \theta z) + \theta(x - \theta t) = 0,$$

and involve the parameter  $\theta$  in the second degree.

If  $\phi = \theta$  then  $\theta = \infty$  gives the point  $X$  of the line corresponding to the line  $XY$  of the regulus, while  $\theta = 0$  gives the point  $Z$  of the line corresponding to the line  $ZT$  of the regulus. The planes are given by

$$y - \theta z + x - \theta t = 0,$$

and involve the parameter  $\theta$  in the first degree.

**23.** Dually, if we have a  $(1, 1)$  correspondence between the planes of a developable of class  $m$  and the generators of a ruled surface of order  $\mu$ , and if there are  $i$  planes of the developable which contain the corresponding generators of the ruled surface, the points of intersection of corresponding elements form a curve of order  $m + \mu - i$ .

A pencil of planes and the generators of a quadric cone when projectively related give a twisted cubic, provided that no plane of the pencil contains

\* von Staudt, *Beiträge zur Geometrie der Lage*, 3 (Nürnberg, 1860), 303; Reye, *Geometrie der Lage*, 2 (Stuttgart, 1907), 168.

the corresponding generator of the cone. A regulus and a pencil of planes when projectively related give a twisted cubic, provided that no line of the regulus lies in the plane which corresponds to it.

24. We may observe still further that if we have three curves in correspondence the planes joining corresponding triads of points, one point of such a triad being on each curve, will form a developable; while if we have three developables in correspondence the points of intersection of triads of corresponding planes, one plane of such a triad belonging to each developable, will form a curve. The generation of the twisted cubic by three related pencils of planes is familiar.

## SECTION II

### THE REPRESENTATION OF A RULED SURFACE IN THREE DIMENSIONS AS A CURVE ON A QUADRIC $\Omega$ IN FIVE DIMENSIONS

25. The lines of three-dimensional space are  $\infty^4$  in aggregate; a line can be determined by four parameters or by the ratios of five parameters. The most convenient representation of the lines of three-dimensional space is, however, by means of the ratios of six parameters which are connected by one relation, thus being reduced effectively to five\*.

Using homogeneous coordinates, let  $(x_1, y_1, z_1, t_1)$  and  $(x_2, y_2, z_2, t_2)$  be two points; and write

$$\begin{aligned} l &= t_1 x_2 - t_2 x_1, & m &= t_1 y_2 - t_2 y_1, & n &= t_1 z_2 - t_2 z_1, \\ l' &= y_1 z_2 - y_2 z_1, & m' &= z_1 x_2 - z_2 x_1, & n' &= x_1 y_2 - x_2 y_1. \end{aligned}$$

\* See Cayley, "On a new analytical representation of curves in space," *Papers*, 4 (1860), 446, where a curve is given by the complex of lines which meet it; Grassmann, *Ausdehnungslehre* (Berlin, 1862), §§ 63-65; Plücker, "On a new geometry of space," *Phil. Trans.* 155 (1865); Cayley, "On the six coordinates of a line," *Papers*, 7 (1869), 66.

From Grassmann it is clearly seen how to determine a system of coordinates for spaces of any dimension in a space of dimension  $n$ .

If we take the two points  $(x_1, y_1, z_1, t_1)$  and  $(x_2, y_2, z_2, t_2)$  of space and form the combinatory product

$$(e_1 x_1 + e_2 y_1 + e_3 z_1 + e_4 t_1)(e_1 x_2 + e_2 y_2 + e_3 z_2 + e_4 t_2),$$

where  $e_r^2 = 0$  and  $e_r e_s = -e_s e_r$ , we have a linear function of the six coordinates of the line joining the two points.

Given a space  $[k]$  in  $[n]$ , we take  $k + 1$  independent points of the  $[k]$  and write down the matrix of  $k + 1$  rows and  $n + 1$  columns formed by the coordinates of these points. Then the coordinates of  $[k]$  are simply the  $(k + 1)$ -rowed determinants of the matrix, which cannot all vanish if the points are independent. These determinants also arise in combinatory products (Grassmann, *ibid.*). They are sometimes spoken of as Grassmann coordinates.

The idea of a line in  $[3]$  being linearly dependent on the six edges of a tetrahedron occurs in Grassmann's *Ausdehnungslehre* (Leipzig, 1844), 167.



Then the six quantities  $(l, m, n, l', m', n')$  are called the *coordinates of the line* joining the two points. We are justified in speaking of them in this way because their mutual ratios are the same whatever two points of the line are taken; if instead of  $(x_1, y_1, z_1, t_1)$  and  $(x_2, y_2, z_2, t_2)$  we take the points

$$(\theta x_1 + \phi x_2, \theta y_1 + \phi y_2, \theta z_1 + \phi z_2, \theta t_1 + \phi t_2)$$

and

$$(\theta' x_1 + \phi' x_2, \theta' y_1 + \phi' y_2, \theta' z_1 + \phi' z_2, \theta' t_1 + \phi' t_2),$$

the six coordinates are each multiplied by  $\theta\phi' - \theta'\phi$ .

The lines whose coordinates satisfy a single relation are said to form a *complex* of lines; in particular, the linear relation

$$a'l + b'm + c'n + a'l' + b'm' + c'n' = 0$$

gives a *linear complex*\*. The lines whose coordinates satisfy simultaneously two linear relations are said to form a *linear congruence* of lines.

The six coordinates of any line satisfy the relation

$$ll' + mm' + nn' = 0.$$

Conversely, it can be shewn that any six quantities connected by this relation can be taken as the coordinates of a line.

**26.** If the line  $\lambda_1$  joining  $(x_1, y_1, z_1, t_1)$  to  $(x_2, y_2, z_2, t_2)$  has coordinates  $(l_1, m_1, n_1, l_1', m_1', n_1')$  and the line  $\lambda_2$  joining  $(\xi_1, \eta_1, \zeta_1, \tau_1)$  to  $(\xi_2, \eta_2, \zeta_2, \tau_2)$  has coordinates  $(l_2, m_2, n_2, l_2', m_2', n_2')$  the condition that the two lines should intersect is the same condition that the four points should be coplanar, or

$$\begin{vmatrix} x_1 & y_1 & z_1 & t_1 \\ x_2 & y_2 & z_2 & t_2 \\ \xi_1 & \eta_1 & \zeta_1 & \tau_1 \\ \xi_2 & \eta_2 & \zeta_2 & \tau_2 \end{vmatrix} = 0,$$

which is  $\varpi_{12} \equiv l_1 l_2' + m_1 m_2' + n_1 n_2' + l_1' l_2 + m_1' m_2 + n_1' n_2 = 0$ .

If this condition is satisfied, the six quantities

$$\begin{aligned} \kappa_1 l_1 + \kappa_2 l_2, \quad \kappa_1 m_1 + \kappa_2 m_2, \quad \kappa_1 n_1 + \kappa_2 n_2, \quad \kappa_1 l_1' + \kappa_2 l_2', \\ \kappa_1 m_1' + \kappa_2 m_2', \quad \kappa_1 n_1' + \kappa_2 n_2', \end{aligned}$$

satisfy the condition

$$\begin{aligned} (\kappa_1 l_1 + \kappa_2 l_2) (\kappa_1 l_1' + \kappa_2 l_2') + (\kappa_1 m_1 + \kappa_2 m_2) (\kappa_1 m_1' + \kappa_2 m_2') \\ + (\kappa_1 n_1 + \kappa_2 n_2) (\kappa_1 n_1' + \kappa_2 n_2') = 0 \end{aligned}$$

\* Geometrical properties of the linear complex were studied before the coordinates of a line were introduced. See Möbius, "Über eine besondere Art dualer Verhältnisse zwischen Figuren im Raume," *Journal für Math.* 10 (1833), 317, or *Gesammelte Werke*, 1, 491.

for all values of  $\kappa_1 : \kappa_2$ ; they are therefore the coordinates of a line which may be denoted symbolically by  $\kappa_1 \lambda_1 + \kappa_2 \lambda_2$ . Since the condition that two lines should intersect is linear in the coordinates of each line, any line which is met by both  $\lambda_1$  and  $\lambda_2$  must also be met by all the lines  $\kappa_1 \lambda_1 + \kappa_2 \lambda_2$ . Hence the line  $\kappa_1 \lambda_1 + \kappa_2 \lambda_2$  is a line passing through the intersection of  $\lambda_1$  and  $\lambda_2$  and lying in their plane; the different lines of this plane pencil are given by the different values of the ratio  $\kappa_1 : \kappa_2$ .

If we have three lines  $\lambda_1, \lambda_2, \lambda_3$  the relation

$$\begin{aligned} &(\kappa_1 l_1 + \kappa_2 l_2 + \kappa_3 l_3) (\kappa_1 l_1' + \kappa_2 l_2' + \kappa_3 l_3') \\ &+ (\kappa_1 m_1 + \kappa_2 m_2 + \kappa_3 m_3) (\kappa_1 m_1' + \kappa_2 m_2' + \kappa_3 m_3') \\ &+ (\kappa_1 n_1 + \kappa_2 n_2 + \kappa_3 n_3) (\kappa_1 n_1' + \kappa_2 n_2' + \kappa_3 n_3') = 0 \end{aligned}$$

is 
$$\varpi_{23} \kappa_2 \kappa_3 + \varpi_{31} \kappa_3 \kappa_1 + \varpi_{12} \kappa_1 \kappa_2 = 0.$$

There are  $\infty^1$  sets of values of  $\kappa_1 : \kappa_2 : \kappa_3$  satisfying this condition; and for such a set of values we have a line which may be represented symbolically by  $\kappa_1 \lambda_1 + \kappa_2 \lambda_2 + \kappa_3 \lambda_3$ . This meets all the lines which meet  $\lambda_1, \lambda_2$  and  $\lambda_3$  and we thus obtain one system of generators of a quadric surface.

This supposes that the lines  $\lambda_1, \lambda_2, \lambda_3$  are of general position; but if they all intersect one another we have  $\varpi_{23} = \varpi_{31} = \varpi_{12} = 0$ , and

$$\kappa_1 \lambda_1 + \kappa_2 \lambda_2 + \kappa_3 \lambda_3$$

is a line for all the  $\infty^2$  values of  $\kappa_1 : \kappa_2 : \kappa_3$ . Now  $\lambda_1, \lambda_2, \lambda_3$  are either concurrent or coplanar; if they are concurrent  $\kappa_1 \lambda_1 + \kappa_2 \lambda_2 + \kappa_3 \lambda_3$  will be a line through their intersection, and the  $\infty^2$  lines through their intersection are obtainable in this way; if they are coplanar  $\kappa_1 \lambda_1 + \kappa_2 \lambda_2 + \kappa_3 \lambda_3$  will be a line in their plane, and the  $\infty^2$  lines in their plane are obtainable in this way.

If we have a linear complex

$$a'l + b'm + c'n + al' + bm' + cn' = 0,$$

for which

$$aa' + bb' + cc' = 0,$$

the linear complex is said to be special. It consists of all the lines meeting the line whose coordinates are  $(a, b, c, a', b', c')$ .

27. *Line geometry in [3] considered as the point geometry of a quadric primal in [5].* If we have six quantities  $(l, m, n, l', m', n')$  satisfying the relation  $ll' + mm' + nn' = 0$  they can be regarded as the homogeneous coordinates of a point in [5] which lies on a quadric primal  $\Omega$ . This quadric  $\Omega$  is a general quadric; the left-hand side of its equation can be written as the sum of six squares.

Thus we have a correspondence between the lines of [3] and the points of  $\Omega$ ; to every line of [3] corresponds a point of  $\Omega$ , while to every point of  $\Omega$  corresponds a line of [3]. There are no exceptions. Hence *line geometry*

in [3] is exactly the same as the geometry of a quadric primal  $\Omega$  in [5]\*, and it will be found that great simplifications arise from this point of view.

28. If we take a definite point  $(\lambda, \mu, \nu, \lambda', \mu', \nu')$  of  $\Omega$  and join it to an arbitrary point  $(l, m, n, l', m', n')$  of [5] the coordinates of any point on the joining line are of the form

$(\theta\lambda + \phi l, \theta\mu + \phi m, \theta\nu + \phi n, \theta\lambda' + \phi l', \theta\mu' + \phi m', \theta\nu' + \phi n')$ , and this will lie on  $\Omega$  if

$$\theta^2 (\lambda\lambda' + \mu\mu' + \nu\nu') + \theta\phi (\lambda l' + \mu m' + \nu n' + \lambda' l + \mu' m + \nu' n) + \phi^2 (l l' + m m' + n n') = 0.$$

This is satisfied by  $\phi = 0$  as we should expect; it will have a double root  $\phi = 0$  if

$$l\lambda' + m\mu' + n\nu' + l'\lambda + m'\mu + n'\nu = 0.$$

This equation, linear in  $(l, m, n, l', m', n')$ , is the equation of a [4]; this [4] is such that the line joining any point of it to  $(\lambda, \mu, \nu, \lambda', \mu', \nu')$  touches  $\Omega$  at this last point. Or the lines which touch  $\Omega$  at any point lie in a prime—the *tangent prime* of  $\Omega$  at the point; and the equation to the tangent prime at  $(\lambda, \mu, \nu, \lambda', \mu', \nu')$  is

$$l\lambda' + m\mu' + n\nu' + l'\lambda + m'\mu + n'\nu = 0.$$

Clearly if a point  $P_1$  of  $\Omega$  lies in the tangent prime at a second point  $P_2$  of  $\Omega$  the point  $P_2$  will lie in the tangent prime at  $P_1$ . The line  $P_1P_2$  will then lie entirely on  $\Omega$ ; all tangents of  $\Omega$  which meet it in points other than their points of contact must lie entirely on  $\Omega$  as meeting it in at least three points. Two points of  $\Omega$  which are such that either lies in the tangent prime at the other will be spoken of as *conjugate*.

It is then clear from the condition  $\omega_{12} = 0$  that two lines  $\lambda_1, \lambda_2$  of [3] which intersect are represented on  $\Omega$  by two conjugate points  $P_1, P_2$ . Then the six quantities

$$\kappa_1 l_1 + \kappa_2 l_2, \quad \kappa_1 m_1 + \kappa_2 m_2, \quad \kappa_1 n_1 + \kappa_2 n_2, \quad \kappa_1 l'_1 + \kappa_2 l'_2, \quad \kappa_1 m'_1 + \kappa_2 m'_2, \\ \kappa_1 n'_1 + \kappa_2 n'_2$$

are the coordinates of a point of the line  $P_1P_2$ ; since this line lies on  $\Omega$ , the point must lie on  $\Omega$  and therefore represents a line of [3] for all values of  $\kappa_1 : \kappa_2$ . Hence the points of a line on  $\Omega$  represent the lines of a plane pencil in [3].

Corresponding to any line  $\lambda$  of [3] we have a point  $P$  of  $\Omega$ ; the lines of [3] which meet  $\lambda$  are represented on  $\Omega$  by its intersection with the tangent prime at  $P$ .

\* Klein, "Über Liniengeometrie und metrische Geometrie," *Math. Ann.* 5 (1872), 257, or *Gesammelte Mathematische Abhandlungen*, 1, 106; Cayley, *Papers*, 9 (1873), 79. The theory is greatly developed by Segre in his paper "Sulla geometria della retta e delle sue serie quadratiche," *Memorie Torino*, 36 (1883). See also Baker, *Principles of Geometry*, 4 (Cambridge, 1925), 40 et seq.

The lines of a linear complex are represented by the points of a prime section of  $\Omega$ ; but for a special linear complex we must take the section by a tangent prime.

29. *The planes\* on  $\Omega$ .* Consider now three points  $P_1, P_2, P_3$  of  $\Omega$ , each pair of which is a pair of conjugate points. Then these represent three lines  $\lambda_1, \lambda_2, \lambda_3$  of [3] for which  $\omega_{23} = \omega_{31} = \omega_{12} = 0$ . The lines  $P_2P_3, P_3P_1, P_1P_2$  all lie on  $\Omega$ ; so that the plane  $P_1P_2P_3$  meets  $\Omega$  in a curve whose order is greater than two, and lies on  $\Omega$  entirely. Then the point whose six coordinates are of the form  $\kappa_1l_1 + \kappa_2l_2 + \kappa_3l_3$  lies on  $\Omega$  for all values of  $\kappa_1 : \kappa_2 : \kappa_3$  and represents a line  $\kappa_1\lambda_1 + \kappa_2\lambda_2 + \kappa_3\lambda_3$  of [3]. We thus obtain every point of the plane  $P_1P_2P_3$ .

We thus have two systems of planes on  $\Omega$ ; the points of a plane of the first system represent the lines through a point of [3], while the points of a plane of the second system represent the lines in a plane of [3]. We shall call the planes of the first system, representing lines through points of [3],  $\omega$ -planes and the planes of the second system, representing lines in planes of [3],  $\rho$ -planes. Since there is one and only one line passing through two given points of [3] two  $\omega$ -planes have one and only one point in common; and since there is one and only one line lying in two given planes of [3] two  $\rho$ -planes have one and only one point in common; two planes of the same system on  $\Omega$  meet in a point. Given a point and a plane of [3], there will not be a line passing through the point and lying in the plane unless the point itself lies in the plane, so that two planes of  $\Omega$  of opposite systems do not in general intersect. If however the point does lie in the plane we have a pencil of lines passing through the point and lying in the plane; hence if two planes of opposite systems on  $\Omega$  do intersect they have a line in common.

On  $\Omega$  there are  $\infty^3$  planes of each system; through any point of  $\Omega$  there pass  $\infty^1$  of each system. There are also on  $\Omega$   $\infty^5$  lines,  $\infty^2$  passing through any given point on  $\Omega$ †.

30. *The quadric point-cone in [4].* We can now give a detailed description of the section of  $\Omega$  by the tangent prime at a point  $P$ . The point  $P$  represents a line  $\lambda$  of [3]; on  $\lambda$  there are  $\infty^1$  points and through  $\lambda$  there pass  $\infty^1$  planes. Hence there are  $\infty^1$   $\omega$ -planes on  $\Omega$  and  $\infty^1$   $\rho$ -planes on  $\Omega$  all passing through  $P$ . Two of these planes which belong to the same system will not intersect except in  $P$ , but two planes of opposite systems

\* The two systems of planes on  $\Omega$  are mentioned explicitly by Cayley, *Papers*, 9 (1873), 79.

† Cf. Segre, "Studio sulle quadriche in uno spazio lineare ad un numero qualunque di dimensione," *Memorie Torino*, 36 (1883), in particular p. 36.

will intersect in a line through  $P$ . The section of  $\Omega$  by the tangent prime at  $P$  is nothing but these two sets of planes, since any point of  $\Omega$  lying in the tangent prime represents a line of [3] meeting  $\lambda$ . Such a line in [3] meets  $\lambda$  in a point and is joined to  $\lambda$  by a plane, so that the representative point on  $\Omega$  lies in two planes, one of each system. The line of intersection of these two planes joins the representative point to  $P$ .

If we take a section by a [3] lying in the tangent prime and not passing through  $P$  we obtain two systems each of  $\infty^1$  lines; two lines of the same system do not intersect but every line of either system meets every line of the other; in other words, we have the two systems of generators of a quadric surface. Thus the section of  $\Omega$  by a tangent prime is the same locus as is obtained by joining the generators of a quadric surface, by planes, to a point outside the [3] in which the quadric surface lies. This locus in [4] is called a *quadric point-cone*\*.

If we have a curve lying on a quadric point-cone in [4] and project it on to a [3] from the vertex of the point-cone we obtain a curve lying on the quadric in which [3] meets the point-cone. Thus, from a knowledge of the properties of this projected curve, we shall be able to deduce certain properties of the curve on the cone.

The section of a quadric point-cone by a [3] through its vertex is an ordinary quadric cone, unless the [3] is that determined by two planes of opposite systems of the point-cone; it then meets the point-cone simply in these two planes. Such a [3] is called a *tangent solid* of the point-cone; there are  $\infty^3$  [3]'s through the vertex,  $\infty^2$  of which are tangent solids.

**31. The representation of a ruled surface.** Since a line of [3] is represented by a point of  $\Omega$ , a ruled surface  $f$  in [3] formed by  $\infty^1$  lines will be represented by the curve  $C$  on  $\Omega$  formed by the  $\infty^1$  representative points†. There is thus a (1, 1) correspondence between  $C$  and the generators of  $f$ , so that the genus of  $C$  is equal to the genus of  $f$ . Moreover, the order of the ruled surface, being equal to the number of its generators which meet an arbitrary line  $\lambda$  of [3], is equal to the number of intersections of  $C$  with the tangent prime at an arbitrary point  $P$  of  $\Omega$ , and this is simply the order of  $C$ . Thus we can say that *a ruled surface  $f$  of order  $n$  and genus  $p$  in [3] is represented on  $\Omega$  by a curve  $C$  of order  $n$  and genus  $p$ .*

We thus see how to begin the classification of ruled surfaces mentioned in § 12. We have first to investigate how the double curve and bitangent developable can be studied by means of the curve  $C$  on  $\Omega$ . Then, taking  $C$  to be of a given order and genus, different positions of  $C$  on  $\Omega$  will give different kinds of double curves and bitangent developables for  $f$ .

\* Cf. Baker, *Principles of Geometry*, 4 (Cambridge, 1925), 120–121.

† Cf. Voss, "Zur Theorie der windschiefen Flächen," *Math. Ann.* 8 (1874), 54. Segre, *Memorie Torino*, 36 (1883), 97.

A given generator  $g$  of  $f$  is represented on  $\Omega$  by a point  $P$  of  $C$ . All generators of  $f$  which meet  $g$  are represented on  $\Omega$  by points of  $C$  which lie in the tangent prime of  $\Omega$  at  $P$ ; but this tangent prime, meeting  $C$  twice at  $P$ , will meet it in  $n - 2$  other points, where  $n$  is the order of  $f$  and  $C$ . Hence every generator of  $f$  is met by  $n - 2$  others (cf. § 10).

**32. Torsal generators.** If we set up a correspondence between the points  $P$  and  $Q$  of  $C$ , saying that two points  $P$  and  $Q$  correspond when the chord  $PQ$  lies on  $\Omega$ , we have a symmetrical  $(n - 2, n - 2)$  correspondence. This correspondence is of valency 2 since the  $n - 2$  points  $Q$  which correspond to any point  $P$ , when taken together with  $P$  counted twice, form the complete intersection of  $C$  with a prime. Hence the number of coincidences in the correspondence is

$$n - 2 + n - 2 + 2 \cdot 2 \cdot p = 2(n + 2p - 2),$$

where  $p$  is the genus of  $C$ .

This means that, for  $2(n + 2p - 2)$  points  $P$  of  $C$ , there are only  $n - 3$  chords of  $C$  passing through  $P$  and lying on  $\Omega$ ; the tangent of  $C$  at  $P$  lies on  $\Omega$ , and the tangent prime of  $\Omega$  at  $P$  meets  $C$  in three points there.

Then on the ruled surface  $f$  we can say that there are  $2(n + 2p - 2)$  generators which meet their "consecutive generators." Such generators are called *torsal generators*; the tangent plane to the ruled surface is the same for all points of such a generator. Thus a ruled surface of order  $n$  and genus  $p$  has, in general\*,  $2(n + 2p - 2)$  torsal generators.

Incidentally we have proved that, if a curve of order  $n$  and genus  $p$  lies on a quadric primal in space of any number of dimensions, there are  $2(n + 2p - 2)$  tangents of the curve lying on the quadric\*. This is easily verified for simple curves on an ordinary quadric surface in [3].

**33. The double curve and bitangent developable.** The degree of the double curve of the ruled surface  $f$  in [3] is the number of points in which it meets an arbitrary plane. Now the lines of such a plane are represented by the points of a  $\rho$ -plane on  $\Omega$ , while the lines (including two generators of  $f$ ) which pass through a point of the double curve are represented by the points of a  $\omega$ -plane (meeting  $C$  in two points). Thus corresponding to each intersection of the double curve with a definite plane of [3] we have a  $\omega$ -plane meeting  $C$  twice and meeting a definite  $\rho$ -plane in a line; or we have a chord of  $C$  lying on  $\Omega$  and meeting the  $\rho$ -plane. Conversely, corresponding to each chord of  $C$  which meets a definite  $\rho$ -plane (and incidentally lies on  $\Omega$  as meeting it in at least three points) we have two generators of  $f$  intersecting on a definite plane of [3]. Thus the order of the double curve of  $f$  is equal to the number of chords of  $C$  which meet any given  $\rho$ -plane of

\* If  $C$  has  $\kappa$  cusps then the number of tangents of  $C$  which lie on  $\Omega$  is only  $2(n + 2p - 2) - 2\kappa$ , see § 349 below.

general position. Similarly, the class of the bitangent developable of  $f$  is equal to the number of chords of  $C$  which meet any given  $\omega$ -plane of general position. These two results are fundamental.

The chords of  $C$  which lie on  $\Omega$  form a ruled surface  $R_2$  on which  $C$  is a multiple curve of multiplicity  $n - 2$ . The two points of  $C$  on a chord which lies on  $\Omega$  represent two intersecting generators of  $f$ , and conversely. The generators of  $R_2$  are thus in (1, 1) correspondence both with the points of the double curve and with the planes of the bitangent developable of  $f$ . A prime section of  $R_2$  is thus a curve whose genus is equal to that of the double curve of  $f$  and also to that of the bitangent developable of  $f$ . We can calculate this genus in the general case.

Denote a prime section of  $R_2$  by  $C'$ , and consider the  $(2, n - 2)$  correspondence between  $C$  and  $C'$ , two points of  $C$  and  $C'$  corresponding when the line joining them is a chord of  $C$ . The number of branch-points of the correspondence on  $C'$  is simply the number of tangents of  $C$  which lie on  $\Omega$ ; this we have seen to be  $2(n + 2p - 2)$ . The number of branch-points of the correspondence on  $C$  is equal to the number of points  $P$  of  $C$  at which two of the  $n - 2$  generators of  $R_2$  coincide. This is equal to the number of times two of the  $n - 2$  points  $Q$  coincide in an  $(n - 2, n - 2)$  correspondence of valency 2 on a curve of genus  $p$ ; this number\* is

$$2(n - 2)(n - 3) + 2(n - 6)p.$$

Hence, applying Zeuthen's formula to the correspondence between  $C$  and  $C'$ , we have, if  $P$  is the genus of  $C'$ ,

$$\begin{aligned} 2(n - 2)(n - 3) + 2(n - 6)p - 2(n + 2p - 2) \\ = 4(P - 1) - 2(n - 2)(p - 1), \end{aligned}$$

or 
$$2P - 2 = (n - 5)(n + 2p - 2) \dagger,$$

giving, for a ruled surface of order  $n$  and genus  $p$ , the genus  $P$  of the double curve and the bitangent developable.

34. When the ruled surface  $f$  in [3] is not completely general for its order and genus the double curve and bitangent developable may break up; when this happens the ruled surface  $R_2$  formed by the chords of  $C$  lying on  $\Omega$  will have to break up correspondingly. For each part of the double curve and bitangent developable we have a set of pairs of intersecting generators of  $f$  and thus a set of chords of  $C$  lying on  $\Omega$  and forming

\* See § 14.

† This result is deducible from formulæ given by Salmon for the theory of reciprocal surfaces; see his *Geometry of Three Dimensions*, 2 (Dublin, 1915), 301. The application to a ruled surface is given by Cayley, *Papers*, 8 (1871), 396. If  $q$  is the rank and  $b$  the order of the double curve

$$q = 2b + 2P - 2 \text{ and } b = \frac{1}{2}(n - 1)(n - 2) - p.$$

The actual form of the result as here stated is given by Wiman, *Acta Mathematica*, 19 (1865), 66.

a ruled surface, say  $S_2$ , which is the corresponding part of  $R_2$ . The order of this part of the double curve of  $f$  is equal to the number of generators of  $S_2$  which meet a definite  $\rho$ -plane, and the class of the corresponding part of the bitangent developable of  $f$  is equal to the number of generators of  $S_2$  which meet a definite  $\varpi$ -plane. The genus of the prime sections of  $S_2$  gives the genus of these parts of the double curve and bitangent developable.

As well as investigating the properties of the component parts of the double curve we can investigate their relations with one another, and similarly for the parts of the bitangent developable. To do this we have to study the relations between the different component ruled surfaces which constitute  $R_2$ .

35. Given a ruled surface in [3] there is, in general, a finite number of points which are *triple points* of the surface; through such points there pass three generators of the surface and they are also triple points of the double curve. Similarly we have, in general, a finite number of planes which are *tritangent planes* of the surface; in such planes there lie three generators of the surface and they are triple planes of the bitangent developable.

When the ruled surface is represented as a curve  $C$  on  $\Omega$  the triple points give  $\varpi$ -planes of  $\Omega$  trisecant to  $C$ , while the tritangent planes give  $\rho$ -planes of  $\Omega$  trisecant to  $C$ ; there is a finite number of trisecant planes of  $C$  lying on  $\Omega$ . There are  $\infty^3$  planes on  $\Omega$ ; and since  $\Omega$  is a four-dimensional locus one condition will be necessary in order that a plane and a curve which lie on it should intersect; thus three conditions must be imposed on a plane of  $\Omega$  to make it a trisecant plane of  $C$ , so that we naturally expect a finite number of such planes.

Denote for the moment the curve  $C$  of order  $n$  and genus  $p$  by  $C_n^p$ ; the number of its trisecant planes lying on  $\Omega$  can be calculated directly by correspondence theory.

If we take any point  $X$  on  $C_n^p$  there are  $n - 2$  chords of  $C_n^p$  passing through it which lie on  $\Omega$ ; these meet  $C_n^p$  again in points  $X_1, X_2, \dots, X_{n-2}$ . We have already noticed that the correspondence between  $X$  and  $X_r$  is a symmetrical  $(n - 2, n - 2)$  correspondence of valency 2.

In the same way each point  $X_r$  gives rise to  $n - 2$  points

$$X, X_r^{(1)}, \dots, X_r^{(n-3)}.$$

Then the correspondence between  $X$  and  $X_r^{(s)}$  is also symmetrical and both its indices are  $(n - 2)(n - 3)$ . The points corresponding to  $X$  are found by taking the square\* of the former correspondence and leaving

\* See Severi, *Memorie Torino* (2), 54 (1904), 5-9.



out the point  $X$  each of the  $n - 2$  times it occurs. Hence the valency is\*

$$- (2)^2 + n - 2 = n - 6.$$

Since both the correspondences are symmetrical the number of common pairs of points is half that given by Brill's formula†, and is therefore

$$\begin{aligned} N &= \frac{1}{2} \{ (n - 2)^2 (n - 3) + (n - 2)^2 (n - 3) - 2 \cdot 2 (n - 6) p \} \\ &= (n - 2)^2 (n - 3) - 2 (n - 6) p. \end{aligned}$$

This result means that there are  $N$  points  $X$  on  $C_n^p$  such that some point  $X_\alpha$  coincides with some point  $X_\beta^{(\gamma)}$ . In particular the suffixes  $\alpha$  and  $\beta$  may be the same; and then the tangent of  $C_n^p$  at  $X_\alpha$  lies on  $\Omega$ . There are  $2(n + 2p - 2)$  such points on  $C_n^p$ ; and through each one of these there pass  $n - 3$  proper chords of  $C_n^p$  which lie on  $\Omega$ ; each of these chords meets  $C_n^p$  again in a point  $X$  which is included in the above  $N$  points. Thus the remaining points  $X$  are in number

$$\begin{aligned} (n - 2)^2 (n - 3) - 2 (n - 6) p - 2 (n - 3) (n + 2p - 2) \\ = (n - 2) (n - 3) (n - 4) - 6 (n - 4) p. \end{aligned}$$

Then for any one of these points  $X$  we have  $X_\alpha$  and  $X_\beta^{(\gamma)}$  coinciding, where  $\alpha$  and  $\beta$  are not the same. This means that the chords  $XX_\alpha$ ,  $XX_\beta$ ,  $X_\alpha X_\beta$  all lie on  $\Omega$ ; so that  $XX_\alpha X_\beta$  is a trisecant plane of  $C_n^p$  lying on  $\Omega$ .

Thus these points  $X$  occur in groups of three, each group determining a trisecant plane of  $C_n^p$  which lies on  $\Omega$ . Hence the number of trisecant planes of  $C_n^p$  which lie on  $\Omega$  is

$$(n - 4) \left\{ \frac{1}{3} (n - 2) (n - 3) - 2p \right\}.$$

These trisecant planes of  $C$  which lie on  $\Omega$  belong half to one system of planes and half to the other. Thus, given a ruled surface of order  $n$  and genus  $p$  in [3], the number of its triple points is‡

$$\frac{1}{6} (n - 2) (n - 3) (n - 4) - (n - 4) p,$$

and this is also the number of its tritangent planes.

36. A conic  $C$  on  $\Omega$  represents a ruled surface in [3] of the second order, or the points of  $C$  represent *one system* of generators of a quadric surface. We have in fact already noticed¶ that such a system of lines in [3] is represented by coplanar points on  $\Omega$ . It is implied that the plane of  $C$  does not lie entirely on  $\Omega$ . The coordinates of any point of  $C$  being of the form  $\kappa_1 l_1 + \kappa_2 l_2 + \kappa_3 l_3$ , the equation to the tangent prime there is

$$\kappa_1 T_1 + \kappa_2 T_2 + \kappa_3 T_3 = 0,$$

\* Severi, *ibid.* The "product" of two correspondences whose valencies are  $\gamma_1$  and  $\gamma_2$  has the valency  $-\gamma_1 \gamma_2$ .

† § 15.

‡ See the references to Cayley and Wiman in § 33. Also Castelnuovo, *Palermo Rendiconti*, 3 (1889), 33.

¶ § 26.

where  $T_1 = 0$ ,  $T_2 = 0$ ,  $T_3 = 0$  are the tangent primes at three definite points of  $C$ , so that the tangent primes of  $\Omega$  at all the points of  $C$  have in common the plane  $T_1 = T_2 = T_3 = 0$ . This plane meets  $\Omega$  in a second conic  $C'$ , and the tangent primes of  $\Omega$  at all the points of  $C'$  all contain the plane of  $C$ . The lines represented by the points of  $C'$  meet all the lines represented by the points of  $C$  and conversely; thus  $C$  and  $C'$  give in [3] complementary reguli on the same quadric surface. There is no double curve or bitangent developable.

If the plane of  $C$  is a  $\pi$ -plane we have the generators of a quadric cone, and if it is a  $\rho$ -plane we have the tangents of a plane conic. Clearly we can have point-cones and plane-envelopes represented on  $\Omega$  in this way by curves of all orders and genera; in future these will be ignored.

**37. Cubic ruled surfaces.** As an application of the theory we will investigate the ruled surfaces of the third order in [3]. For this we have to consider cubic curves on  $\Omega$ , and the only relevant curve is the twisted cubic  $C$  of  $S_3$ . The plane cubic is irrelevant, because if a plane cubic curve lies on a quadric the whole of its plane must do so, and we have the case mentioned at the end of § 36.

In general,  $S_3$  meets  $\Omega$  in a quadric surface  $Q$ ;  $C$  will meet all the generators of one system of  $Q$  in two points and all of the other system in one point\*. Through  $S_3$  there pass two tangent primes of  $\Omega$ , touching it in two points  $O$  and  $O'\dagger$ . These two points represent two lines  $R$  and  $R'$ , and every generator of the surface meets these two lines. The surface has therefore two directrices, where we define a directrix to be a line which is met by every generator of the surface.

The tangent prime at  $O$  meets  $\Omega$  in a quadric point-cone containing  $Q$ ; the two systems of planes on the point-cone passing through the two systems of generators on  $Q$ . We may suppose that that system of generators which are chords of  $C$  lie in the  $\pi$ -planes through  $O$ ; these same generators will then lie in the  $\rho$ -planes through  $O'$ .

Hence through any point of  $R$  there pass two generators lying in a plane through  $R'$ , while any plane through  $R'$  contains two generators meeting in a point of  $R$ .

We have thus established a geometrical connection between the facts that

(a) a twisted cubic on a quadric meets all the generators of one system in two points and all of the other system in one point;

(b) a cubic ruled surface has two directrices; through each point of one there pass two generators, while through each point of the other there passes one generator.

\* Salmon, *Geometry of Three Dimensions*, 1 (Dublin, 1914), 347.

† Just as there are two tangent planes of an ordinary quadric passing through a line of [3]. The poles of the primes which pass through  $S_3$  all lie on a line; this meets  $\Omega$  in the two points  $O$ ,  $O'$  at which the tangent primes contain  $S_3$ .

The double curve consists of the points of  $R$ , the bitangent developable of the planes through  $R'$ . An arbitrary plane of  $\Omega$  meets  $S_3$  in a point of  $Q$ , and there is only one chord of  $C$  lying on  $\Omega$  and meeting this plane—that generator of  $Q$  which meets  $C$  twice and passes through the point where the plane meets  $S_3$ . Thus the double curve is of the first order and the bitangent developable of the first class.

38. It may happen however that  $S_3$  occupies a special position in regard to  $\Omega$ , meeting it in a quadric cone with vertex  $V$ . Then only one tangent prime of  $\Omega$  passes through  $S_3$ , this being the tangent prime at  $V^*$ . A cubic curve on the cone necessarily passes through  $V$ . Hence the second species of cubic ruled surface has a directrix line  $R$  which is also a generator. Any plane of  $\Omega$  passing through  $V$  lies in the tangent prime at  $V$  and therefore meets  $S_3$  in a line. This line will be a generator of the cone and will meet the curve  $C$  in one point other than  $V$ . Thus through any point of  $R$  there passes one generator other than  $R$ , while any plane through  $R$  contains one generator other than  $R$ .

This is in fact the cubic ruled surface of § 20.

We have thus obtained the two kinds of cubic ruled surfaces in [3] given by Cayley †.

### SECTION III

## THE PROJECTION OF RULED SURFACES FROM HIGHER SPACE

39. Just as a curve is said to be normal when it cannot be obtained by projection from a curve of the same order in space of higher dimension, so a ruled surface is said to be normal when it cannot be obtained by projection from a ruled surface of the same order in space of higher dimension.

It is well known how the descriptive theory of curves has been amplified and simplified by considering curves as the projections of normal curves, and it is natural to expect that the theory of ruled surfaces will benefit similarly by considering ruled surfaces as projections of normal ruled surfaces. There are, however, as we should again naturally expect, more complicated relations to consider in the theory of ruled surfaces than in the theory of curves; for example, the normal space for a curve of order  $n$  and genus  $p$  is unique so long as  $n > 2p - 2$ , but for a ruled surface of order  $n$  and genus  $p$  the normal space is only unique when  $p = 0$  and

\* The poles of the primes containing  $S_3$  now lie on a line which touches  $\Omega$  at  $V$ .

† *Papers*, 5, 212–213.

$p = 1$ . Again, rational curves of order  $n$  are all projectively equivalent, but rational ruled surfaces of order  $n$  are not all projectively equivalent.

**40.** We shall, from this until § 50, be concerned solely with ruled surfaces which are rational. A rational ruled surface of order  $n$  is normal in  $[n + 1]^*$ , all rational ruled surfaces of order  $n$  can be obtained as projections of these normal surfaces.

In the first place no surface of order  $n$ , whether ruled or not, can lie in a space of dimension greater than  $n + 1$  †. If it did so every prime section would be a curve lying in a space of dimension greater than its order.

In the second place, given any rational ruled surface of order  $n$ , there exists in  $[n + 1]$  a normal surface of which it is the projection. For consider such a surface in  $[m]$ , where  $2 < m < n + 1$ , and take two prime sections. If the primes are of general position in regard to the surface the sections will both be rational curves of order  $n$ ; the  $m + 1$  homogeneous coordinates of a point on either curve can be expressed as polynomials of order  $n$  in a parameter. Since the curves are placed in (1, 1) correspondence by the generators of the surface we can take the same parameter  $\lambda$  for both curves; so that for a point of the first curve

$$x_i = \phi_i(\lambda) \quad (i = 0, 1, \dots, m),$$

and for a point of the second curve

$$x_i = \psi_i(\lambda) \quad (i = 0, 1, \dots, m),$$

where  $\phi$  and  $\psi$  are polynomials of order  $n$ .

Thus the coordinates of a point of the ruled surface can be expressed as rational functions of two parameters  $\lambda$  and  $\nu$  in the form

$$x_i = \phi_i(\lambda) + \nu\psi_i(\lambda) \quad (i = 0, 1, \dots, m).$$

Now the two prime sections have  $n$  common points, viz. the points in which the surface is met by the  $[m - 2]$  common to the two primes. Then for these points the polynomials  $\phi_i$  must be proportional to the polynomials  $\psi_i$ , so that

$$\frac{\phi_1(\lambda_j)}{\psi_1(\lambda_j)} = \frac{\phi_2(\lambda_j)}{\psi_2(\lambda_j)} = \dots = \frac{\phi_m(\lambda_j)}{\psi_m(\lambda_j)} = k_j \quad (j = 1, 2, \dots, n - 1, n),$$

where  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the  $n$  values of  $\lambda$  giving the common points of the two prime sections. We can write the coordinates of a point on the ruled surface in the homogeneous form

$$x_i = \mu\phi_i(\lambda, \mu) + \nu\psi_i(\lambda, \mu) \quad (i = 0, 1, \dots, m).$$

Now let us choose further pairs of polynomials  $\phi$  and  $\psi$  of degree  $n$  in  $\lambda$ , the values of the ratios  $\phi/\psi$  being also equal to the quantities  $k$  for the

\* Veronese, *Math. Ann.* 19 (1882), 228.

† Veronese, *ibid.* 166.

$n$  values of  $\lambda$  giving the common points of the two prime sections; and then take further expressions

$$x_i = \mu\phi_i(\lambda, \mu) + \nu\psi_i(\lambda, \mu) \quad (i = m + 1, \dots).$$

If  $(\lambda, \mu, \nu)$  are regarded as the homogeneous co-ordinates of a point in a plane the quantities  $x_i$  when equated to zero represent curves of order  $n + 1$ ; these curves all have:

- (a) a multiple point of order  $n$  at  $\lambda = \mu = 0$ ,
- (b) a common point at  $\mu = \nu = 0$ ,
- (c) common points at the  $n$  points  $\mu = 1, \lambda = \lambda_j, \nu = -k_j$ .

Thus the number of coordinates  $x_i$  it is possible to choose which are linearly independent is the same as the number of linearly independent plane curves of order  $n + 1$  which have in common  $n + 1$  ordinary points and have also a given point of multiplicity  $n$ . But the number of such curves is

$$\frac{1}{2}(n + 1)(n + 4) + 1 - (n + 1) - \frac{1}{2}n(n + 1) = n + 2.$$

We can therefore choose linearly independent coordinates

$$x_i = \mu\phi_i(\lambda, \mu) + \nu\psi_i(\lambda, \mu) \quad (i = 0, 1, \dots, n + 1),$$

$n + 2$  in number and no more.

Then these quantities  $x_i$  are the coordinates of a point on a rational ruled surface of order  $n$  in  $[n + 1]$ , the generators being given by

$$\lambda/\mu = \text{const.}$$

This surface is normal and cannot be obtained by projection from a ruled surface of the same order in higher space. The original surface lies in the  $[m]$  whose equations are  $x_{m+1} = x_{m+2} = \dots = x_{n+1} = 0$  and is the projection of the normal surface from the  $[n - m]$  whose equations are

$$x_0 = x_1 = \dots = x_m = 0.$$

To a general point of the ruled surface there corresponds one point of the plane, while to a general point of the plane there corresponds one point of the ruled surface. The correspondence between the plane and ruled surface is birational save for a certain number of exceptional points.

To a prime section of the ruled surface corresponds on the plane a curve of order  $n + 1$  with a fixed  $n$ -ple point and  $n + 1$  other fixed points. To the points in which the surface is met by a space of dimension two less than the space to which it belongs we have the variable intersections of two such plane curves; the number of these is

$$(n + 1)^2 - n^2 - n - 1 = n,$$

which is, as it should be, the order of the ruled surface.

A ruled surface of order  $n$  in  $[n + 1]$  is necessarily a rational surface because its prime sections are rational curves.

It can be shewn\* that every surface of order  $n$  in  $[n + 1]$  is necessarily a ruled surface except when  $n = 4$ .

41. The properties of the rational normal ruled surfaces were first studied by Segre in his paper on rational ruled surfaces published at Turin in 1884 †. The results obtained by him are fundamental for our work, and as we shall be using them constantly we give here some account of them.

In the first place we can immediately establish the existence of rational ruled surfaces of order  $n$  in  $[n + 1]$  of several types which are projectively distinct. For take two spaces  $[m]$  and  $[n - m]$  in  $[n + 1]$  which do not intersect; in  $[m]$  take a rational normal curve of order  $m$  and in  $[n - m]$  take a rational normal curve of order  $n - m$ . Then if the two curves are placed in (1, 1) correspondence the ruled surface formed by joining pairs of corresponding points is of order  $n$  ‡; it is rational and no two of its generators can intersect §.

Take homogeneous coordinates  $(x_0, x_1, \dots, x_{n+1})$  so that the equations of  $[m]$  are  $x_{m+1} = x_{m+2} = \dots = x_{n+1} = 0$  and the equations of  $[n - m]$  are  $x_0 = x_1 = \dots = x_m = 0$ . Then a point of the curve in  $[m]$  can be given by

$$x_0 = \lambda^m, x_1 = \lambda^{m-1}, \dots, x_m = 1, x_{m+1} = 0, x_{m+2} = 0, \dots, x_{n+1} = 0,$$

while the corresponding point of the curve in  $[n - m]$  is given by

$$x_0 = 0, x_1 = 0, \dots, x_m = 0, x_{m+1} = \lambda^{n-m}, x_{m+2} = \lambda^{n-m-1}, \dots, x_{n+1} = 1.$$

The coordinates of a point on the ruled surface are then given rationally in terms of two parameters  $\lambda$  and  $\mu$  by the equations

$$x_0 = \lambda^m, x_1 = \lambda^{m-1}, \dots, x_m = 1, x_{m+1} = \lambda^{n-m}\mu, x_{m+2} = \lambda^{n-m-1}\mu, \dots, x_{n+1} = \mu,$$

so that the coordinates of every point of the ruled surface satisfy the equations

$$\frac{x_0}{x_1} = \frac{x_1}{x_2} = \dots = \frac{x_{m-1}}{x_m} = \frac{x_{m+1}}{x_{m+2}} = \dots = \frac{x_n}{x_{n+1}};$$

and conversely every point whose coordinates satisfy these equations is a point of the ruled surface.

The equations to the ruled surface are therefore

$$\left\| \begin{array}{cccccc} x_0 & x_1 & \dots & x_{m-1} & x_{m+1} & \dots & x_n \\ x_1 & x_2 & \dots & x_m & x_{m+2} & \dots & x_{n+1} \end{array} \right\| = 0.$$

In particular, a ruled surface of order 2 in  $[3]$  is obtained by a (1, 1) correspondence between two skew lines. If the lines are  $x_0 = x_1 = 0$  and  $x_2 = x_3 = 0$  the equation to the ruled surface can be taken as

$$\frac{x_0}{x_1} = \frac{x_2}{x_3}.$$

\* Del Pezzo, "Sulle superficie di ordine  $n$  immerse nello spazio di  $n + 1$  dimensioni," *Rend. dell' Accad. di Napoli*, 24 (1885), 212.

† "Sulle rigate razionali in uno spazio lineare qualunque," *Atti Torino*, 19 (1884), 355.

‡ § 19.

§ For if they did the spaces  $[n]$  and  $[n - m]$  would also intersect.

42. We shall always assume that the ruled surface in  $[n + 1]$  is not a cone and that it does not break up into separate surfaces. Further, it does not lie in any space of dimension less than  $n + 1$ .

If there is on the surface a curve of order less than or equal to  $n$  it can only meet each generator of the surface in one point; for if it met each generator in more than one point any space containing the curve would contain the whole surface. But the surface lies in a space of dimension greater than  $n$ , whereas the curve must lie in a space of dimension less than or equal to its order. Since then the curve only meets each generator in one point it is a rational curve.

A curve on a ruled surface meeting every generator in one point will be called a *directrix*.

Thus on a ruled surface of order  $n$  in  $[n + 1]$  every curve  $C_\mu$  of order  $\mu < n$  is a rational curve. But, further, every such curve is a rational normal curve. For if it were contained in a  $[\mu - b]$ , where  $b > 0$ , we could take, through the curve and through any  $n - \mu + b$  generators, a space  $[n]$  which would meet the surface in a curve of order

$$\mu + n - \mu + b = n + b > n;$$

and this is impossible.

Moreover, we cannot have on the surface two curves the sum of whose orders is less than  $n$ . For if we have two curves  $C_\mu$  and  $C_{\mu'}$  of respective orders  $\mu$  and  $\mu'$ , where  $\mu + \mu' < n$ , they are contained in spaces  $[\mu]$  and  $[\mu']$ : whence the space  $[\mu + \mu' + 1]$  containing these would contain the whole surface, which is not contained in a space of dimension less than  $n + 1$ .

Similarly, if we have on the surface two curves the sum of whose orders is  $n$  they cannot intersect.

43. When we generate a normal ruled surface by means of two curves of orders  $m$  and  $n - m$  we can always suppose, except when  $m = n - m = \frac{1}{2}n$ , that

$$m < n - m,$$

or

$$m < \frac{n}{2}.$$

Then there can be no other curve on the surface of order as small as  $m$ ; or the curve of order  $m$  is the *directrix of minimum order* on the surface. We can call it for brevity the *minimum directrix*.

Now in generating the surface we can clearly take any value for  $m$  such that  $1 < m < \frac{n}{2}$  (ignoring  $m = 0$  which gives a cone). Two surfaces whose minimum directrices are of different orders cannot be projectively equivalent.

Thus if  $n$  is odd we have  $\frac{n-1}{2}$  projectively distinct types of rational normal ruled surfaces of order  $n$  in  $[n+1]$ ; each surface has a directrix of minimum order equal to  $m$ , where  $m$  is one of the numbers  $1, 2, \dots, \frac{n-1}{2}$ .

If  $n$  is even we have  $\frac{n}{2}$  projectively distinct types of rational normal ruled surfaces of order  $n$  in  $[n+1]$ ; one surface has minimum directrices of order  $\frac{n}{2}$ , while the others have each a directrix of minimum order equal to  $m$ , where  $m$  is one of the numbers  $1, 2, \dots, \frac{n}{2} - 1$ .

**44.** We now shew that every rational normal ruled surface of order  $n$  in  $[n+1]$  is of one of these types. To do this we must shew:

(a) that every ruled surface of order  $n$  in  $[n+1]$  has on it a curve of order  $m < \frac{n}{2}$ ;

(b) that such a surface has on it curves of order  $n - m$ .

Then taking curves  $C_m$  and  $C_{n-m}$  of these orders they cannot have any intersections, and the ruled surface can be given by a  $(1, 1)$  correspondence between these curves.

The proof of (a) is immediate; for taking a ruled surface of order  $n$  in  $[n+1]$  we know that a prime  $[n]$  can be taken to contain  $n+1$  independent points. If then  $n$  is odd we can take a prime through  $\frac{n+1}{2}$  generators arbitrarily chosen; the remaining intersection of the prime with the surface is a curve of order  $\frac{n-1}{2}$  which may or may not contain other generators as parts of itself. It always includes, however, a curve which is a directrix; so that on a ruled surface of odd order  $n$  in  $[n+1]$  there is always a directrix curve of order less than or equal to  $\frac{n-1}{2}$ . Similarly, on a ruled surface of even order  $n$  in  $[n+1]$  there is always a directrix curve of order less than or equal to  $\frac{n}{2}$ .

Consider now a surface  $F_2^n$  with a minimum directrix  $\gamma^m$  of order  $m$ . If a prime  $[n]$  contains more than  $m$  generators of  $F_2^n$  it will meet  $\gamma^m$  in more than  $m$  points and so contain it entirely, and its intersection with the ruled surface will consist of  $\gamma^m$  and  $n - m$  generators. This further illustrates the fact that there are no curves, other than  $\gamma^m$  itself, on  $F_2^n$  of order less than  $n - m$ . If, however, a prime  $[n]$  is made to contain exactly  $m$  generators of  $F_2^n$  (as it always can since  $m < \frac{n+1}{2}$ ) it will



not contain  $\gamma^m$  and will meet  $F_2^n$  further in a curve of order  $n - m$ ; unless, however, every prime through the  $m$  generators necessarily contains a further generator. But this cannot be; for we can take a prime through  $m$  arbitrary generators and  $n - 2m + 1$  arbitrary points of  $F_2^n$ ; if such a prime necessarily contained a further generator it would also contain  $\gamma^m$  and therefore the  $n - 2m + 1$  generators of  $F_2^n$  through the chosen points; its intersection with  $F_2^n$  would then be a curve of order at least

$$m + m + n - 2m + 1$$

or  $n + 1$ , which is impossible.

We have then clearly established the existence of curves of order  $n - m$  on  $F_2^n$ ; so that all rational normal ruled surfaces of order  $n$  in  $[n + 1]$  can be obtained as in § 41.

45. Since any directrix curve of order  $n - m$  and any  $m$  generators lie together in an  $[n]$  all the curves of order  $n - m$  can be obtained by means of primes through  $m$  generators arbitrarily chosen. We can now state the following:

*On the surface  $F_2^n$  whose minimum directrix is of order  $m$  there are  $\infty^{n-2m+1}$  curves of order  $n - m$ . These curves are all obtained by primes through any  $m$  fixed generators of the surface; and through any  $n - 2m + 1$  points of general position on the surface there passes one such curve.*

Any two curves of order  $n - m$  will have  $n - 2m$  intersections; a prime containing one of them meets  $F_2^n$  further in  $m$  generators and meets the other curve in  $n - m$  points of which  $m$  are on these generators, the remaining  $n - 2m$  being intersections of the two curves.

In particular *the ruled surface of even order which has minimum directrices of order  $\frac{n}{2}$  has  $\infty^1$  of them; through any point of the surface there passes one such curve and no two of them can intersect. The curves can all be obtained by means of primes through any  $\frac{n}{2}$  generators.*

In a similar way it can be shewn that, if  $0 < k < m$ , there are, on a surface  $F_2^n$  with a minimum directrix  $\gamma^m$  of order  $m$ ,  $\infty^{n-2k+1}$  directrix curves of order  $n - k$  such that through any  $n - 2k + 1$  points of general position on the surface there passes just one. All these curves can be obtained by means of primes through any  $k$  fixed generators. Two directrices of orders  $n - k$  and  $n - k'$  intersect in  $n - k - k'$  points.

46. A space  $[n - 1]$  of general position will meet  $F_2^n$  in  $n$  points; we can, however, consider spaces containing generators of  $F_2^n$ . If an  $[n - 1]$  contains  $k$  generators it will contain  $k$  points of  $\gamma^m$ ; so that if  $k > m$  it will contain the whole of  $\gamma^m$ , and its intersection with  $F_2^n$  will consist of  $\gamma^m$  and a certain number of generators. If, however,  $k < m$ ,

$[n - 1]$  will not contain  $\gamma^m$  and its intersection with  $F_2^n$  will consist of the  $k$  generators and a certain number of isolated points. A prime through  $[n - 1]$  meets  $F_2^n$  in the  $k$  generators and a curve of order  $n - k$ ; this curve will meet  $[n - 1]$  in  $n - k$  points of which  $k$  lie on the generators, the remaining  $n - 2k$  giving the isolated intersections of  $F_2^n$  with  $[n - 1]$ . Hence *an  $[n - 1]$  through  $k$  generators of  $F_2^n$ , where  $k < m$ , meets  $F_2^n$ , in general, in these generators and in  $n - 2k$  points.*

47. Having then such a knowledge of the properties of a normal surface in  $[n + 1]$  we can deduce the properties of a rational ruled surface of order  $n$  in  $[3]$  by projecting from a space  $[n - 3]$ . The spaces  $[n - 2]$  joining  $[n - 3]$  to the points of the normal surface  $F$  meet the  $[3]$  in the points of the projected surface  $f$ , while the spaces  $[n - 1]$  joining  $[n - 3]$  to the generators of  $F$  meet the  $[3]$  in the generators of  $f$ . Surfaces  $f$  derived by projection from the same type of normal surface are projectively equivalent; but surfaces  $f$  derived from different types of normal surfaces are not. Knowing a method of generating  $F$  by a  $(1, 1)$  correspondence between two directrix curves we can deduce a method of generating  $f$ , and so if we wish obtain the equation to  $f$ .

In order to give a first illustration of this work we shall obtain the two cubic ruled surfaces of  $[3]$ , already met with in §§ 37, 38, by projection of the normal cubic ruled surface in  $[4]^*$ .

48. In  $[4]$  there is only one type of cubic ruled surface  $F$ ; it has a directrix line  $\lambda$  and  $\infty^2$  directrix conics. Through any two points of  $F$  (not on the same generator and neither of them on  $\lambda$ ) there passes one directrix conic; while any two directrix conics have a single intersection. Project  $F$  from a point  $O$  of  $S_4$  on to a solid  $\Sigma$ ; we get a cubic ruled surface  $f$  in  $\Sigma$ . The lines joining  $O$  to the points of  $F$  meet  $\Sigma$  in the points of  $f$ , while the planes joining  $O$  to the generators of  $F$  meet  $\Sigma$  in the generators of  $f$ .

Through a general point of  $S_4$  there passes one plane containing a directrix conic of  $F$ . There cannot be more than one, because the intersection of two such planes is the intersection of the two directrix conics which they contain, and this lies on  $F$ . But there certainly exists one such plane; for any solid through the point meets  $F$  in a twisted cubic, one of whose chords passes through the point, and the two points in which the chord meets  $F$  determine a directrix conic whose plane contains the chord.

If then  $O$  is a general point of  $S_4$  there is a plane  $\pi$  passing through it which contains a directrix conic  $\Gamma$ ; and  $\pi$  meets  $\Sigma$  in a line  $R$  which is a double directrix of  $f$ , two generators of  $f$  passing through every point of  $R$ . The plane joining  $O$  to  $\lambda$ , the line directrix of  $F$ , meets  $\Sigma$  in a line  $R'$  which

\* Cf. Veronese, *loc. cit.* 229-232.

is a second directrix of  $f$ ; through any point of  $R'$  there passes one generator of  $f$  (see Fig. 1).

Any plane of  $\Sigma$  passing through  $R'$  is joined to  $O$  by a solid containing  $\lambda$  and therefore meeting  $F$  in  $\lambda$  and two generators  $g_1$  and  $g_2$ \*. This solid meets  $\pi$  in a line through  $O$ , and this line must meet  $\Gamma$  in the two points  $g_1\Gamma$  and  $g_2\Gamma$ . Then  $g_1$  and  $g_2$  project into two generators of  $f$  which intersect in a point of  $R$  and lie in the plane through  $R'$  from which we started. Hence any plane through  $R'$  contains two generators of  $f$  which meet in a point of  $R$ .




Fig. 1.

This shews that  $f$  is a cubic ruled surface in  $\Sigma$  belonging to the first of our two species. Since  $F$  can be generated by placing its directrix line in (1, 1) correspondence with any of its directrix conics\* we deduce at once that  $f$  can be generated by placing two lines  $R, R'$  in (1, 2) correspondence.

49. This general type of surface has naturally been obtained by selecting a general point of projection. Let us then take a point  $O$  in a plane through  $\lambda$  which contains a generator  $g$  of  $F$ , and project from  $O$  on to a solid  $\Sigma$ . Then the plane  $O\lambda g$  meets  $\Sigma$  in a line  $R$  which is a directrix and also a generator of  $f$ .

It is clear from the projection that any plane through  $R$  contains one other generator of  $f$ , while through any point of  $R$  there passes one other

\* § 44.

generator of  $f$ . Thus we have our second type of cubic ruled surface in three-dimensional space.

From the known generation of  $F$  we deduce that  $f$  can be generated by a line and conic with a point of intersection placed in (1, 1) correspondence without a united point\*. If we regard  $F$  as generated by a (1, 1) correspondence between its directrix line  $\lambda$  and one of its directrix conics  $\Gamma$  this point of intersection is the projection of the point  $g\Gamma$ . The point of intersection is not a united point because the two points in which  $g$  is met by  $\lambda$  and  $\Gamma$  project into different points of  $R$  †.

50. *Irrational ruled surfaces.* It is proved by Segre ‡ that all ruled surfaces of order  $n$  which are elliptic ( $p = 1$ ) can be obtained by projection from normal surfaces in  $[n - 1]$ ; he has made a complete study of these surfaces, and we shall give some account of his results and apply them to surfaces of the fifth and sixth orders. But when we come to surfaces for which  $p = 2$  the normal space is no longer unique; this is clearly exemplified in our study of sextic surfaces. It can be shewn that all ruled surfaces of order  $n$  and genus  $p$  which are not contained in a space of dimension less than  $n - p + 1$  are cones§; and on the other hand that all ruled surfaces of order  $n$  and genus  $p$  have a normal space|| of dimension at least  $n - 2p + 1$ . Thus it would seem that, in order to obtain all the ruled surfaces of order  $n$  and genus  $p$  by projection, we have to consider the possible normal surfaces in  $p$  different spaces.

51. *The chords and tangents of a ruled surface.* If we have a surface in higher space then, just as in [3], all the tangent lines at a non-singular point of the surface lie in a plane¶. The tangent plane at any point of a ruled surface must clearly contain the generator of the ruled surface which passes through the point; and it is the fact that the tangent planes at the different points of the generator form a pencil of planes related to the range of points on the generator and all lie in a [3]¶. We shall speak of this [3] as the *tangent solid* of the ruled surface along the generator.

We thus have a four-dimensional locus  $M_4$  formed by these  $\infty^1$  tangent solids of the ruled surface; we can also regard  $M_4$  as consisting of the  $\infty^2$  tangent planes or of the  $\infty^3$  tangent lines of the ruled surface. If the ruled surface lies in [4] there will be a finite number of its tangent solids (or planes or lines) passing through any point of [4]. If the ruled surface lies

\* Cf. § 20.

† Concerning this surface we may also refer to Reye, *Die Geometrie der Lage*, 3 (Leipzig, 1923), 156.

‡ "Ricerche sulle rigate ellittiche di qualunque ordine," *Atti Torino*, 21 (1886), 868.

§ Segre, *ibid.* § 2.

|| Segre, *Math. Ann.* 34 (1889), 4.

¶ Del Pezzo, *Palermo Rendiconti*, 1 (1887), 243-245.

in  $[r]$ , where  $r > 4$ , there will be a finite number of its tangents meeting an  $[r - 4]$  of general position, and this number is the order of  $M_4$ .

The chords of the ruled surface form a five-dimensional locus  $M_5$  on which the locus  $M_4$  formed by the tangents lies. If the surface is in  $[4]$  there will be  $\infty^1$  chords of it through a general point of  $[4]$ . If the surface is in  $[5]$  there will be a finite number of its chords passing through a general point of  $[5]$ ; this number is the number of "apparent double points" of the surface. If the surface is in  $[r]$ , where  $r > 5$ , there will be a finite number of its chords meeting an  $[r - 5]$  of general position, and this number is the order of  $M_5$ .

For example, on the rational ruled surface  $F_2^n$  in  $[n + 1]$

$$\left\| \begin{array}{cccccc} x_0 & x_1 & \dots & x_{m-1} & x_{m+1} & \dots & x_n \\ x_1 & x_2 & \dots & x_m & x_{m+2} & \dots & x_{n+1} \end{array} \right\| = 0,$$

the coordinates of any two points can be taken as

$$(a^m, a^{m-1}, \dots, 1, a^{n-m}b, \dots, ab, b),$$

and

$$(a^m, a^{m-1}, \dots, 1, a^{n-m}\beta, \dots, a\beta, \beta).$$

Then the coordinates of any point on the line joining these are seen to satisfy

$$\left\| \begin{array}{cccccc} x_0 & x_1 & \dots & x_{m-2} & x_{m+1} & \dots & x_{n-1} \\ x_1 & x_2 & \dots & x_{m-1} & x_{m+2} & \dots & x_n \\ x_2 & x_3 & \dots & x_m & x_{m+3} & \dots & x_{n+1} \end{array} \right\| = 0,$$

which represents a locus of five dimensions and of order  $\frac{1}{2}(n - 2)(n - 3)$ .

52. In the actual projection of a ruled surface from higher space these two loci  $M_4$  and  $M_5$  are useful for studying any double curve which there may be on the projected surface. When the surface is in  $[r]$  we project on to  $[3]$  from an  $[r - 4]$ ; the locus  $M_5$  meets  $[r - 4]$  in a curve  $\mathfrak{D}$ ; through each point of this curve there passes a chord of  $F$  and the  $[r - 3]$  joining  $[r - 4]$  to such a chord meets the  $[3] \Sigma$  on to which we are projecting in a double point of the projected surface  $f$ . We thus have the double curve of  $f$  in  $(1, 1)$  correspondence with the curve in which  $M_5$  meets  $[r - 4]$ . The chords of  $F$  meeting  $[r - 4]$  give on  $F$  a curve  $C$  in  $(2, 1)$  correspondence with the double curve of  $f$ ; the number of branch-points of this correspondence on  $C$  is the number of tangents of  $C$  which meet  $[r - 4]$ , and this is simply the order of  $M_4$ . Thus the genus of the double curve of  $f$  is the same as that of the curve  $\mathfrak{D}$  in which  $M_5$  meets  $[r - 4]$ ; it can also be calculated by Zeuthen's formula when we know the genus of  $C$ .

There will, of course, not be a curve in  $[r - 4]$  unless  $r > 5$ ; if  $r = 5$  we have a finite number of chords of  $F$  passing through each point of a line, while if  $r = 4$  we have an infinity of chords of  $F$  passing through a single point: in both these cases, however, we still have the curve  $C$  on  $F$ .

The projection of the cubic ruled surface in [4] exemplifies this; the chords of  $F$  passing through an arbitrary point  $O$  lie in a plane and meet  $F$  in the points of a directrix conic  $\Gamma$ . There are two tangents of  $\Gamma$  passing through  $O$ .

53. We already know, by means of the representation of  $f$  as a curve on  $\Omega$ , that  $f$  will have, in general, a finite number of triple points: this is confirmed by the projection. The triple points of  $f$  must arise from spaces  $[r - 3]$  passing through the centre of projection  $[r - 4]$  and meeting  $F$  each in three points. Now the spaces  $[r - 3]$  in  $[r]$  which contain  $[r - 4]$  are in aggregate  $\infty^3$ , and one condition is necessary for an  $[r - 3]$  to meet a surface in  $[r]$ . Hence there are, containing  $[r - 4]$ ,  $\infty^2$  spaces  $[r - 3]$  each meeting  $F$  in one point,  $\infty^1$  spaces  $[r - 3]$  each meeting  $F$  in two points and a finite number of spaces  $[r - 3]$  each meeting  $F$  in three points. The first set simply joins  $[r - 4]$  to the  $\infty^2$  points of  $F$ ; the second set contains the  $\infty^1$  chords of  $F$  which meet  $[r - 4]$ ; the third gives the triple points of  $f$ . The singularities of a general ruled surface in [3] are a double curve and a finite number of triple points, these points being also triple points of the double curve.

For a ruled surface in [4] the only singularities, in general, are a finite number of double points. If the surface is a projection of a non-singular ruled surface in [5] this is clear at once, for we have already seen that there is a finite number of chords of this surface passing through a general point of [5]. If the surface is the projection of a surface in  $[r]$ , when  $r > 5$ , the centre of projection is an  $[r - 5]$ , and this, if it is of general position, meets the  $M_5$  formed by the chords of the surface in a finite number of points.

54. *Conclusion.* We have now two powerful methods of obtaining the different kinds of ruled surfaces in [3]; the first by considering the surfaces as curves on  $\Omega$ , the second by projection of normal surfaces from higher space. For surfaces which are the most general of their order and genus we must choose the curve to have a general position on  $\Omega$ , and the centre of projection to have a general position in regard to the normal surface. To obtain the other surfaces we specialise the position of the curve on  $\Omega$ , and the position of the centre of projection in regard to the normal surfaces.

The first method is only applicable to the ruled surfaces in [3], whereas the second is equally applicable to ruled surfaces in any space; but the duality of the surface is lost in the second method, whereas it is retained in the first; it is only in [3] that the line is the self-dual element. The two methods will have to lead to the same results; and we may confidently expect a greater efficiency than usual when we have two such different methods confirming the workings of one another.

We proceed then to the main task of classifying ruled surfaces in [3], beginning with those of the fourth order.

## CHAPTER II

### QUARTIC RULED SURFACES

#### INTRODUCTORY

55. The object of this chapter is to give a detailed and exhaustive classification of the quartic ruled surfaces of three-dimensional space. The most general type of surface is mentioned by Chasles\*; the surfaces were studied and classified, though not quite exhaustively, by Cayley†, who obtained his different types by means of directing curves and gave algebraic equations for them. The complete classification was first given by Cremona‡, who generated his surfaces by means of correspondences between two curves.

We shall illustrate the general methods of this volume by obtaining the quartic ruled surfaces of [3] in two ways:

(a) by regarding their generators as represented by the points of a quartic curve on a quadric  $\Omega$  in [5] (§ 31);

(b) by regarding them as projections of normal quartic ruled surfaces in higher space.

56. Lines in [3] are  $\infty^4$  in aggregate; there are thus  $\infty^1$  lines satisfying three conditions. It is one condition for a line to meet a curve; so that the lines which meet each of three given curves  $C_1, C_2, C_3$  are  $\infty^1$  in aggregate and form a ruled surface. Salmon remarked§ that if the curves  $C_1, C_2, C_3$  are of respective orders  $m_1, m_2, m_3$  then the order of the ruled surface is  $2m_1m_2m_3$ ; further, the three curves are multiple curves on the surface; through every point of  $C_1$  there pass  $m_2m_3$  generators, through every point of  $C_2$   $m_3m_1$  generators and through every point of  $C_3$   $m_1m_2$  generators. Cayley added the further statement that if  $C_2$  and  $C_3$  have  $\alpha$  intersections,  $C_3$  and  $C_1$  have  $\beta$  intersections and  $C_1$  and  $C_2$  have  $\gamma$  intersections, then the order of the ruled surface is reduced to  $2m_1m_2m_3 - m_1\alpha - m_2\beta - m_3\gamma$ , while through every point of  $C_1$  there pass  $m_2m_3 - \alpha$  generators, through

\* *Comptes Rendus*, 52 (1861), 1094.

† *Papers*, 5 (1864), 214–219, and 6 (1868), 312–328.

‡ *Memorie dell' Accademia di Bologna* (2), 8 (1868), 235; *Opere*, 2, 420. Concerning quartic ruled surfaces we may refer also to Reye, *Die Geometrie der Lage*, 2 (Stuttgart, 1907), 301, and Sturm, *Liniengeometrie*, 1 (Leipzig, 1892), 48.

§ *Cambridge and Dublin Mathematical Journal*, 8 (1853), 45; cf. *Geometry of Three Dimensions*, 2 (Dublin, 1915), 90.

every point of  $C_2$   $m_3 m_1 - \beta$  generators and through every point of  $C_3$   $m_1 m_2 - \gamma$  generators\*.

Cayley then chose sets of directing curves which would give ruled surfaces of the fourth order. For example, one such set can be taken to consist of two conics  $C_1$  and  $C_2$  with two common points and a line  $C_3$  meeting one of the conics, say  $C_1$ . Then

$$\begin{aligned} m_1 &= 2, & m_2 &= 2, & m_3 &= 1, \\ \alpha &= 1, & \beta &= 0, & \gamma &= 2, \end{aligned}$$

and we have a ruled surface of order 4 on which  $C_2$  is a double conic and  $C_3$  a double line. He did not succeed however in discovering all the quartic ruled surfaces in this way †.

Incidentally we also have a ruled surface formed by the chords of one curve which meet another given curve; or by the lines trisecant to a given curve. Formulae can be given for the orders of such ruled surfaces; we shall see below that a quartic ruled surface is formed by the chords of a twisted cubic which meet a given line.

57. Cremona on the other hand generated the quartic ruled surfaces by means of a (1, 1) correspondence between two conics. In general the conics will not degenerate and will not intersect; but further types of surfaces are obtained if one or both of the conics degenerates into a line counted doubly, and still further types can arise if the two curves intersect.

It can at once be shewn by elementary methods that the ruled surface is of the fourth order. For let the two conics be  $C$  and  $C'$  and suppose that an arbitrary line meets their planes in  $p$  and  $q$ . Then an arbitrary plane through the line gives a pair of points  $x, y$  on  $C$  and a pair of points  $z', u'$  on  $C'$ . Then as the plane varies in the pencil the pairs of points  $x, y$  will describe an involution on  $C$ , the join of every pair of points passing through  $p$ ; while the pairs of points  $z', u'$  describe an involution on  $C'$ , the join of every pair of points passing through  $q$ ; the pencils of lines through  $p$  and  $q$  in the planes of the two conics are thus homographically related. But to the points  $x, y$  of  $C$  there will correspond, in the (1, 1) correspondence between the conics, points  $x', y'$  of  $C'$ ; and as the points  $x, y$  describe the involution on  $C$  the points  $x', y'$  will describe an involution on  $C'$ . The join of  $x'$  and  $y'$  thus passes through a fixed point  $p'$ , and the pencil of lines through  $p'$  in the plane of  $C'$  is thus homographic with the pencil

\* *Papers*, 5 (1864), 203.

† There are, as we shall see, ten species of quartic ruled surfaces with rational plane sections and two with elliptic plane sections; in Cayley's first paper we find the two elliptic ones and six of the rational ones. This paper was seen by Cremona before he published his own, and meanwhile Cayley discovered two other rational quartic ruled surfaces. He did not know of the existence of the remaining two species until he was informed of them by Cremona.



of lines through  $p$  in the plane of  $C$ . We have then in the plane of  $C'$  two homographic pencils of lines with vertices  $p'$  and  $q$ ; the locus of intersections of corresponding lines is a conic. This conic meets  $C'$  in four points, and the plane joining any one of these four points to  $pq$  contains a pair of corresponding points of  $C$  and  $C'$  and therefore a generator of the ruled surface. We have thus four generators of the surface meeting  $pq$ , so that the ruled surface is of the fourth order.

It is clear that the planes of  $C$  and  $C'$  are bitangent planes of the ruled surface, each meeting it in a conic and two generators. The plane of  $C$ , for example, meets  $C'$  in two points and therefore contains the two generators which join those points to the corresponding points of  $C$ .

Further, there are three double points of the ruled surface in an arbitrary plane. In the plane of  $C$  we have the intersection of the two generators lying therein and two other points, namely, those intersections of the generators with  $C$  which do not correspond to the two intersections of the plane with  $C'$ . The double curve of the ruled surface is thus a twisted cubic\*.

The (1, 1) correspondence between  $C$  and  $C'$  determines a projectivity between their planes †; there are certain lines in either plane which intersect their corresponding lines in the other plane, and the planes of such pairs of intersecting lines are known to form a developable of the third class ‡. Such a plane necessarily contains two generators of the ruled surface; and conversely every bitangent plane of the ruled surface will meet the planes of  $C$  and  $C'$  in lines which correspond to one another in the projectivity. Thus we see that the quartic ruled surface has a bitangent developable of the third class.

Just as the points of the surface which lie in a plane containing two generators, but not themselves on either of the generators, lie on a conic, so the tangent planes of the surface which pass through a point of intersection of two generators, but do not themselves contain either of these generators, touch a quadric cone.

After these preliminary remarks we proceed to the classification of the quartic ruled surfaces by the two methods; the reader is referred for further investigations to Cremona's paper.

\* The double curve, being algebraic, meets every plane of the space containing the ruled surface in the same number of points. It is then sufficient, in order to be able to say that the double curve is a twisted cubic, to prove that it meets the plane of  $C$  in three points.

† von Staudt, *Geometrie der Lage*, 2 (Nürnberg, 1857), 149; Reye, *Geometrie der Lage*, 2 (Stuttgart, 1907), 10.

‡ von Staudt, *Geometrie der Lage*, 3 (Nürnberg, 1860), 326; Reye, *Geometrie der Lage*, 2 (Stuttgart, 1907), 163.

## SECTION I

RATIONAL QUARTIC RULED SURFACES CON-  
SIDERED AS CURVES ON  $\Omega$ 

58. The generators of a rational quartic ruled surface are represented on  $\Omega$  by the points of a rational quartic curve  $C$ . Such a curve necessarily lies in a space  $S_4$  and may be contained in a space  $S_3$ , so that we consider five possibilities as follows:

I.  $C$  is a rational normal quartic lying in an  $S_4$  which has no special relation in regard to  $\Omega$ .

II.  $C$  lies in a tangent prime  $T$  of  $\Omega$ .

III.  $C$  lies in a tangent prime  $T$  of  $\Omega$  and passes through  $O$ , the point of contact of  $\Omega$  and  $T$ .

IV.  $C$  lies in an  $S_3$ , the intersection of tangent primes to  $\Omega$  at two points  $O$  and  $O'$ .

V.  $C$  lies in an  $S_3$  through which there passes only one tangent prime of  $\Omega$ ,  $S_3$  meeting  $\Omega$  in a quadric cone with vertex  $V$ .

59. Let us now examine the general case I.

To find the degree of the double curve of the ruled surface we take a plane  $\rho$  on  $\Omega$  and find how many planes  $\omega$  there are which meet this plane  $\rho$  in a line and also meet  $C$  in two points\*. The chord joining the points of  $C$  in such a plane  $\omega$  would have to meet  $\rho$  (on its line of intersection with  $\omega$ ). But  $\rho$  meets  $S_4$  in a line; and we know that any line in  $S_4$  is met by three chords of  $C$ †, because when we project  $C$  from the line on to a plane we obtain a rational quartic, i.e. a plane quartic with three double points. A chord of  $C$  which meets the line lies entirely on  $\Omega$  as meeting it in three points, and therefore a plane  $\omega$  passes through it. We therefore obtain three planes  $\omega$  such as we require.

Hence the double curve is of the third order.

Similarly the bitangent developable is of the third class.

The tangent prime of  $\Omega$  at any point  $P$  of  $C$  meets  $S_4$  in a solid which contains the tangent of  $C$  at  $P$  and meets  $C$  in two points other than  $P$ . Hence every generator of the surface is met by two others. On any generator there lie two points of the double curve, while through any generator there pass two planes of the bitangent developable.

Since the points of  $C$  lie in an  $S_4$  which does not touch  $\Omega$  the generators of the surface belong to a linear complex which is not special. The surface was thus given by Cayley, as that formed by the chords of a twisted cubic belonging to a linear complex ‡.

\* § 33.

† Cf. § 8.

‡ *Papers*, 6, 316.

The chords of a twisted cubic are  $\infty^2$  in aggregate; they are therefore represented on  $\Omega$  by the points of a surface  $V$ . Since one chord of the cubic can be drawn through any point of the space [3] in which it lies, the surface  $V$  meets every plane  $\omega$  on  $\Omega$  in one point; and since there are three chords of the cubic in any plane of the [3], the surface  $V$  meets every plane  $\rho$  on  $\Omega$  in three points. Hence  $V$  is of order 4, meeting an arbitrary solid in four points. This surface  $V$  is in fact the surface known as Veronese's surface\*.

The statement that the chords of a twisted cubic belonging to a linear complex form a rational quartic ruled surface is the same as the statement that the section of Veronese's surface by a prime is a rational quartic curve.

**60. Quartic ruled surfaces of type II.** A tangent prime  $T$  of  $\Omega$  meets  $\Omega$  in a quadric point-cone †; this contains the two systems of planes on  $\Omega$  through the point of contact  $O$ . Any solid of  $T$  meets  $\Omega$  in a quadric surface; and the lines joining  $O$  to every point of this quadric surface lie on  $\Omega$ . The rational quartic  $C$  is projected from  $O$  into a rational quartic on this quadric. Remembering that the generators of the quadric lie on the planes of the two systems through  $O$  we can at once subdivide type II into three parts:

II (A).  $C$  meets every plane  $\omega$  through  $O$  in three points and every plane  $\rho$  through  $O$  in one point.

II (B).  $C$  meets every plane  $\omega$  through  $O$  in one point and every plane  $\rho$  through  $O$  in three points.

II (C). A chord of  $C$  passes through  $O$ ;  $C$  meets every plane of  $\Omega$  through  $O$  in two points.

The point  $O$  of  $\Omega$  represents a line  $R$ ; this is a directrix of the quartic surface, being met by every generator.

When the surface is of the type II (A) there are three generators passing through every point of  $R$ , while there is one generator lying in each plane through  $R$ .

To find the double curve we take, as before, an arbitrary plane  $\rho$ , and we consider those chords of  $C$  which meet this plane. These are the three chords of  $C$  which meet the line of intersection of the plane  $\rho$  with the tangent prime  $T$ , and are therefore those chords of  $C$  which lie in the plane  $\omega$  joining this line to  $O$ .

If we now interpret this result in the space  $S_3$  containing the ruled surface the plane  $\rho$  represents an arbitrary plane of  $S_3$ ; and the three planes  $\omega$  which represent the three intersections of this plane of  $S_3$  with the double

\* For this surface, regarded as representing the chords of a twisted cubic, see Baker, *Principles of Geometry*, 4, 52-55, where references to the literature concerning it will also be found.

† Cf. § 30.

curve all coincide in a single plane through  $O$ . Hence the three points of intersection of the plane of  $S_3$  and the double curve all coincide in the intersection of this plane with the line  $R$ .

Hence the double curve is the line  $R$  counted three times.

To find the bitangent developable we take an arbitrary plane  $\omega$ ; this meets  $T$  in a line which is met by three chords of  $C$ , and the bitangent developable of the surface is a general developable of the third class.

When the surface is of the type II (B) there is one generator passing through each point of  $R$  and three generators lying in any plane through  $R$ . The double curve is a non-degenerate twisted cubic, while the bitangent developable consists of the pencil of planes through  $R$  counted three times.

When the surface is of the type II (C) there are two generators which pass through any point of  $R$  and two which lie in any plane through  $R$ .

An arbitrary plane  $\rho$  meets  $T$  in a line; one of the three chords of  $C$  which meet the line is contained in the plane  $\omega$  joining it to  $O$ . Hence the double curve consists of  $R$  and a conic.  $R$  and the conic intersect; their point of intersection is represented by the  $\omega$ -plane which contains the chord of  $C$  passing through  $O$ . Similarly, the bitangent developable consists of the pencil of planes through  $R$  together with the tangent planes of a quadric cone, one tangent plane of the cone passing through  $R$ . This tangent plane of the cone is represented on  $\Omega$  by the  $\rho$ -plane which contains the chord of  $C$  passing through  $O$ .

**61. Quartic ruled surfaces of type III.** We can subdivide III into two parts; since the projection of  $C$  from  $O$  on to a solid in  $T$  is now a twisted cubic we have:

III (A).  $C$  meets every plane  $\omega$  through  $O$  in two points and every plane  $\rho$  through  $O$  in one point other than  $O$ .

III (B).  $C$  meets every plane  $\omega$  through  $O$  in one point and every plane  $\rho$  through  $O$  in two points other than  $O$ .

The point  $O$  represents a line  $R$  which is a directrix and also a generator of the ruled surface.

In the type III (A) there are two generators other than  $R$  passing through each point of  $R$  and one generator other than  $R$  lying in each plane through  $R$ . An arbitrary plane  $\rho$  meets  $T$  in a line; but the three chords of  $C$  meeting this line all lie in the plane  $\omega$  joining it to  $O$ . Hence the double curve is the line  $R$  counted three times. An arbitrary plane  $\omega$  meets  $T$  in a line, and the plane  $\rho$  joining this line to  $O$  contains one chord of  $C$ . There will be two other chords of  $C$  meeting the line, so that the bitangent developable consists of the pencil of planes through  $R$  together with the tangent planes of a quadric cone.

In the type III (B) there are two generators other than  $R$  lying in any plane through  $R$  and one generator other than  $R$  passing through any point of  $R$ . The double curve consists of  $R$  and a conic, the bitangent developable of the pencil of planes through  $R$  counted three times.

**62. Quartic ruled surfaces of type IV.** In IV we have a rational quartic  $C$  lying on the quadric surface in which  $\Omega$  is met by  $S_3$ . We have then two possibilities:

IV (A).  $C$  meets all generators of one system in three points and all of the other system in one point.

IV (B).  $C$  has a double point and meets every generator in two points.

Through a general point of  $S_3$  we can draw three chords to either of these rational quartics; in the second case we include the line to the double point.

The points  $O$  and  $O'$  of  $\Omega$  represent lines  $R$  and  $R'$ , which are both met by every generator; the surface has two directrices.

The planes of the two systems on  $\Omega$  which pass through  $O$  and  $O'$  meet  $S_3$  in the two systems of generators of the quadric surface. Each generator is the intersection of a plane of one system through  $O$  with a plane of the opposite system through  $O'$ .

In IV (A) we may suppose that the generators trisecant to  $C$  lie in the  $\omega$ -planes through  $O$  and in the  $\rho$ -planes through  $O'$ . Then through any point of  $R$  there pass three generators lying in a plane through  $R'$ ; any plane through  $R'$  contains three generators meeting in a point of  $R$ . The double curve consists of the points of the line  $R$  counted three times; the bitangent developable consists of the planes through the line  $R'$  counted three times.

In IV (B) the double point of  $C$  represents a double generator  $G$  of the surface. Through any point of  $R$  there pass two generators lying in a plane through  $R'$  and through any point of  $R'$  there pass two generators lying in a plane with  $R$ . The double curve therefore consists of the points of  $R$ ,  $R'$ ,  $G$ , while the bitangent developable consists of the three pencils of planes through  $R$ ,  $R'$ ,  $G$ .

**63. Quartic ruled surfaces of type V.** In V  $C$  lies on a quadric cone  $Q$  with vertex  $V$  and must therefore have a double point. If two quadrics in  $S_3$  touch there are three cones through their curve of intersection, one of which has its vertex at the double point; so that V can be subdivided according as

V (A), the double point of  $C$  is not at  $V$ ,

or V (B), the double point of  $C$  is  $V$ .

Through every generator of  $Q$  there passes a plane of  $\Omega$  of either system.

In V (A) the surface has a directrix line  $R$ ; through any point of  $R$  there pass two generators which lie in a plane with  $R$ , while any plane through

$R$  contains two generators meeting in a point of  $R$ .  $R$  is represented on  $\Omega$  by the point  $V$ .

An arbitrary plane of  $\Omega$  meets  $S_3$  in a point  $P$  on  $Q$ , and it is clear that the projection of  $C$  from  $P$  on to a plane of  $S_3$  is a quartic curve with a double point and a tacnode, the tangent at the tacnode being the intersection of the plane with the tangent plane of the cone along the generator through  $P$ . Hence the three chords of  $C$  which pass through  $P$  consist of the generator  $PV$  counted twice and the line to the double point.

Hence the double curve consists of the directrix  $R$  counted twice and a double generator, while the bitangent developable consists of the planes through  $R$  counted twice and the planes through the double generator.

In  $V(B)$  the surface has a directrix line  $R$  which is also a double generator.  $C$  meets every generator of  $Q$  in one point other than  $V$ ; so that through any point of  $R$  there passes one generator other than  $R$ , while every plane through  $R$  contains one generator other than  $R$ .

An arbitrary plane of  $\Omega$  meets  $S_3$  in a point  $P$  on  $Q$ , and the projection of  $C$  from  $P$  on to a plane of  $S_3$  has a triple point. Hence the three chords of  $C$  passing through  $P$  all coincide with  $PV$ . The double curve is therefore the line  $R$  counted three times, while the bitangent developable consists of the planes through  $R$  counted three times.

64. This completes the determination of the rational quartic ruled surfaces of ordinary space. We have obtained in all ten species, and these are the same as the ten species obtained by Cremona. A table of the surfaces is given on p. 303.

65. *The representation of the double curve and bitangent developable.* Two generators of the surface which intersect are represented on  $\Omega$  by two points of  $C$  such that the chord joining them lies entirely upon  $\Omega$ . This chord will meet an arbitrary prime in a point which can be taken as representative either of the point of the double curve in which the two generators intersect or of the plane of the bitangent developable in which the two generators lie\*. Hence the chords of  $C$  which lie on  $\Omega$  trace out in the prime a curve  $D$  which is in (1, 1) correspondence with both the double curve and the bitangent developable.

We assume that  $C$  is general unless the contrary is stated.

The chords of  $C$  form a locus  $U_3$  of three dimensions; any point of this locus which is on  $\Omega$  and not on  $C$  must be on a chord of  $C$  that lies entirely on  $\Omega$ . Hence these chords form a ruled surface in the  $S_4$  containing  $C$ , this ruled surface being the intersection of  $\Omega$  and  $U_3$ . The section of this ruled surface by a prime or, what is the same thing, by an  $S_3$  lying

\* Cf. § 33.

in  $S_4$ , gives us the curve  $D$ . The  $S_3$  will meet  $\Omega$  in a quadric surface and  $U_3$  in another surface; and  $D$  is the intersection of these two surfaces.

If  $C$  is projected from a line in  $S_4$  on to a plane we have a quartic curve with three double points, so that the line must be met by three chords of  $C$ . Hence  $U_3$  is of the third order and may be denoted by  $U_3^3$ . Moreover, if  $C$  is projected from a line which meets it on to a plane we have a plane cubic with one double point, so that the line is only met by one chord of  $C$  besides those passing through its intersection with  $C$ . Hence  $C$  is a double curve on  $U_3^3$ . Since no plane can meet  $C$  in four points no two chords of  $C$  can intersect except on  $C$  itself, so that there is no double surface on  $U_3^3$ . The section of  $U_3^3$  by an  $S_3$  is a four-nodal cubic surface.

The curve  $D$  in  $S_3$  is therefore the intersection of a four-nodal cubic surface with a quadric passing through the nodes; this is a sextic curve with four double points. It lies on the quadric and meets every generator in three points; so that if it is projected from a point of the quadric on to a plane we obtain a sextic with two triple points and four double points. This is a rational curve.

**66.** If the curve  $D$  should happen to break up into one or more parts we expect the double curve and the bitangent developable to break up into the same number of parts.

Let us consider in particular the case II (C).  $C$  lies in a prime  $T$  touching  $\Omega$  in a point  $O$ ; a chord of  $C$  passes through  $O$ , while every plane of  $\Omega$  through  $O$  meets  $C$  in two points.

Every chord of  $C$  which lies entirely upon  $\Omega$  is such that the plane joining it to  $O$  also lies entirely upon  $\Omega$ . Hence we can clearly separate these chords into two distinct classes, those joined to  $O$  by planes  $\omega$  and those joined to  $O$  by planes  $\rho$ . We therefore expect  $D$  to break up, and moreover to break up into two similar parts.

Consider those points of an arbitrary  $S_3$  of  $T$  lying on chords which are joined to  $O$  by  $\omega$ -planes; they form a curve  $D_1$  lying on the quadric  $Q$  in which  $S_3$  meets  $\Omega$ . One system of generators of  $Q$  is joined to  $O$  by  $\omega$ -planes, and clearly there is one point of  $D_1$  on each of these generators. A generator of  $Q$  of the opposite system is joined to  $O$  by a  $\rho$  plane: this plane contains one chord of  $C$ , so that the generator of  $Q$  is met by two other chords; these must be joined to  $O$  by  $\omega$ -planes. Hence on this generator we have two points of  $D_1$ .

Hence  $D_1$  is a twisted cubic meeting the generators of  $Q$  in  $\omega$ -planes in one point and the generators of  $Q$  in  $\rho$ -planes in two points.

Similarly, those points of  $S_3$  lying on chords that are joined to  $O$  by  $\rho$ -planes form a twisted cubic  $D_2$  which meets the generators of  $Q$  in  $\omega$ -planes in two points and the generators of  $Q$  in  $\rho$ -planes in one point.

Through any point of  $C$  there pass two chords which lie entirely on  $\Omega$ , and these will be joined to  $O$  by planes of opposite systems. This holds in particular for the four points of intersection with  $S_3$ .

Thus we have again the intersection of a quadric with a four-nodal cubic surface; but here the intersection  $D$  breaks up into two twisted cubics,  $D_1$  and  $D_2$ , each passing through the four nodes.  $D_1$  and  $D_2$  have one other common point lying on the chord of  $C$  which passes through  $O$ .

The pairs of generators of the surface which meet in the points of  $R$  and whose planes touch a quadric cone are represented on  $\Omega$  by the pairs of points of  $C$  which lie in the  $\varpi$ -planes through  $O$ . The chords of  $C$  joining these points form a cubic ruled surface  $\phi_1^*$ , so that the points of  $R$  and the tangent planes of the cone are in (1, 1) correspondence with the generators of  $\phi_1$ . The section of  $\phi_1$  by  $S_3$  is the twisted cubic  $D_1$ .  $\phi_1$  has a directrix line  $l_1$  and the plane  $\varpi_0$  passing through  $l_1$  represents the vertex of the quadric cone.

Similarly, the pairs of generators of the surface which lie in the planes through  $R$  and whose points of intersection lie on a conic are represented on  $\Omega$  by the pairs of points of  $C$  which lie in the  $\rho$ -planes through  $O$ . The chords of  $C$  joining these points form a cubic ruled surface  $\phi_2$ , so that the planes through  $R$  and the points of the conic are in (1, 1) correspondence with the generators of  $\phi_2$ . The section of  $\phi_2$  by  $S_3$  is the twisted cubic  $D_2$ .  $\phi_2$  has a directrix line  $l_2$  and the plane  $\rho_0$  passing through  $l_2$  represents the plane of the conic.

The planes  $\varpi_0$  and  $\rho_0$  do not lie in  $T$ .

67. We can also consider in this way the curves  $C$  of type III. Suppose for definiteness that  $C$  is of the type III (B). The ruled surface formed by the chords of  $C$  which lie on  $\Omega$  breaks up into two parts; the cubic cone projecting  $C$  from  $O$  and the cubic ruled surface formed by those chords of  $C$  which lie in the  $\rho$ -planes through  $O$  but do not themselves pass through  $O$ . The tangent of  $C$  at  $O$  belongs both to the cone and to the ruled surface, and the plane  $\varpi$  through this tangent represents a point of intersection of  $R$  and the double conic. The plane of the double conic is represented on  $\Omega$  by the  $\rho$ -plane which contains the directrix of the cubic ruled surface.

\* These chords join the pairs of an involution on  $C$ . The joins of the pairs of points of an involution on a rational curve of order  $n$  form a ruled surface of order  $n - 1$  (see footnote to § 19).



## SECTION II

RATIONAL QUARTIC RULED SURFACES CONSIDERED  
AS PROJECTIONS OF NORMAL SURFACES IN  
HIGHER SPACE

68. The rational quartic ruled surfaces  $f$  of ordinary space can all be obtained as projections of the rational normal quartic ruled surfaces  $F$  of  $S_5$ ; the projection will be from a line  $l$  not meeting  $F$ . There are two kinds of surfaces  $F$ ; the general surface with  $\infty^1$  directrix conics and the surface with a directrix line.

We first confine our attention to the general surface  $F$ , and find that by suitably choosing  $l$  we can obtain six types of surfaces  $f$ . From the known methods of generating  $F$  we deduce methods of generating these six types of surfaces; all these methods are given in Cremona's paper.

The other four types are obtained by projection from the surface with a directrix line.

69. *The normal surface in [5] with directrix conics.* The general surface  $F$  has  $\infty^1$  directrix conics; any such conic is determined by one point of  $F$  and no two conics intersect.

We first assume that  $l$  has a general position, and project from  $l$  on to a solid  $\Sigma$ .

A prime through  $l$  meets  $F$  in a rational normal quartic curve, three of whose chords will meet  $l$ . Hence in any prime through  $l$  there are three planes that pass through  $l$  and meet two generators of  $F$ . Projecting: on any plane section of  $f$  there are three points through which two generators pass. Thus the double curve of  $f$  is a twisted cubic.

The planes of the  $\infty^1$  conics on  $F$  meet any prime in lines forming a ruled surface  $A$ . The prime meets  $F$  in a rational normal quartic, and the generators of  $A$  must be chords of this curve since each one meets  $F$  in the two points where it cuts the corresponding directrix conic. Further, these chords join pairs of an involution on the curve because each conic is determined by one point of  $F$ . Thus they form a cubic ruled surface. This shews that the planes of the directrix conics form a locus  $V_3^3$ .

The line  $l$ , having a general position, does not meet the  $V_3^3$ ; but a plane  $\omega$  drawn through  $l$  meets the planes of three directrix conics  $\Gamma_1, \Gamma_2, \Gamma_3$ .  $\omega$  meets  $\Sigma$  in a point  $P$ . Then through  $\omega$  and  $\Gamma_1$  there passes a prime which meets  $F$  in  $\Gamma_1$  and two generators; the projections of these two generators will lie in a plane through  $P$ . There are two other bitangent planes through  $P$  arising from the primes  $\omega\Gamma_2$  and  $\omega\Gamma_3$ . Hence there are three bitangent planes of  $f$  passing through a general point  $P$  of  $\Sigma$ , so that the bitangent developable of  $f$  is of the third class.

$f$  is therefore of the type I.

$F$  can be generated by a (1, 1) correspondence between any two of its directrix conics. Hence the most general quartic ruled surface of ordinary space can be generated by a (1, 1) correspondence between two conics.

70.  $F$  contains  $\infty^3$  directrix cubics, any such curve being determined by three points of  $F$ . All these can be obtained as prime sections of  $F$  residual to any given generator. Any one of these curves lies in a solid; it is easily seen that for general positions of  $l$  none of these solids contains  $l$ .

Suppose, however, that  $l$  is taken in one of these solids  $\sigma$ . Then the projected surface  $f$  has a directrix line  $R$ , the intersection of  $\Sigma$  and  $\sigma$ . A plane through  $l$  lying in  $\sigma$  meets  $F$  in the three points where it meets the directrix cubic of  $\sigma$ ; hence through any point of  $R$  there pass three generators of  $f$ . A prime containing  $\sigma$  meets  $F$  again in one generator; hence every plane through  $R$  contains one generator of  $f$ .

Thus  $f$  is of the type II (A).

$F$  can be generated by placing a directrix cubic and a directrix conic in (1, 1) correspondence with a united point. Hence the surface  $f$  can be generated by taking a line and a conic with a common point  $P$  and placing them in (1, 3) correspondence with  $P$  as a united point. To  $P$  regarded as a point of the conic corresponds  $P$  regarded as a point of the line; while to  $P$  regarded as a point of the line correspond three points of the conic of which  $P$  is one.

71. Now let  $l$  be taken so as to meet a plane  $\gamma$  of  $V_3^3$ ,  $\gamma$  containing a directrix conic  $\Gamma$ . The solid  $l\gamma$  meets  $\Sigma$  in a line  $R$  which is a directrix of  $f$ . A plane through  $l$  lying in the solid  $l\gamma$  meets  $\gamma$  in a line having two points of intersection with  $\Gamma$ , and therefore meets two generators of  $F$ . Hence through any point of  $R$  there pass two generators of  $f$ . Moreover, a prime through  $l\gamma$  meets  $F$  in  $\Gamma$  and two generators, so that any plane through  $R$  contains two generators of  $f$ .

Thus  $f$  is of the type II (C).

$f$  can be generated by placing a line and a conic in (1, 2) correspondence.

With this last choice of  $l$  the solid  $l\gamma$  will not in general contain a generator of  $F$ , but we may clearly choose  $l$  so that it does. Then  $f$  has a directrix line  $R$  which is also a generator. Any plane through  $R$  contains one other generator, while through any point of  $R$  there pass two generators other than  $R$  itself.

Here  $f$  is of the type III (A).

To determine a generation for  $f$  suppose that  $F$  is generated by means of  $\Gamma$  and some other directrix conic  $\Gamma'$ ; the solid  $l\gamma$  containing a generator  $g$  which meets  $\Gamma$  in  $X$  and  $\Gamma'$  in  $X'$  (Fig. 2).

The solid  $l\Gamma$  meets  $\Sigma$  in the directrix  $R$ , while the planes joining  $l$  to the points of  $\Gamma'$  meet  $\Sigma$  in the points of a conic  $C$ .

Now consider the plane  $lX'$ . Since it lies in the solid  $lg$  or  $l\gamma$  it meets the plane  $\gamma$  in a line, and therefore meets  $\Sigma$  in a point on  $R$ . Hence  $R$  and  $C$  have a common point  $P$ . The point  $X'$  on  $\Gamma'$  gives the corresponding point  $X$  on  $\Gamma$ , which is clearly projected into a point  $Q$  of  $R$  different from  $P$ .

Hence the (1, 1) correspondence between  $\Gamma$  and  $\Gamma'$  gives a (1, 2) correspondence between  $R$  and  $C$ ,  $P$  not being a self-corresponding point.

Thus  $f$  may be generated by taking a line and a conic with a common point and placing them in (1, 2) correspondence without a united point.




Fig. 2.

72. Further, we may choose  $l$  to meet two planes  $\gamma$  and  $\gamma'$  of  $V_3^3$ , these planes containing directrix conics  $\Gamma$  and  $\Gamma'$ . Then  $f$  has two directrices  $R$  and  $R'$ , the intersections of the solids  $l\gamma$  and  $l\gamma'$  with  $\Sigma$ . Let  $l$  meet  $\gamma$  and  $\gamma'$  in  $P$  and  $P'$  respectively.

The lines of  $\gamma$  through  $P$  give an involution on  $\Gamma$ . This gives an involution of pairs of generators of  $F$ , and thus an involution on  $\Gamma'$  also. But the lines of  $\gamma'$  through  $P'$  give a second involution on  $\Gamma'$  which will have a pair of points in common with the former. This pair of points gives a pair of generators  $g$  and  $g'$  on  $F$ ; and clearly  $g, g', l$  lie in a solid as having two common transversals. Thus the projected surface  $f$  has a double generator.

A prime through  $l$  and  $\gamma$  meets  $F$  in  $\Gamma$  and two generators which meet  $\Gamma'$  in a pair of points collinear with  $P'$ . Thus a plane through  $R$  contains two generators of  $f$  which intersect in a point of  $R'$ . Similarly a plane through  $R'$  meets  $f$  in a pair of generators which intersect in a point of  $R$ .

Hence  $f$  is of the type IV (B).

We may clearly generate  $f$  by a (2, 2) correspondence between two lines  $R$  and  $R'$ ; but this will not be the most general (2, 2) correspondence, it must be specialised so as to give the double generator.

Conversely, take two generators  $g$  and  $g'$  of  $F$ ; the solid  $gg'$  does not meet  $F$  again\*. Take  $l$  in the solid  $gg'$ .

Any point  $p$  of  $g$  determines a conic of  $F$  meeting  $g'$  in a point  $p'$ ; the plane of this conic meets the solid  $gg'$  in the line  $pp'$ . Also, through  $p$  there passes a transversal of  $g, g', l$  meeting  $g'$  in a point  $q'$ .

The ranges ( $p'$ ) and ( $q'$ ) on  $g'$ , both being homographic with the range ( $p$ ) on  $g$ , are homographic with each other; in general, they will have two self-corresponding points. Thus  $l$  is met by two planes of  $V_3^3$  as before.




Fig. 3.

But it may happen that the two self-corresponding points coincide, so that  $l$  meets only one plane of  $V_3^3$ . The surface  $f$  has now a double generator and a directrix  $R$ , but  $R$  must be regarded as a coincidence of two directrices. Any plane through  $R$  contains two generators, while through any point of  $R$  there pass two generators,  $f$  being of the type V (A).

The  $V_3^3$  formed by the planes of the directrix conics meets an arbitrary solid in a cubic curve passing through the four points of intersection of the solid with  $F$ . But a solid containing a pair of generators  $g$  and  $g'$  of  $F$  is met by the planes of the directrix conics in lines which give homographic ranges on  $g$  and  $g'$  and therefore form a regulus.

A line  $l$  in the solid meets the regulus in two points, and on projecting we have a surface  $f$  of the type IV (B); but if  $l$  is taken to touch the quadric surface on which the regulus lies we have a surface  $f$  of the type V (A).

73. We can give another generation for a surface of type IV (B) which can immediately be specialised to give the type V (A).

\* § 46.

We have generated  $f$  by placing its two directrices  $R, R'$  in (2, 2) correspondence with united elements  $X, X'$  corresponding to one another; whence the double generator  $XX'$ .

Let  $C$  be the projection of  $\Gamma_0$ , a third directrix conic of  $F$  (Fig. 3).

The plane of  $C$  is the intersection of  $\Sigma$  with the prime containing  $l$  and  $\Gamma_0$ . This prime meets  $\gamma$  in a line through  $P$  and  $\gamma'$  in a line through  $P'$ ; hence the two generators which it contains must be the pair  $g$  and  $g'$  that determine a solid containing  $l$ . Thus the plane of  $C$  contains the line  $XX'$ .

We may therefore generate  $f$  by a (1, 2) correspondence between  $R$  and  $C$ ; to the point  $X$  in which  $R$  meets the plane of  $C$  there must correspond two points of  $C$  collinear with  $X$ , these giving the double generator  $XX'$ .

To the points of  $R$  correspond pairs of points of  $C$  which form an involution; thus the chords of  $C$  which join these pairs of points all pass through a fixed point. This point will lie on  $XX'$  and is in fact  $X'$ . For the two generators of  $f$  which issue from any point of  $R$  both meet  $R'$ , so that their plane contains  $R'$  and meets the plane of  $C$  in a line through  $X'$ .

Similarly the points of  $R'$  give rise to pairs of points on  $C$  whose joins all pass through  $X$ .

This at once suggests the following generation for a surface of the type V (A): take a line  $R$  and a conic  $C$  in (1, 2) correspondence, every point of  $R$  giving a corresponding pair of points of  $C$  whose join passes through the point in which  $R$  meets the plane of  $C$ .

74. *The normal surface in [5] with a directrix line.* We now consider the projections of the surface  $F$  which has a directrix line  $\lambda$ .

Taking first a general position of  $l$ , the projected surface  $f$  has a directrix line  $R$ , the intersection of  $\Sigma$  with the solid  $l\lambda$ . A prime through  $l\lambda$  meets  $F$  in  $\lambda$  and three generators, so that any plane through  $R$  contains three generators of  $f$ . A plane through  $l$  lying in the solid  $l\lambda$  meets  $\lambda$  in one point, so that through any point of  $R$  there passes one generator of  $f$ .

Hence  $f$  is of the type II (B).

$F$  can be generated by a (1, 1) correspondence between  $\lambda$  and a directrix cubic. Hence  $f$  can be generated by a (1, 1) correspondence between a line  $R$  and a twisted cubic.

The planes through  $\lambda$  which contain the generators of  $F$  form a locus  $U$  of three dimensions, which does not meet a line of general position.

But if  $l$  is chosen to meet a plane  $\pi$  of  $U$  the solid  $l\pi$  meets  $\Sigma$  in a line  $R$  which is a directrix and also a generator of  $f$ . The primes through the solid  $l\lambda$  now meet  $F$  in sets of three generators of which one is always the generator in  $\pi$ . Thus any plane through  $R$  contains two generators of  $f$  other than  $R$ , while through any point of  $R$  there passes one generator of  $f$  other than  $R$ .

Hence  $f$  is of the type III (B).

By reasoning similar to that employed in obtaining the generation for the type III (A) we see that  $f$  can be generated by a line and twisted cubic with a point of intersection, placed in (1, 1) correspondence without a united point.

Further,  $l$  may meet two planes of  $U$ . Then  $f$  has a directrix  $R$  which is also a double generator. Any plane through  $R$  contains one other generator, while through any point of  $R$  there passes one other generator.

Here  $f$  is of the type V (B).

$f$  can be generated by taking a twisted cubic and one of its chords, and placing them in (1, 1) correspondence without any united point.

It is easily seen that the locus  $U$  is of the third order, meeting an arbitrary solid in the points of a cubic curve. If  $l$  met three planes of  $U$  it would lie entirely on  $U$  and so meet  $F$ , so that we do not consider this possibility.

75.  $F$  contains  $\infty^3$  directrix cubics, they can all be obtained as prime sections residual to any given generator. In general  $l$  does not lie in a solid with any of these curves; if, however, it happens that  $l$  and one of the cubics do lie in the same solid  $\sigma$  the projected surface  $f$  will have two directrices,  $R$ , the intersection of  $\Sigma$  with the solid  $l\lambda$ , and  $R'$ , the intersection of  $\Sigma$  with the solid  $\sigma$ .

A prime through  $l\lambda$  meets  $F$  in  $\lambda$  and three generators and meets  $\sigma$  in a plane through  $l$ . Hence any plane through  $R$  contains three generators meeting in a point of  $R'$ . Through any point of  $R$  there passes one generator.

Thus  $f$  is of the type IV (A).

$f$  is generated by placing two lines  $R$  and  $R'$  in (3, 1) correspondence. This last generation is mentioned by Cremona.

We have now completed the determination of the rational quartic ruled surfaces of ordinary space; the results obtained by the two methods are in complete agreement with one another and also with Cremona's results.

### SECTION III

#### ELLIPTIC QUARTIC RULED SURFACES

76. We must now give a short account of those surfaces which are elliptic; i.e. those whose plane sections are elliptic quartic curves. The generators of such a surface will be represented on  $\Omega$  by the points of an elliptic quartic curve  $C$ , which necessarily lies in a solid  $S_3$ .

77. In general,  $S_3$  will be the intersection of the tangent primes to  $\Omega$  at two points  $O$  and  $O'$ ; these two points represent lines  $R$  and  $R'$  which are directrices of the surface.  $S_3$  meets  $\Omega$  in a quadric  $Q$ ; every generator

of  $Q$  is met by  $C$  in two points and is the intersection of a plane of  $\Omega$  through  $O$  with a plane of the opposite system through  $O'$ .

Hence through any point of  $R$  there pass two generators which lie in a plane through  $R'$ , while through any point of  $R'$  there pass two generators which lie in a plane through  $R$ .

The double curve consists of the lines  $R$  and  $R'$ , while the bitangent developable consists of the pencils of planes through  $R$  and  $R'$ . A plane section of the surface is a quartic with double points on  $R$  and  $R'$ .

This is the surface which is generated by means of the most general (2, 2) correspondence between two lines  $R$  and  $R'$ . It is Cayley's first species and Cremona's eleventh.

This general type of elliptic quartic ruled surface is the most general type of Segre's "rigate biquadratische\*."

An elliptic quartic curve is determined by eight points on a quadric†; hence a ruled quartic surface can be made to contain eight lines of a linear congruence.



78. Suppose now that only one tangent prime of  $\Omega$  passes through  $S_3$ ,  $S_3$  meeting  $\Omega$  in a quadric cone with vertex  $V$ .  $C$  lies on this cone and does not pass through  $V$ . Each generator of the cone meets  $C$  in two points and is the intersection of two planes of  $\Omega$ , one of each system.

The point  $V$  represents a line  $R$  which is a directrix of the surface.

Any plane of  $\Omega$  meets  $S_3$  in a point  $P$  of the cone. The projection of  $C$  from  $P$  on to a plane of  $S_3$  is clearly a quartic with a tacnode; so that the two chords of  $C$  which can be drawn through  $P$  consist of the generator  $PV$  counted twice.

The double curve is thus the line  $R$  counted twice, while the bitangent developable consists of the planes through  $R$  counted twice. Through any point of  $R$  there pass two generators which lie in a plane with  $R$ . A plane section of the surface is a quartic with a tacnode on  $R$ .

This surface is Cayley's fourth species and Cremona's twelfth.

79. An elliptic quartic ruled surface cannot contain a simple directrix or a conic; but if we take a plane through a generator we obtain a cubic curve on the surface. We have in this way  $\infty^2$  cubic curves on the surface. Through any two general points on the surface there pass two of these curves, because the line joining the two points meets the surface in two other points, and through either of the two generators passing through the latter points there is a plane containing the former points. In general two of the cubic curves will intersect in two points on the line of intersection of their planes; the other two points of intersection of this line

\* *Memorie Torino*, 36 (1885), 142.

† Salmon, *Geometry of Three Dimensions*, 1 (Dublin, 1914), 360.

with the surface being on the generators which lie in the planes of the cubic curves.

To generate the surface with two directrices we take a line  $R$  and a plane cubic curve without a double point, the curve passing through the point  $P$  in which  $R$  meets its plane. We then place the line and cubic in (1, 2) correspondence with  $P$  for a united point. The pairs of points of the cubic which correspond to the points of  $R$  must be collinear with a fixed point  $P'$  of the curve\*; the range of points on  $R$  is related projectively to the pencil of lines through  $P'$ , the point  $P$  of the range corresponding to the ray  $P'P$  of the pencil. The planes joining the points of the range to the corresponding rays of the pencil all pass through a line  $R'$  which passes through  $P'$  †.

From this we can deduce at once the generation for an elliptic quartic ruled surface with one directrix. We take a line  $R$  and an elliptic cubic curve with a point of intersection  $P$  and place them in (1, 2) correspondence with  $P$  as a united point. But here the pairs of points of the curve which correspond to the points of the line are collinear with  $P$  itself; and the two generators which pass through any point of  $R$  are co-planar with  $R$ . The correspondence is at once determined by a projectivity between the points of the range on  $R$  and the lines of the pencil through  $P$ , the point  $P$  of the range corresponding to the tangent of the cubic curve at  $P$ . This latter is the generator of the surface which lies in the plane of the cubic curve.

#### SECTION IV

### ALGEBRAICAL RESULTS CONNECTED WITH QUARTIC RULED SURFACES

80. We first obtain the equations of the different kinds of quartic ruled surfaces by the methods which we have given for generating them. The results may be compared with those in Salmon's *Geometry of Three Dimensions* ‡.

The surface of the type I is generated by the chords of a twisted cubic which belong to a linear complex.

We can take the coordinates of any point on the cubic to be  $(\theta^3, \theta^2, \theta, 1)$ ; the six coordinates of the line joining the two points for which the parameter has the values  $\lambda$  and  $\mu$  are then

$$\lambda^2 + \lambda\mu + \mu^2, \quad \lambda + \mu, \quad 1, \quad -\lambda\mu, \quad \lambda\mu(\lambda + \mu), \quad -\lambda^2\mu^2.$$

\* The joins of the pairs of points of a  $g_2^1$  on a plane cubic without a double point all meet the curve again in the same point.

† If there were not a united element the planes would touch a quadric cone.

‡ 2 (Dublin, 1915), §§ 546–554.



Suppose now that the chord belongs to the linear complex

$$bl + 2fm + cn - (b + 2g)l' + 2hm' - am' = 0.$$

Then

$$a\lambda^2\mu^2 + b(\lambda + \mu)^2 + c + 2f(\lambda + \mu) + 2g\lambda\mu + 2h\lambda\mu(\lambda + \mu) = 0.$$

Now the coordinates of any point on the chord are

$$\lambda^3 + k\mu^3, \quad \lambda^2 + k\mu^2, \quad \lambda + k\mu, \quad 1 + k,$$

for which

$xz - y^2 = k\lambda\mu(\lambda - \mu)^2$ ,  $xt - yz = k(\lambda + \mu)(\lambda - \mu)^2$ ,  $yt - z^2 = k(\lambda - \mu)^2$ ,  
so that the locus of the chords is the quartic ruled surface

$$a(xz - y^2)^2 + b(xt - yz)^2 + c(yt - z^2)^2 + 2f(xt - yz)(yt - z^2) + 2g(yt - z^2)(xz - y^2) + 2h(xz - y^2)(xt - yz) = 0,$$

on which the existence of the twisted cubic as a double curve is clear. This then may be taken to represent the general surface\* of type I.

If, however, we have

$$b(b + 2g) - 4fh + ac = 0$$

the surface degenerates into one formed by the chords of the cubic which meet the line whose coordinates are

$$\{-(b + 2g), \quad 2h, \quad -a, \quad b, \quad 2f, \quad c\}$$

and is of the type II (B).

Let us now take the surface of the type II (A).

We take a line and a conic in (1, 3) correspondence with a united point †. The sets of three points of the conic which correspond to the points of the line form a singly infinite set of triangles whose sides all touch another conic. The two conics have four common tangents; these touching the first conic in double points of four of the sets of three points. Take two of these four points as  $X$  and  $Y$ ; the plane of the conic being  $z = t$  and the line being  $x = y = 0$ . Let  $Z$  be the point of the line which gives rise to the set of three points with the double point  $X$  and  $T$  the point of the line which gives rise to the set of three points with the double point  $Y$ . Let the equation of the conic be

$$yz + zx + xy = z - t = 0.$$

Then any point of the conic can be written  $(\theta, 1 - \theta, \theta^2 - \theta, \theta^2 - \theta)$ , while any point of the line is  $(0, 0, \phi, 1)$ ; and the general (1, 3) relation

$$\phi(a\theta^3 + b\theta^2 + c\theta + d) = A\theta^3 + B\theta^2 + C\theta + D$$

must become

$$\phi(\theta - 1)^2(a\theta + \beta) = \theta^2(a\theta + \gamma).$$

Any point of the surface is

$$(\theta, \quad 1 - \theta, \quad \theta^2 - \theta + \lambda\phi, \quad \theta^2 - \theta + \lambda),$$

so that

$$z - \phi t = x(\theta - 1)(1 - \phi),$$

and

$$\phi = \frac{x^2\{ax + \gamma(x + y)\}}{y^2\{ax + \beta(x + y)\}}.$$

Thus

$$(x + y)(z - \phi t) + xy(1 - \phi) = 0,$$

$$(yz + zx + xy)y^2\{ax + \beta(x + y)\} = \{xy + t(x + y)\}x^2\{ax + \gamma(x + y)\},$$

which divides by  $x + y$ , giving

$$zy^2\{ax + \beta(x + y)\} - tx^2\{ax + \gamma(x + y)\} = xy\{ax(x - y) + \gamma x^2 - \beta y^2\},$$

$$z\{\beta y^3 + (\alpha + \beta)xy^2\} - t\{\gamma x^2y + (\gamma + \alpha)x^3\} = xy\{(\gamma + \alpha)x^2 - \alpha xy - \beta y^2\}.$$

\* Salmon, § 549.

† § 70.

Writing then  $z = \zeta - x$  and  $t = \tau - y$  we find\*

$$\zeta (ay^3 + bxy^2) + \tau (cx^2y + dx^3) = ex^2y^2.$$

For a surface of the type II (C) we must take a line and a conic in (1, 2) correspondence †. Let the line meet the plane of the conic in  $Y$ , and let  $X$  and  $Z$  be the points of contact of the tangents from  $Y$ . Then the conic is

$$xz - y^2 = t = 0.$$

Take  $T$  on the line.

Then any point of the conic is  $(\theta^2, \theta, 1, 0)$ , while any point of the line is  $(0, \phi, 0, 1)$ . The points of the line give rise to pairs of an involution on the conic; one of these pairs will be on a line through  $Y$ , and we can take  $T$  to correspond to this pair. Since this pair divides  $X$  and  $Z$  harmonically (the join passing through the pole of  $XZ$ ) we can take their parameters to be  $+1$  and  $-1$ , so that the (1, 2) relation will be of the form

$$\phi (a\theta^2 + b\theta + c) = \theta^2 - 1.$$

Any point of the surface is  $(\lambda\theta^2, \lambda\theta + \mu\phi, \lambda, \mu)$ , giving

$$x = \theta^2z, \quad y = \theta z + \phi t = \theta z + \frac{\theta^2 - 1}{a\theta^2 + b\theta + c} t,$$

$$(y - \theta z) (a\theta^2 + b\theta + c) = (\theta^2 - 1) t,$$

$$ax (y - \theta z) + byz\theta - bzx + cz (y - \theta z) = t (x - z);$$

or

$$axy - bzx + cyz + t (z - x) = \theta (azx - byz + cz^2).$$

Squaring this we have the equation to the surface in the form

$$[cyz - bzx + axy + zt - tx]^2 = zx (ax - by + cz)^2.$$

This clearly has the double line  $z = x = 0$  and also a double conic, the intersection of the plane

$$ax - by + cz = 0$$

with the quadric

$$cyz - bzx + axy + zt - tx = 0,$$

or with the quadric

$$b (y^2 - zx) + t (z - x) = 0.$$

The planes  $x = 0$  and  $cy + t = 0$  give a torsal generator, as also do the planes  $z = 0$  and  $ay - t = 0$ , the respective tangent planes being  $x = 0$  and  $z = 0$ .

The intersection of the surface with the plane  $t = 0$  is given by

$$\begin{aligned} 0 &= (cyz - bzx + axy)^2 - zx (ax - by + cz)^2 \\ &= \{y (ax - by + cz)^2 - b (zx - y^2)\}^2 - zx (ax - by + cz)^2 \\ &= (zx - y^2) [b^2 (2x - y^2) - 2by (ax - by + cz) - (ax - by + cz)^2] \\ &= (zx - y^2) [(b^2 - 2ac)zx - a^2x^2 - c^2z^2], \end{aligned}$$

which consists of the original conic together with two lines through  $Y$ .

The equation of this pair of lines can be written

$$(ax + by + cz) (ax - by + cz) + b^2 (y^2 - zx) = 0,$$

so that they pass through the two points of intersection of the double conic and the conic  $xz - y^2 = t = 0$ ; the two conics lying on a quadric.

For a surface of the type III (A) we take a line and conic with a point of intersection and place them in (1, 2) correspondence without a united point †.

\* Cf. Salmon, § 546.

† § 71.

The pairs of corresponding points on the conic form an involution; take the two double points of this as  $X$  and  $Y$ , and let the corresponding points be  $Z$  and  $T$ . Then we can take the line to be  $x = y = 0$  and the conic to be

$$yz + zx + xy = z - t = 0.$$

Any point of the line can be written  $(0, 0, \phi, 1)$  and any point of the conic  $(\theta, 1 - \theta, \theta^2 - \theta, \theta^2 - \theta)$ . The  $(1, 2)$  relation must be of the form

$$\phi (\theta - 1)^2 = \theta^2 \alpha.$$

Then the coordinates of any point of the surface are

$$(\theta, 1 - \theta, \theta^2 - \theta + \lambda\phi, \theta^2 - \theta + \lambda)$$

for which

$$z - \phi t = x (\theta - 1) (1 - \phi),$$

so that

$$z (\theta - 1)^2 - \alpha \theta^2 t = x (\theta - 1) [(\theta - 1)^2 - \alpha \theta^2],$$

or

$$(x + y) (zy^2 - \alpha tx^2) = xy (\alpha x^2 - y^2).$$

Writing  $-t = y + \tau$  and  $z = \zeta - x$ , we obtain

$$(y^2 \zeta + \alpha \tau x^2) (x + y) = mx^2 y^2,$$

and we take for the equation of the surface\*

$$mx^2 y^2 = (x + y) (x^2 t + y^2 z).$$

For a surface of the type III (B) we may take a line and a twisted cubic with a point of intersection and place them in  $(1, 1)$  correspondence without a united point†. We will, however, obtain this surface as the reciprocal of III (A).

The equation of the surface III (A) is

$$\mu x^2 y^2 = (x + y) (x^2 t + y^2 z).$$

A tangent plane is

$$lx + my + nz + pt = 0,$$

where

$$n = (x + y) y^2, \quad p = (x + y) x^2,$$

and

$$lx + my = - (nz + pt) = - \mu x^2 y^2.$$

Also

$$nx^2 + py^2 = 2x^2 y^2 (x + y),$$

so that

$$2 (lx + my) (x + y) + \mu (nx^2 + py^2) = 0,$$

or, since this is homogeneous in  $x$  and  $y$  and  $x^2 : y^2 = p : n$ ,

$$lp + mn + (l + m) \sqrt{np} + \mu np = 0,$$

or

$$(lp + mn + \mu np)^2 = (l + m)^2 np.$$

This being the tangential equation of the surface III (A), the point-equation of the surface III (B) can be written †

$$(xt + yz + \mu zt)^2 = zt (x + y)^2.$$

It has a double conic  $x + y = xt + yz + \mu zt = 0$  and a double line  $z = t = 0$  which meets the conic in  $(1, -1, 0, 0)$ .

To generate a surface of the type IV (A) we take two lines  $R$  and  $R'$  in  $(3, 1)$  correspondence‖. There will be four points of  $R'$  for which two of the three corresponding points on  $R$  coincide. Take two of these four points as  $X$  and  $Y$ , the corresponding coincident elements being  $Z$  and  $T$ . Any point of  $R'$  is  $(\theta, 1, 0, 0)$ , while any point of  $R$  is  $(0, 0, \phi, 1)$ ; the  $(1, 3)$  relation between  $\theta$  and  $\phi$  is necessarily of the form

$$\theta = \frac{\alpha \phi^3 + \beta \phi^2}{\gamma \phi + \delta}.$$

\* Cf. Salmon, § 548.

† § 74.

‡ Cf. Salmon, § 548.

‖ § 75.

Then any point of the surface has coordinates  $(\theta, 1, \lambda\phi, \lambda)$ , or

$$[\alpha\phi^3 + \beta\phi^2, \gamma\phi + \delta, \lambda\phi(\gamma\phi + \delta), \lambda(\gamma\phi + \delta)].$$

Hence

$$\frac{x}{y} = \frac{\alpha z^3/t^3 + \beta z^2/t^2}{\gamma z/t + \delta} = \frac{(\alpha z + \beta t) z^2}{(\gamma z + \delta t) t^2}$$

and the equation of the surface is\*

$$x(\gamma z + \delta t) t^2 = y(\alpha z + \beta t) z^2.$$

To generate the surface of the type IV (B) we take two lines  $R, R'$  in (2, 2) correspondence, the correspondence being specialised so as to give a double generator †. In a general (2, 2) correspondence there will be four points of either line giving rise to pairs of coincident points on the other; but here two of the four points will themselves coincide. Take then  $X$  and  $Y$  as the corresponding double elements; and let  $Z$  be another point on  $R$  which gives rise to a pair of coincident points  $T$  on  $R'$ . Any point of  $R$  is  $(\phi, 0, 1, 0)$  and any point of  $R'$  is  $(0, \theta, 0, 1)$ , the (2, 2) relation will be of the form

$$\phi = \alpha\phi^2 + \beta\theta\phi + \gamma\theta^2.$$

Any point of the surface is  $(\lambda\phi, \mu\theta, \lambda, t)$ , whence

$$\frac{x}{z} = \phi, \quad \frac{y}{t} = \theta,$$

and

$$\frac{x}{z} = \alpha \frac{x^2}{z^2} + \beta \frac{xy}{zt} + \gamma \frac{y^2}{t^2},$$

or

$$xzt^2 = \alpha x^2 t^2 + \beta xyzt + \gamma y^2 z^2;$$

and we have for the equation of the surface ‡

$$y^2 z^2 + mxyzt + t^2(\alpha xz + \beta x^2) = 0.$$

There is a second generation which can be used for a surface of the type IV (B)||; we take a line and conic in (1, 2) correspondence, making the two points of the conic which correspond to the point in which the line meets the plane collinear with this point. Take the point of intersection of the line with the plane of the conic as  $Y$ ; the points  $X$  and  $Z$  being the points of contact of the tangents from  $Y$ . Then the conic is  $t = xz - y^2 = 0$ , with points  $(\theta^2, \theta, 1, 0)$ , and we can take the parameters of the two points which correspond to  $Y$  as  $\pm 1$ . Any point of the line is  $(0, \phi, 0, 1)$ , if we take  $T$  on the line. The pairs of points of the conic which correspond to the points of the line form an involution; we shall take  $T$  as the point which gives rise to the pair of the involution including  $Z$ . Then the (1, 2) relation between  $\phi$  and  $\theta$  must be of the form

$$\phi(\theta^2 - 1) = A\theta^2 + B\theta.$$

Any point of the surface is  $(\lambda\theta^2, \lambda\theta + \mu\phi, \lambda, \mu)$ , giving

$$x = \theta^2 z, \quad y = \theta z + \phi t = \theta z + t \frac{A\theta^2 + B\theta}{\theta^2 - 1},$$

\* Cf. Salmon, § 547.

† § 72.

‡ Cf. Salmon, § 553.

|| § 73

i.e. 
$$(y - \theta z) (\theta^2 - 1) = (A\theta^2 + B\theta) t,$$

$$x (y - \theta z) - z (y - \theta z) = t (Ax + Bz\theta),$$

$$xy - yz - Atx = \theta (Bzt + zx - z^2);$$

and on squaring we have the equation of the surface in the form

$$(yz - xy + Atx)^2 = zx (x - z + Bt)^2.$$

This surface has clearly the double lines  $x = z = 0$  and  $x - z = t = 0$ . Writing it in the form

$$\{y (z - x + Bt) + t (Ax - By)\}^2 = zx (x - z + Bt)^2,$$

we see that it has also a double line

$$x - z + Bt = Ax - By = 0.$$

The line  $x - z = t = 0$  is a double generator; it meets the other two lines which are double directrices.

Now if  $B = 0$  the two directrices coincide, and we have a surface of the type V (A) whose equation is

$$(yz - xy + Atx)^2 = xz (x - z)^2.$$

The pair of points of the conic which corresponds to any point of the line is now such that its join passes through  $Y$ .

The section by any plane  $ax + \beta y + \gamma z + At = 0$  is a quartic curve with a tacnode at  $x = z = 0$  and another ordinary node.

To generate a surface of the type V (B) we take a twisted cubic and one of its chords and place them in (1, 1) correspondence without any united points\*. Take  $X$  and  $Y$  to be the points where the line meets the cubic; any point of the line is then  $(\theta, 1, 0, 0)$ . Also,  $\theta$  may be taken as the parameter of the cubic, and if  $Z$  and  $T$  correspond to  $X$  and  $Y$  respectively any point of the curve can be taken to have coordinates

$$\theta (\theta - \beta), \quad \theta (\theta - \alpha), \quad \theta (\theta - \alpha) (\theta - \beta), \quad (\theta - \alpha) (\theta - \beta).$$

Any point of the ruled surface will have coordinates

$$\theta (\theta - \beta) + \lambda\theta, \quad \theta (\theta - \alpha) + \lambda, \quad \theta (\theta - \alpha) (\theta - \beta), \quad (\theta - \alpha) (\theta - \beta).$$

Then 
$$z - \alpha t = (\theta - \alpha) t, \quad z - \beta t = (\theta - \beta) t,$$

$$yz - xt = \theta t \{ \theta^2 - (\alpha + 1) \theta + \beta \};$$

so that 
$$(yz - xt) (z - \alpha t) (z - \beta t) = zt \{ z^2 - (\alpha + 1) zt + \beta t^2 \}$$

is the equation of the surface.

By taking two other planes of the pencil  $z + kt = 0$  instead of  $z = 0$  and  $t = 0$  we can reduce this equation to the form †

$$z^2 t^2 = (az^2 + bzt + ct^2) (yz - xt).$$

To generate the general elliptic quartic surface, which we may call a surface of the type VI (A), we take two lines and place them in (2, 2) correspondence ‡.

\* § 74.

† Cf. Salmon, § 548.

‡ § 77.

On either line there will be four points for which the two corresponding points coincide. Take, on one of the lines, two of these points as  $X$  and  $Y$ , the corresponding double elements on the other line being  $Z$  and  $T$ . Any point of the first line is  $(\theta, 1, 0, 0)$  and any point of the second is  $(0, 0, \phi, 1)$  with a (2, 2) relation

$$a\theta^2 + \theta(c\phi^2 + d\phi + e) + b\phi^2 = 0.$$

Any point of the ruled surface is  $(\lambda\theta, \lambda, \mu\phi, \mu)$  so that  $\theta = \frac{x}{y}$  and  $\phi = \frac{z}{t}$ . The (2, 2) relation gives

$$a \frac{x^2}{y^2} + \frac{x}{y} \left( c \frac{z^2}{t^2} + d \frac{z}{t} + e \right) + b \frac{z^2}{t^2} = 0,$$

and the equation of the surface is

$$ax^2t^2 + xy(cz^2 + dzt + et^2) + by^2z^2 = 0.$$

To generate the type of elliptic quartic ruled surface with one directrix, which we may call the type VI (B), we take a line  $R$  and a plane elliptic cubic curve; this cubic passes through the point  $P$  in which  $R$  meets its plane. We then place the line and the cubic in (1, 2) correspondence with  $P$  as a united point, the pairs of points of the cubic which correspond to the points of  $R$  all having their joins passing through  $P^*$ . Let the cubic curve lie in the plane  $z = 0$ , the point  $P$  being  $x = y = z = 0$  and the tangent to the curve there  $x = z = 0$ . Then the equation to the curve may be written

$$xt^2 + (ax^2 + 2bxy + cy^2)t + Ax^3 + 3Bx^2y + 3Cxy^2 + Dy^3 = 0.$$

We take  $R$ , passing through  $P$ , to be the intersection of the planes  $x = 0$  and  $y = 0$ . We refer the points of  $R$  projectively to the pencil of lines  $x = ky$ ,  $z = 0$  in such a way that the point  $P$  of  $R$  corresponds to the line  $x = 0$  of the pencil, i.e. we take the point  $(0, 0, k, 1)$  of  $R$  to correspond to the line  $x = ky$ ,  $z = 0$  of the pencil. Then the coordinates of a point of the ruled surface are

$$(k\eta, \eta, k\lambda, 1 + \lambda),$$

where  $\eta$  is a root of

$$k + (ak^2 + 2bk + c)\eta + (Ak^3 + 3Bk^2 + 3Ck + D)\eta^2 = 0.$$

If then  $(x, y, z, t)$  denotes the point on the ruled surface we have

$$\frac{x}{y} + \frac{ax^2 + 2bxy + cy^2}{y^2} \frac{xy}{xt - yz} + \frac{Ax^3 + 3Bx^2y + 3Cxy^2 + Dy^3}{y^3} \frac{x^2y^2}{(xt - yz)^2} = 0,$$

and the equation to the surface is†

$$(xt - yz)^2 + (ax^2 + 2bxy + cy^2)(xt - yz) + (Ax^3 + 3Bx^2y + 3Cxy^2 + Dy^3)x = 0.$$

The section of this surface by a plane is a quartic curve having a tacnode at the point where  $R$  meets the plane.

\* § 79.

† Cf. Salmon, § 554.

Thus the equations of the different types of quartic ruled surfaces are as follows:

- I  $a(xz - y^2)^2 + b(xt - yz)^2 + c(yt - z^2)^2 + 2f(xt - yz)(yt - z^2) + 2g(yt - z^2)(xz - y^2) + 2h(xz - y^2)(xt - yz) = 0.$
- II (A)  $zy^2(ay + bx) + tx^2(cy + dx) = ex^2y^2.$
- II (B) As in I, but with the relation  $b^2 + 2bg - 4hf + ac = 0.$
- II (C)  $(cyz - bzx + axy + zt - tx)^2 = zx(ax - by + cz)^2.$
- III (A)  $mx^2y^2 = (x + y)(x^{2t} + y^{2z}).$
- III (B)  $(xt + yz + \mu zt)^2 = zt(x + y)^2.$
- IV (A)  $x(\gamma z + \delta t)t^2 = y(\alpha z + \beta t)z^2.$
- IV (B)  $\left. \begin{aligned} y^2z^2 + mxyz + t^2(az + bx)x = 0 \\ (yz - xy + Atx)^2 = zx(x - z + Bt)^2 \end{aligned} \right\}.$
- V (A)  $(yz - xy + Atx)^2 = zx(x - z)^2.$
- V (B)  $z^2t^2 = (az^2 + bzt + ct^2)(yz - xt).$
- VI (A)  $ax^2t^2 + xy(cz^2 + dzt + et^2) + by^2z^2 = 0.$
- VI (B)  $(xt - yz)^2 + (ax^2 + 2bxy + cy^2)(xt - yz) + (Ax^3 + 3Bx^2y + 3Cxy^2 + Dy^3)x = 0.$

**§1.** If in [5] we take two conics and place them in (1, 1) correspondence we obtain the most general rational normal quartic ruled surface.

Let the planes of the conics be  $x_0 = x_1 = x_2 = 0$  and  $x_3 = x_4 = x_5 = 0$ . We may then take a point of the first conic as  $(0, 0, 0, \theta^2, \theta, 1)$  and the corresponding point of the second as  $(\theta^2, \theta, 1, 0, 0, 0)$ . Then any point on the ruled surface has coordinates of the form

$$(\theta^2, \theta, 1, \lambda\theta^2, \lambda\theta, \lambda),$$

so that the equations of the ruled surface are\*

$$\frac{x_0}{x_1} = \frac{x_1}{x_2} = \frac{x_3}{x_4} = \frac{x_4}{x_5}.$$

If  $\theta$  is constant we have a generator of the surface; if  $\lambda$  is constant we have a conic on the surface. We thus obtain  $\infty^1$  directrix conics. The equations to the plane of any one of these are

$$x_3 = \lambda x_0, \quad x_4 = \lambda x_1, \quad x_5 = \lambda x_2,$$

so that the equations of the  $V_3^3$  formed by the planes are

$$\frac{x_0}{x_3} = \frac{x_1}{x_4} = \frac{x_2}{x_5}.$$

This clearly contains the quartic surface.

All directrix conics can be obtained as residuals of prime sections through any two fixed generators. The equation of the prime, which contains the conic whose points are  $(\theta^2, \theta, 1, k\theta^2, k\theta, 1)$  and the two generators given by  $(\alpha^2, \alpha, 1, \lambda\alpha^2, \lambda\alpha, 1)$  and  $(\beta^2, \beta, 1, \mu\beta^2, \mu\beta, 1)$ , is

$$k[x_0 - (\alpha + \beta)x_1 + \alpha\beta x_2] = x_3 - (\alpha + \beta)x_4 + \alpha\beta x_5.$$

Take a point  $(\xi_0, \xi_1, \xi_2, \xi_3, \xi_4, \xi_5)$  on  $V_3^3$  and any other point whatever  $(x_0, x_1, x_2, x_3, x_4, x_5)$ .

Then the coordinates of any point on the line joining them are

$$(\xi_0 + \mu x_0, \xi_1 + \mu x_1, \xi_2 + \mu x_2, \xi_3 + \mu x_3, \xi_4 + \mu x_4, \xi_5 + \mu x_5),$$

and if this point lies on  $V_3^3$  we have

$$\frac{\xi_0 + \mu x_0}{\xi_3 + \mu x_3} = \frac{\xi_1 + \mu x_1}{\xi_4 + \mu x_4} = \frac{\xi_2 + \mu x_2}{\xi_5 + \mu x_5}.$$

These equations are satisfied by  $\mu = 0$  as they should be. There will be two roots  $\mu = 0$ , or the line will touch the locus, if we have

$$\begin{aligned}\xi_0 x_4 + x_0 \xi_4 &= \xi_1 x_3 + x_1 \xi_3, \\ \xi_1 x_5 + x_1 \xi_5 &= \xi_2 x_4 + x_2 \xi_4, \\ \xi_2 x_3 + x_2 \xi_3 &= \xi_0 x_5 + x_0 \xi_5;\end{aligned}$$

and in virtue of the equations satisfied by  $\xi$  these may be written

$$\begin{aligned}\frac{x_4}{\xi_4} + \frac{x_0}{\xi_0} &= \frac{x_3}{\xi_3} + \frac{x_1}{\xi_1}, \\ \frac{x_5}{\xi_5} + \frac{x_1}{\xi_1} &= \frac{x_4}{\xi_4} + \frac{x_2}{\xi_2}, \\ \frac{x_3}{\xi_3} + \frac{x_2}{\xi_2} &= \frac{x_5}{\xi_5} + \frac{x_0}{\xi_0};\end{aligned}$$

so that the tangents of  $V_3^3$  at  $\xi$  lie in a solid—the tangent solid of  $V_3^3$ , whose equations are

$$\frac{x_0}{\xi_0} - \frac{x_3}{\xi_3} = \frac{x_1}{\xi_1} - \frac{x_4}{\xi_4} = \frac{x_2}{\xi_2} - \frac{x_5}{\xi_5}.$$

This tangent solid will, of course, contain the plane of  $V_3^3$  which passes through  $\xi$ ; to find the points common to the solid and  $V_3^3$  we write

$$\frac{x_0}{x_3} = \frac{x_1}{x_4} = \frac{x_2}{x_5} = \lambda$$

in the equations of the solid, which become

$$x_3 \left( \frac{\lambda}{\xi_0} - \frac{1}{\xi_3} \right) = x_4 \left( \frac{\lambda}{\xi_1} - \frac{1}{\xi_4} \right) = x_5 \left( \frac{\lambda}{\xi_2} - \frac{1}{\xi_5} \right);$$

and since  $\xi_0/\xi_3 = \xi_1/\xi_4 = \xi_2/\xi_5 = \kappa$ , say, we have

$$\frac{x_3}{\xi_3} \left( \frac{\lambda}{\kappa} - 1 \right) = \frac{x_4}{\xi_4} \left( \frac{\lambda}{\kappa} - 1 \right) = \frac{x_5}{\xi_5} \left( \frac{\lambda}{\kappa} - 1 \right).$$

Now these equations are satisfied by  $\lambda = \kappa$ ; we have thus the plane of  $V_3^3$  passing through  $\xi$ . But they are also satisfied by

$$\frac{x_3}{\xi_3} = \frac{x_4}{\xi_4} = \frac{x_5}{\xi_5},$$

which make also, from the equations of the solid,

$$\frac{x_0}{\xi_0} = \frac{x_1}{\xi_1} = \frac{x_2}{\xi_2},$$

so that we obtain the points of a straight line.

Incidentally through every point  $\xi$  of  $V_3^3$  there passes a line whose equations are

$$\frac{x_0}{\xi_0} = \frac{x_1}{\xi_1} = \frac{x_2}{\xi_2}, \quad \frac{x_3}{\xi_3} = \frac{x_4}{\xi_4} = \frac{x_5}{\xi_5}.$$



We have thus  $\infty^2$  lines on  $V_3^3$ . Each of them is a directrix, meeting every generating plane. The  $\infty^1$  generators of the quartic surface are a particular set of these lines.

The coordinates of any point on the chord of the surface which joins the points  $(\theta^2, \theta, 1, \lambda\theta^2, \lambda\theta, \lambda)$  and  $(\phi^2, \phi, 1, \mu\phi^2, \mu\phi, \mu)$  may be taken as

$$(\theta^2 + \nu\phi^2, \theta + \nu\phi, 1 + \nu, \lambda\theta^2 + \mu\nu\phi^2, \lambda\theta + \mu\nu\phi, \lambda + \mu\nu),$$

and the equations of the chord can be taken as

$$x_0 - x_1(\theta + \phi) + x_2\theta\phi = 0,$$

$$x_3 - x_4(\theta + \phi) + x_5\theta\phi = 0,$$

$$\lambda(x_1 - \phi x_2) = x_4 - \phi x_5,$$

$$\mu(x_1 - \theta x_2) = x_4 - \theta x_5,$$

the equation of any other prime through the chord being a linear combination of these four.

These equations shew that through a general point of [5] there passes one and only one chord of the surface; if the coordinates of the point are substituted in the four equations the first two will give  $\theta$  and  $\phi$  and the last two  $\lambda$  and  $\mu$ .

The only exception is when

$$\frac{x_0}{x_3} = \frac{x_1}{x_4} = \frac{x_2}{x_5},$$

or the point lies on  $V_3^3$ , as is obvious geometrically.

Thus a general rational quartic ruled surface in [4] has one double point, but there is also a rational quartic ruled surface in [4] with a double line. This latter is obtained by projecting the normal surface from a point of  $V_3^3$  and is generated by a line and a conic in (2, 1) correspondence.

# CHAPTER III

## QUINTIC RULED SURFACES

### SECTION I

#### RATIONAL QUINTIC RULED SURFACES CONSIDERED AS CURVES ON $\Omega$

82. There is a classification of quintic ruled surfaces given by Schwarz\*, the surfaces being classified by means of their double curves. When this has been obtained we can at once deduce, by the principle of duality, a second classification of quintic ruled surfaces by means of their bitangent developables. These two classifications are quite different; two surfaces which belong to different classes according to one classification can very well belong to the same class according to the other—this is sufficiently clear from Cremona's table of quartic ruled surfaces and is quite evident from the classification of the quintic surfaces themselves that we shall obtain. Thus, although Schwarz's classification is exhaustive, it is such that another, which is not included in it, can be deduced immediately from it. It is then surely desirable to obtain the more precise classification which includes both of these; and when this has been obtained the application of the principle of duality can only reproduce it.

The work is more complicated than that for the quartic ruled surfaces; one cause of this is the higher degree of the double curve and class of the bitangent developable. For the rational quartic ruled surface the double curve is a twisted cubic, and if it breaks up it must contain a line as a part of itself; this is always easy to detect when we represent the generators of the ruled surface as the points of a curve  $C$  on  $\Omega$ . But for the rational quintic ruled surface the double curve is a sextic, and if this breaks up it does not necessarily contain a line.

It will, of course, be necessary to make use of the properties of quintic curves; for some of these we can refer to a paper by Marletta†, while others that are required will be obtained in the course of the work; certain loci connected with the curve are investigated in so far as their properties are required. At the end of his paper Marletta mentions this representation of the generators of a rational quintic ruled surface, dividing the surfaces into three main classes and referring to Schwarz's paper.

\* "Über die geradlinigen Flächen fünften Grades," *Journal für Math.* 67 (1867), 23–57.

† "Sulle curve razionali del quinto ordine," *Palermo Rendiconti*, 19 (1905), 94–119.

*The rational quintic curves on a quadric  $\Omega$  in [5]*

83. Any rational quintic curve may be regarded as the projection of a rational normal quintic in [5]\*. We may project on to a [4] from a point on a chord of the normal curve; this shews that it is possible to have a rational quintic in [4] with a double point. Similarly, by projecting from a line meeting two chords of the normal curve, we may have a rational quintic in [3] with two double points and, by projecting from a line in a trisecant plane of the normal curve, we may have a rational quintic in [3] with a triple point.

We shall then divide the rational quintic curves on  $\Omega$  into seven classes:

- I. The rational normal curve in [5].
- II. The curve lies in a prime which does not touch  $\Omega$ .
- III. The curve lies in a tangent prime of  $\Omega$  but does not pass through the point of contact.
- IV. The curve lies in a tangent prime of  $\Omega$  and passes through the point of contact.
- V. The curve lies in a tangent prime of  $\Omega$  and has a double point at the point of contact.
- VI. The curve lies on the section of  $\Omega$  by an  $S_3$  through which pass two tangent primes of  $\Omega$ .
- VII. The curve lies on the quadric cone in which  $\Omega$  is met by an  $S_3$  which touches it.

84. We can at once subdivide these classes. If a rational quintic  $C$  lies in a tangent prime  $T$  touching  $\Omega$  in a point  $O$  then, when projected from  $O$  on to any solid of  $T$ , it becomes a rational quintic lying on a quadric surface. If the curve passes through  $O$  it becomes a rational quartic, while if it has a double point at  $O$  it becomes a twisted cubic. The rational curves of the third and fourth orders which lie on a quadric surface are well known; we give here those of the fifth order. There are two kinds:

(a) The residual intersection of the quadric with a quartic surface passing through three of its generators of the same system. This curve meets all the generators of one system in four points and all of the other system in one point.

(b) The residual intersection of the quadric with a cubic surface which passes through a generator and touches the quadric in two points. This curve has two double points; it meets all generators of one system in three points and all of the other system in two points.

There are also two kinds of rational quintic curves which lie on a quadric cone:

(a) The residual intersection of the cone with a cubic surface passing through a generator and touching the cone in two points. This curve has

\* § 8.

two double points; it passes through the vertex and meets every generator in two points other than the vertex.

(b) The residual intersection of the cone with a nodal cubic surface passing through a generator, the node being at the vertex of the cone. This curve has a triple point at the vertex and meets every generator of the cone in one point other than the vertex.

We are now in a position to give a more minute classification of the rational quintic curves  $C$  on  $\Omega$ .

It may be remarked here that if we have a rational quintic curve in [5] there is one trisecant plane passing through a general point of [5]\*. Hence the rational quintic in [4], if it has not a double point, has a trisecant chord.

85. We now classify the rational quintic curves which lie on  $\Omega$  as follows:

I. The rational normal curve in [5].

II. (A)  $C$  lies in a prime which does not touch  $\Omega$ .

(B)  $C$  lies in a prime which does not touch  $\Omega$ , and has a double point.

III.  $C$  lies in a tangent prime  $T$  of  $\Omega$  but does not pass through the point of contact  $O$ .

(A)  $C$  meets every plane  $\omega$  through  $O$  in four points and every plane  $\rho$  through  $O$  in one point.

(B)  $C$  meets every plane  $\omega$  through  $O$  in one point and every plane  $\rho$  through  $O$  in four points.

(C)  $C$  meets every plane  $\omega$  through  $O$  in three points, every plane  $\rho$  through  $O$  in two points, and has a double point;  $O$  lying on a chord of  $C$ .

(D)  $C$  meets every plane  $\omega$  through  $O$  in two points, every plane  $\rho$  through  $O$  in three points, and has a double point;  $O$  lying on a chord of  $C$ .

(E)  $C$  meets every plane  $\omega$  through  $O$  in three points and every plane  $\rho$  through  $O$  in two points; two chords of  $C$  passing through  $O$ .

(F)  $C$  meets every plane  $\omega$  through  $O$  in two points and every plane  $\rho$  through  $O$  in three points; two chords of  $C$  passing through  $O$ .

IV.  $C$  lies in a tangent prime  $T$  of  $\Omega$  and passes through the point of contact  $O$ .

(A)  $C$  meets every plane  $\omega$  through  $O$  in three points and every plane  $\rho$  through  $O$  in one point other than  $O$ .

(B)  $C$  meets every plane  $\omega$  through  $O$  in one point and every plane  $\rho$  through  $O$  in three points other than  $O$ .

(C)  $C$  meets every plane of  $\Omega$  through  $O$  in two points other than  $O$  and has a double point.

\* Marletta, *loc. cit.* § 12.

(D)  $C$  meets every plane of  $\Omega$  through  $O$  in two points other than  $O$  and has its trisecant passing through  $O$ .

V.  $C$  lies in a tangent prime  $T$  of  $\Omega$  and has a double point at the point of contact  $O$ .

(A)  $C$  meets every plane  $\omega$  through  $O$  in two points and every plane  $\rho$  through  $O$  in one point other than  $O$ .

(B)  $C$  meets every plane  $\omega$  through  $O$  in one point and every plane  $\rho$  through  $O$  in two points other than  $O$ .

VI.  $C$  lies on the quadric  $Q$  in which  $\Omega$  is met by an  $S_3$  through which pass two tangent primes of  $\Omega$ .

(A)  $C$  meets all generators of  $Q$  of one system in four points and all of the other system in one point.

(B)  $C$  meets all generators of  $Q$  of one system in three points and all of the other system in two points, and has two double points.

VII.  $C$  lies on the quadric cone in which  $\Omega$  is met by an  $S_3$  touching it in a point  $V$ .

(A)  $C$  passes through  $V$ , meets every generator of the cone in two points other than  $V$ , and has two double points.

(B)  $C$  has a triple point at  $V$  and meets every generator of the cone in one point other than  $V$ .

*The general surface in [3]*

86. The generators of a ruled surface of the fifth order in [3] will be represented on  $\Omega$ , if the surface is general, by the points of a rational normal quintic curve  $C$ . If  $C$  is projected from any plane on to another plane we obtain a plane quintic curve which is rational, and has therefore six double points. Hence, in the [5] containing  $C$ , a plane is met by six chords of the curve. This is true in particular for a plane of  $\Omega$ .

Hence the double curve of the ruled surface is of order six, while the bitangent developable is of the sixth class\*.

87. *The genus of the double curve.* We have just seen that there are six chords of  $C$  meeting an arbitrary plane. Hence the chords of  $C$  form a locus  $U_3^6$ , of three dimensions and of the sixth order. If we project  $C$  from a plane which meets it in one point on to another plane we obtain a rational quartic with three double points; so that the plane is only met by three chords of  $C$  other than those which pass through its point of intersection with  $C$ . Hence  $C$  is a triple curve on  $U_3^6$ . There is no double surface on  $U_3^6$  because no two chords of  $C$  can intersect in a point not lying on  $C$ .

\* See § 33.

† For if they did there would be a [4] meeting  $C$  in more than five points. If the curve is given by  $x_0:x_1:x_2:x_3:x_4:x_5 = \theta^5:\theta^4:\theta^3:\theta^2:\theta:1$ , the equations of  $U_3^6$  are

$$\left\| \begin{array}{cccc} x_0 & x_1 & x_2 & x_3 \\ x_1 & x_2 & x_3 & x_4 \\ x_2 & x_3 & x_4 & x_5 \end{array} \right\| = 0.$$

The tangent prime to  $\Omega$  at any point of  $C$  meets  $C$  in three other points; hence through any point of  $C$  there pass three chords lying on  $\Omega$ . The chords of  $C$  lying on  $\Omega$  form a ruled surface with  $C$  as a triple curve; this is merely the intersection of  $\Omega$  and  $U_3^6$ . The curve of intersection of this ruled surface with a prime is in (1, 1) correspondence with the points of the double curve and the planes of the bitangent developable\*.

Now a prime  $S_4$  meets  $C$  in five points  $a, b, c, d, e$ , and meets  $U_3^6$  in a sextic surface with these five points as triple points and no other singularities. Also  $S_4$  meets  $\Omega$  in a quadric primal through  $a, b, c, d, e$ . Hence the chords of  $C$  which lie on  $\Omega$  meet  $S_4$  in a curve of order 12 with these five points for triple points.

Project this curve from the line  $ab$  on to a plane in  $S_4$ . We obtain a sextic with three triple points. Any solid through  $ab$  will meet  $U_3^6$  in a sextic curve of which  $ab$  is a part; but if the solid contains the plane  $abc$  it will meet  $U_3^6$  in the lines  $bc, ca, ab$  and a twisted cubic passing through  $a, b, c$ . This cubic can only meet a plane through  $ab$  in one other point. Thus it is clear that no plane through  $ab$  can meet the curve of order 12 in more than one point in addition to  $a$  and  $b$ . Thus the plane sextic which we obtain has no singular points other than three triple points, and is therefore an elliptic curve. Hence the curve of order 12 is also an elliptic curve.

Thus we have proved that the double curve of the ruled surface is an elliptic curve, while the planes of the bitangent developable form an elliptic family.

**88.** *The triple point and the tritangent plane.* By a result already found † for curves of any order and genus we see that if a rational quintic curve  $C$  lies on  $\Omega$  there are two of its trisecant planes also lying on  $\Omega$ .

It is at once seen that these two planes cannot belong to the same system of planes on  $\Omega$ . For suppose, if possible, that we have two planes  $\omega_1$  and  $\omega_2$  of the same system, both trisecant to  $C$ .  $\omega_1$  and  $\omega_2$  intersect in a point  $A$  and lie in a [4], and since no [4] can meet  $C$  in more than five points  $A$  must lie on  $C$ . But  $\omega_1$  and  $\omega_2$  both lie in the tangent prime of  $\Omega$  at  $A$ , which only meets  $C$  in three points other than  $A$ . Hence our supposition is false.

There is then one plane of each system of  $\Omega$  which is trisecant to  $C$ ; so that the rational quintic ruled surface has one triple point and one tritangent plane.

**89.** We now know that the double curve of the ruled surface is an elliptic sextic curve with a triple point. If we project this from a point of itself on to a plane we obtain an elliptic quintic curve; this has a triple point which is the projection of the triple point on the sextic and it must have

\* Cf. §§ 33 and 65.

† § 35.

two further double points. Hence through any point of the double curve there pass two of its trisecants. These are precisely the two generators of the ruled surface which intersect in this point of the double curve; and we can therefore generate the surface by the trisecants of an elliptic sextic curve with a triple point\*.

It is not difficult to see which are the three generators of the surface passing through the triple point. The plane containing two of the tangents at the triple point will meet the curve in a single further point  $P$ , and if the curve is projected from  $P$  on to a plane the triple point becomes, on the resulting quintic curve, a triple point at which two tangents coincide. Thus this quintic curve will only have one other double point, so that only one proper trisecant of the sextic curve passes through  $P$ . The line joining  $P$  to the triple point is a generator of the ruled surface. The other two generators passing through the triple point are at once determined similarly.

*Further consideration of the surfaces whose generators do not belong to a linear complex*

90. Considering again the rational normal quintic  $C$ , let us take a plane meeting  $C$  in a single point  $P$ . In general there are three points of this plane, other than  $P$ , through which pass chords of  $C$ . But suppose that there is an infinite number of chords of  $C$  which meet the plane; the plane then meets  $U_3^6$  in a curve  $\Gamma$ .

If we project  $C$  from  $P$  on to a space [4] we obtain a rational normal quartic, and there is no point of [4], which is not on the curve, through which two chords of the curve can pass. Hence no line through  $P$  can meet  $U_3^6$  in more than one point other than  $P$ . This shews that  $\Gamma$ , if of order  $n$ , has a point at  $P$  of multiplicity  $n - 1$ , and  $n \geq 2$ .

If we project  $C$  from a line in the plane of  $\Gamma$  on to a space [3] we obtain a rational quintic with  $n$  double points. Hence we must have  $n \leq 2$ .

\* For the degree of a ruled surface formed by trisecants of a curve see Zeuthen, *Annali di Matematica* (2), 3 (1869), 183–185.

The result given by Zeuthen is only true for curves having the singularities which he prescribes; it does not hold, for example, for a curve with a triple point. In the first correspondence proof given by Zeuthen a triple point gives rise to 6 ( $m - 3$ ) coincidences of the points  $x$  and  $n$ ; and, taking an elliptic sextic curve with a triple point, we find a quintic ruled surface as we should do.

If we have an elliptic sextic with three double points, then Zeuthen's formula gives four for the degree of the ruled surface formed by the trisecants. There are two trisecants passing through any point of the curve, and this would seem at first to be at variance with our work on quartic ruled surfaces, where we shewed that the double curve was of the third order. But this quartic ruled surface is none other than the unique quadric containing the curve, counted twice. The trisecants are the generators of this quadric. Since a quadric which is made to contain  $2n$  arbitrary points of an elliptic curve of order  $n$  will contain the curve entirely, this quadric is determined by the three double points and six other points of the curve.

There is no quadric containing the curve with a triple point.

Thus  $n = 2$ , and  $\Gamma$  is a conic passing through  $P$ .

The chords of  $C$  through the points of  $\Gamma$  form a ruled surface.

There is no doubt that such planes actually exist. For consider an involution  $g_2^1$  of pairs of points on  $C$ . It is known that the chords joining the pairs of  $g_2^1$  form a quartic ruled surface\*. This ruled surface cannot have a directrix line; for, if it had, any three of its generators, having a common transversal, would belong to a [4]; and no [4] can meet  $C$  in six points. Hence it has  $\infty^1$  directrix conics, one conic passing through each point of the surface†. All these conics can be obtained as residuals of prime sections through any two fixed generators, so that each conic meets  $C$  in one point.

The planes of these conics are planes such as we require; such a plane will be called a secant plane (of  $U_3^6$ ). There are  $\infty^2$  involutions  $g_2^1$  on  $C$ ; each gives a quartic ruled surface lying on  $U_3^6$  and  $\infty^1$  secant planes of  $U_3^6$ . There are thus  $\infty^3$  secant planes of  $U_3^6$ . Through a general point of [5] there will pass a finite number of these secant planes; but through a point of  $U_3^6$  there pass  $\infty^1$ , and through a point of  $C$  there pass  $\infty^2$ .

If any point  $X$  is taken in a secant plane which meets  $C$  in  $P$ , the line  $XP$  meets the conic  $\Gamma$  in a second point  $P'$  through which there passes a chord  $k$  of  $C$ . The plane  $PkX$  is thus the trisecant plane of  $C$  which passes through  $X$ . Hence if a secant plane passes through  $X$  it must contain one of the three lines which join  $X$  to the three points of  $C$  lying in the trisecant plane through  $X$ .

**91. The geometry of the secant planes.** There are  $\infty^3$  secant planes, each meeting  $C$  in one point; if we have a quadric  $\Omega$  containing  $C$  it will not, in general, be possible to choose the parameters on which a secant plane depends so that it lies entirely on  $\Omega$ . But the general quadric  $\Omega$  containing  $C$  is linearly dependent from ten quadrics, so that we should be able to choose  $\Omega$  to contain a secant plane.

The condition that a quadric in [5] should represent the lines of a space [3] is simply that it should be a general quadric and not a cone; if we take a general quadric passing through  $C$ , and containing a secant plane, to represent the lines of a space [3], the curve  $C$  will represent the generators of a quintic ruled surface in [3] whose double curve and bitangent developable

\* See footnote to § 19. If the  $g_2^1$  is given by  $A - B(\theta_1 + \theta_2) + C\theta_1\theta_2 = 0$  on the curve  $x_0:x_1:x_2:x_3:x_4:x_5 = \theta^5:\theta^4:\theta^3:\theta^2:\theta:1$ , the quartic ruled surface is

$$\left\| \begin{array}{ccc} x_0 & x_1 & x_2 \\ x_1 & x_2 & x_3 \\ x_2 & x_3 & x_4 \\ x_3 & x_4 & x_5 \\ A & B & C \end{array} \right\| = 0.$$

† § 43.



break up, because the chords of  $C$  which lie on the quadric form a ruled surface which breaks up.

Suppose now that  $\Omega$  contains a secant plane meeting  $C$  in a point  $P$ , this being a  $\rho$ -plane. The involution determined on  $C$  by the conic  $\Gamma$  in which the secant plane meets  $U_3^6$  gives a chord  $PA$  through  $P$ ; this, in common with all the other generators of the quartic surface, lies on  $\Omega$ . Through  $P$  there pass two other chords  $PQ, PR$  which lie on  $\Omega$  (Fig. 4).

Through  $Q$  there passes a chord of  $C$  meeting  $\Gamma$  in  $Q'$  and through  $R$  there passes a chord of  $C$  meeting  $\Gamma$  in  $R'$ . Then the three lines  $PQ, QQ', PQ'$  lie on  $\Omega$ , so that  $PQQ'$  is a plane  $\omega_1$  which is trisecant to  $C$ . Similarly




Fig. 4.

$PRR'$  is a plane  $\omega_2$  which is trisecant to  $C$ . But there is only one plane  $\omega$  trisecant to  $C$ , so we must conclude that this plane is  $PQR$ , the chord  $QR$  meeting  $\Gamma$ .

Hence, if we have a quadric  $\Omega$  containing  $C$  and also a secant plane  $\rho$  which meets  $C$  in  $P$ , the plane  $\omega$  of  $\Omega$  which is trisecant to  $C$  passes through  $P$ .

Let this trisecant  $\omega$ -plane meet  $C$  in  $P, Q, R$  and suppose that  $\Omega$  contains two secant planes  $\rho_1$  and  $\rho_2$  through  $P$ . Each of these determines an involution on  $C$ , the common pair of the two involutions being  $Q, R$ . Hence we have two other chords  $PP_1$  and  $PP_2$  through  $P$  and lying on  $\Omega$ , one arising from each involution. But it is impossible for four chords  $PP_1, PP_2, PQ, PR$  all passing through  $P$  to lie on  $\Omega$ .

Hence it is impossible for  $\Omega$  to contain two secant  $\rho$ -planes passing through  $P$ .

The planes of the directrix conics of a quartic ruled surface in [5] form a locus  $V_3^3$  of three dimensions and the third order\*, and therefore meet an arbitrary solid  $S_3$  in the points of a twisted cubic.

Suppose then that  $\Omega$  contains  $C$  and also the quartic ruled surface arising from an involution  $g_2^1$  on  $C$ . Then a solid  $S_3$  meets  $\Omega$  in a quadric  $Q$  and meets the  $V_3^3$  formed by the planes of the directrix conics in a twisted cubic which meets  $Q$  in six points. Four of these six points will be the points of intersection of  $S_3$  with the quartic surface; the other two give planes which lie on  $\Omega$  entirely, since each meets it in a conic and a point. Since these two planes cannot intersect they are of opposite systems.

Hence, if  $\Omega$  contains a secant  $\rho$ -plane it must also contain a secant  $\omega$ -plane; the two planes meet  $U_3^6$  in conics belonging to the same quartic ruled surface. This secant  $\omega$ -plane will meet  $C$  in a point  $P'$ , this being one of the three points in which  $C$  is met by its trisecant  $\rho$ -plane.

92. Since the quadrics containing  $C$  are linearly dependent from ten quadrics there will be five linearly independent quadrics containing  $C$  and a secant plane. We may then suppose that there is a general quadric  $\Omega$  containing  $C$  and two of its secant planes, these planes being of opposite systems on  $\Omega$  and both arising from the same involution on  $C$ .

Then the ruled surface formed by the chords of  $C$  which lie on  $\Omega$  breaks up into a quartic ruled surface passing through  $C$  (and met by the secant planes on  $\Omega$  in two of its directrix conics) and an octavic ruled surface having  $C$  as a double curve. Through any point of  $C$  there pass one generator of the quartic and two of the octavic; by considering the correspondence thus set up on  $C$  it is seen that these ruled surfaces have two common generators, say  $g$  and  $g'$ .

The section of the composite ruled surface by a [4] gives an octavic curve with five double points and a quartic curve passing through these points; both these curves are rational.

It is easily shewn† that any general plane of  $\Omega$  will meet the quartic ruled surface in two points and the octavic in four points. Thus the points of  $C$  represent the generators of a quintic ruled surface in [3]; the double curve breaks up into a conic and a rational quartic, while the bitangent developable breaks up into a quadric cone and a developable of the fourth class whose planes form a rational family.

Suppose that  $PQR$  is the  $\omega$ -plane which is trisecant to  $C$  and let the secant  $\rho$ -plane pass through  $P$ . Then  $PQ$  and  $PR$  are generators of the

\* § 69.

† A plane  $\omega$  of  $\Omega$  meets the secant  $\omega$ -plane in a point  $O$ . These two planes lie in the tangent [4] of  $\Omega$  at  $O$ , which meets  $\Omega$  in a quadric point-cone and the quartic ruled surface in a directrix conic and two generators. These two generators each meet the first  $\omega$ -plane.

octavic surface while  $QR$  is on the quartic. Thus  $PQR$  represents the triple point of the ruled quintic; the double conic passes through this point while the double quartic has a double point there. Further, the two parts of the double curve intersect in two other points, these being represented on  $\Omega$  by the  $\omega$ -planes through  $g$  and  $g^*$ .

Similarly the quartic developable has a double plane which touches the quadric cone (this being the tritangent plane of the ruled quintic), while they have also two other common tangent planes.

The plane of the double conic is represented on  $\Omega$  by the secant  $\rho$ -plane; it therefore contains a generator of the ruled quintic passing through the triple point. The vertex of the quadric cone lies on a generator which is in the tritangent plane.

**93.** *The surface with three double conics.* The quadrics containing  $C$  and a secant plane are linearly dependent from five quadrics. Such a quadric meets any other secant plane in two definite points; one of these is on  $C$  and the other on that chord of  $C$  which belongs to both the involutions determined on  $C$  by the two secant planes. If the quadric is made to pass through four further points of the second secant plane (not lying on the same conic with the two fixed points) it will contain the secant plane entirely. Thus there will be a quadric containing  $C$  and two secant planes.

Suppose then that  $\Omega$  contains  $C$ , a secant plane  $\rho_1$  meeting  $C$  in  $P$  and a secant plane  $\rho_2$  meeting  $C$  in  $Q$ . There is a plane  $\omega$  of  $\Omega$  meeting  $C$  in three points  $P, Q, R$ . The ruled surface formed by the chords of  $C$  lying on  $\Omega$  is in general of order 12 with  $C$  for a triple curve; here it consists of a quartic ruled surface containing  $C$  and the chord  $QR$ , a quartic ruled surface containing  $C$  and the chord  $RP$ , and therefore also of a third quartic ruled surface containing  $C$  and the chord  $PQ$ . There is a directrix conic of this last surface passing through  $R$ ; the plane of this conic meets  $\Omega$  in the conic itself and also in a line of the plane  $PQR$ ; it therefore lies on  $\Omega$  entirely and is a secant plane  $\rho_3$ .

$\Omega$  will also contain three secant planes  $\omega_1, \omega_2, \omega_3$  belonging to the same three quartic surfaces; these meet  $C$  in three points  $P', Q', R'$  and the trisecant plane  $P'Q'R'$  is a  $\rho$ -plane of  $\Omega$ .

A general plane of  $\Omega$  of either system meets each quartic ruled surface in two points. Thus the curve  $C$  represents the generators of a quintic ruled surface in a space [3]; the double curve of this surface consists of three conics and the bitangent developable of three quadric cones. The planes of the conics intersect in the point represented on  $\Omega$  by the plane  $PQR$ , these planes themselves being represented by the planes  $\rho_1, \rho_2, \rho_3$ .

\* Cf. Schwarz, *loc. cit.* p. 37.

The intersection of the planes of the three conics lies on all the conics, and is the triple point. Similarly the tritangent plane touches all the cones and is the plane joining the three vertices.

Any two involutions on  $C$  have a common pair of points, so that any two of the three quartic surfaces have a common generator. Thus any two of the three double conics have a second common point on the line of intersection of their planes, this point being represented on  $\Omega$  by the plane  $\omega$  passing through the common generator of the two quartic surfaces concerned\*. Similarly any two of the three quadric cones have a second common tangent plane passing through the line joining their two vertices.

*Surfaces whose generators belong to a linear complex  
which is not special*

94. We turn now to the quintic ruled surface in [3] whose generators are represented by the points of a curve  $C$  on  $\Omega$ , this curve lying in a space [4] which does not touch  $\Omega$ . We assume, unless the contrary is stated, that  $C$  has not a double point; we are dealing then with a surface of the type II (A).

If we project  $C$  from a line on to a plane in [4] we obtain a quintic curve which, as a rational curve, must have six double points. Hence six chords of  $C$  meet the line, so that the chords of  $C$  form a locus  $V_3^6$  of three dimensions and the sixth order.

Any plane of  $\Omega$  meets the space [4] in a line lying on  $\Omega$ ; this line is met by six chords of  $C$ , and no other chords of  $C$  can meet the plane. Hence we have a quintic ruled surface whose double curve is of the sixth order and whose bitangent developable is of the sixth class.

95. *The locus  $V_3^6$ .* The curve  $C$  has a trisecant chord  $t$ † meeting it in three points  $P, Q, R$ .

If we project  $C$  from a line which meets it on to a plane of [4] we obtain a rational quartic with three double points; hence the line is met by three chords of  $C$  other than those which pass through its point of intersection with  $C$ . Hence  $C$  is a triple curve on  $V_3^6$ .

If we project from a line meeting  $t$  we obtain a rational quintic with a triple point and three double points, so that the line is met by three chords of  $C$  other than  $t$ . Hence  $t$  is a triple line on  $V_3^6$ .

If we project from a line through  $P$  we obtain a rational quartic with three double points, one of these arising from  $t$ . Hence the line is only met by two chords of  $C$  which do not pass through  $P$ , so that  $P, Q, R$  are quadruple points on  $V_3^6$ .

If we project  $C$  from one of its chords on to a plane we obtain a cubic with a double point; hence every chord of  $C$  is met by one other chord in

\* Cf. Schwarz, *loc. cit.* p. 44.

† Marletta, *loc. cit.* p. 101.

a point not on  $C$ , these two chords lying in a quadrisecant plane of  $C$ . This at once suggests a double surface  $F_2$  on  $V_3^6$ . The order of  $F_2$  is the number of points in which it is met by a plane of [4]; the plane meets  $V_3^6$  in a sextic curve having this same number of double points.

This number of double points is known when we know the genus of a plane section of  $V_3^6$ . Now, since  $C$  is the projection of a rational normal curve in [5],  $V_3^6$  is the projection of the locus  $U_3^6$  of § 87. Hence the genus of the plane sections of  $V_3^6$  is the same as that of the sections of  $U_3^6$  by solids.

We then consider in [5] the curve in which a solid  $\Sigma$  is met by the chords of a rational normal quintic curve  $C_0$ . If we take any four trisecant planes of  $C_0$  then it is clear that the chords of  $C_0$  are obtained as the intersections of corresponding primes of four doubly infinite projectively related systems of primes, each of the four trisecant planes being the base of one of the doubly infinite systems. The section by  $\Sigma$  then gives four projectively related "stars" of planes, each with a base point. If  $\Sigma$  is met by a chord of  $C_0$  the point of meeting is an intersection of corresponding planes of the four stars. But the locus of such points is known to be a sextic curve of genus *three\**, and this is then the section† of  $U_3^6$  by  $\Sigma$ .

The plane sections of  $V_3^6$  are therefore also of genus three, and thus have *seven* double points. Hence we have on  $V_3^6$  a double surface  $F_2^7$  of order seven.

The section of  $V_3^6$  by a quadrisecant plane of  $C$  consists of the six chords of  $C$  lying in the plane; the section of  $F_2^7$  consists of the four points of  $C$  together with the three diagonal points of the quadrangle formed by them.

A trisecant plane  $PXY$  of  $C$  meets  $V_3^6$  in the three chords  $PX$ ,  $PY$ ,  $XY$  and a cubic curve having a double point at  $P$ . This curve meets  $XY$  again in a point  $W$ ; the section of  $F_2^7$  by the plane  $PXY$  consists of the points  $X$ ,  $Y$ ,  $W$  together with  $P$  counted four times.  $P$ ,  $Q$ ,  $R$  are quadruple points on  $F_2^7$ .

A trisecant plane  $XYZ$  of  $C$  meets  $V_3^6$  in the three chords  $YZ$ ,  $ZX$ ,  $XY$  and in a cubic curve passing through  $X$ ,  $Y$ ,  $Z$  and having a double point  $O$ . This curve meets the lines  $YZ$ ,  $ZX$ ,  $XY$  again in points  $U$ ,  $V$ ,  $W$ . The plane  $XYZ$  meets  $F_2^7$  in the seven points  $X$ ,  $Y$ ,  $Z$ ,  $U$ ,  $V$ ,  $W$ ,  $O$ .

\* Schur, *Math. Ann.* 18 (1881), 15.

† The surface which is the prime section of  $U_3^6$  is a sextic surface in [4] generated by four doubly infinite systems of solids projectively related to each other. It is a particular case of that considered by Veronese, *Math. Ann.* 19 (1882), 232–233. The projection of this sextic surface in [4] on to a solid is the same as the surface which is the section of  $V_3^6$  by a solid. This surface in [3] has a double curve of order 7 (the section of  $F_2^7$ ) and, on this double curve, a triplanar point, this being the intersection of [3] with the trisecant  $t$  of  $C$ . Further, since the solid meets  $C$  in five points, and the ten chords joining these points are lines of  $V_3^6$ , the sextic surface has ten lines on it.

The section of  $V_3^6$  by a plane through  $t$  and a point  $X$  of  $C$  consists of  $t$  counted three times together with the three chords  $XP$ ,  $XQ$ ,  $XR$ . The point  $X$  is the only point of  $F_2^7$ , other than the points of  $t$ , which can lie in this plane. A solid containing  $t$  will meet  $F_2^7$  in  $t$  counted three times and in a quartic curve passing through  $P$ ,  $Q$ ,  $R$ .

$t$  is a triple line on  $F_2^7$ , but  $C$  is only a simple curve.

96. *The genus of the double curve.* A quadric containing  $C$  will also contain  $t$ ; its intersection with  $F_2^7$  therefore consists of  $C$ ,  $t$  counted three times, and a sextic curve; the total intersection being a curve of order 14. Hence those chords of  $C$  which lie on  $\Omega$  form a ruled surface of order 12 in [4], the intersection of  $\Omega$  and  $V_3^6$ . This ruled surface has  $C$  for a triple curve and  $t$  for a triple generator; it has also a double sextic curve meeting every generator in one point and passing through  $P$ ,  $Q$ ,  $R$ .

If we take the section of this ruled surface by a [3] we obtain a curve in (1, 1) correspondence with the double curve of the ruled quintic. The curve is of order 12, having six triple points and six double points; it is the intersection of a quadric and a sextic surface, so that it meets each generator of the quadric in six points. If we project the curve from a point of the quadric on to a plane of the [3] we obtain a curve of order 12 with six triple points, six double points, and two sextuple points; its genus is therefore

$$55 - 18 - 6 - 30 = 1,$$

so that it is elliptic.

Thus the ruled quintic has an elliptic double curve of order 6, while the planes of its bitangent developable form an elliptic family of class 6. Moreover, the double curve has a triple point; the three generators passing through it lie in a plane and are represented on  $\Omega$  by the points  $P$ ,  $Q$ ,  $R^*$ . Also the bitangent developable has a tritangent plane; the three generators therein pass through a point. For this surface the tritangent plane passes through the triple point; the plane  $\rho$  of  $\Omega$  passing through  $t$  represents the tritangent plane, while the plane  $\omega$  of  $\Omega$  passing through  $t$  represents the triple point.

It is clear, either by the geometry of the planes on  $\Omega$  or by the geometry of the linear complex, that at a triple point on any ruled surface whatever whose generators all belong to a linear complex the three generators which intersect there lie in a plane. Similarly, the three generators in any tritangent plane pass through a point.

97. *An associated ruled surface.* We have just seen that the prime sections of the ruled surface formed by the chords of  $C$  which lie on  $\Omega$  are elliptic curves; this enables us to shew that the sextic curve which forms part of the intersection

\* Cf. Marletta, *loc. cit.* p. 117.

of  $\Omega$  and  $F_2^7$  is also an elliptic curve\*. Its points thus represent the generators of an elliptic sextic ruled surface  $f^6$  in the original space [3]; this passes through the double curve of the quintic ruled surface and is touched by the planes of the bitangent developable of the quintic surface, and it contains the three generators which meet in the triple point and lie in the tritangent plane.

Suppose then that we have in [3] a rational quintic ruled surface whose generators belong to a linear complex. Take any point  $x$  on the double curve, and the plane of the two generators which meet in  $x$ . Then this plane meets the surface again in a cubic curve. This cubic meets each generator in three points; one point on each generator is a point where the plane (which is a bitangent plane) touches the surface, the other two are points of the double curve. The remaining point  $y$  in which the plane meets the double curve is a double point on the cubic curve. If we had started with the point  $y$  instead of the point  $x$  then the bitangent plane through  $y$  would have met the surface in two generators and a cubic curve having a double point at  $x$ . The lines such as  $xy$ , chords of the double curve and axes of the bitangent developable, generate an elliptic sextic ruled surface  $f^6$  whose generators belong to the same linear complex as do those of the quintic surface.

The curve of intersection of the two ruled surfaces, in all of order 30, will consist of the double curve of the quintic surface counted twice, a certain number of common generators, and another curve. Since every generator of  $f^6$  can only meet the quintic surface in one point other than the two points in which it meets the double curve, this other curve meets the generators of  $f^6$  each in one point and is therefore elliptic. Moreover, it meets each generator of the quintic ruled surface in three points, and an elliptic curve on a rational quintic ruled surface which meets each generator in three points is of order  $9\ddagger$ . Hence the two ruled surfaces have in common this curve of order 9, the double curve of the quintic ruled surface counted twice, and nine common generators. These last consist of the three generators through the triple point of the quintic surface and six others $\ddagger$ .

\* There is a (1, 2) correspondence between the points of this curve and the chords of  $C$  on  $\Omega$ , to which we can apply Zeuthen's formula with  $a = 1$ ,  $a' = 2$ ,  $p' = 1$ ,  $\eta = \eta' = 0$ .

The assumption that  $\eta = 0$  is the assumption that the two chords of  $C$  which intersect at a point of the sextic curve never coincide. We shall see later that there are four special chords of  $C$  lying on  $F_2^7$ ; if  $\Omega$  were to contain any of these chords we could not assume  $\eta$  to be zero. But, in general, the assumption is true. The points  $P, Q, R$  do not count as branch-points of the correspondence.

If  $\Omega$  contained one or more of the special chords of  $C$  we should not have a sextic curve of intersection on  $F_2^7$ .

† § 17.

‡ This result shews that the sextic curve of intersection of  $\Omega$  and  $F_2^7$  has six intersections with  $C$  other than  $P, Q, R$ .

In the first paper of Segre's, referred to in § 17, we find a formula for the number of intersections of two simple (non-multiple) curves on a ruled surface. There is a corresponding formula for the number of intersections of two multiple curves on a ruled surface, but it is subject to modification as certain intersections may be included more than once. For two curves of orders  $m$  and  $m'$  of multiplicities  $s$  and  $s'$  on a ruled surface of order  $n$ , the curves meeting each generator of the surface in  $k$  and  $k'$  points respectively, the formula gives the number of intersections as

$$i = msk' + m's'k - nkk'.$$

If we apply this formula to the triple quintic curve and the double sextic curve

98. Suppose that we take any point  $X$  on  $C$ ; is there a line through  $X$  which is not a chord of  $C$  and yet lies on  $V_3^6$ ? If we project  $C$  from  $X$  on to a space [3] we obtain a rational quartic without a double point. If such a line through  $X$  existed it would meet [3] in a point through which an infinite number of chords of the quartic pass; this would then be the vertex of a quadric cone containing the quartic, and there are no such cones\*. But this reasoning does not hold for the three points  $P, Q, R$ . If we project from  $P$  we obtain a rational quartic with a double point; there are two quadric cones containing this quartic whose vertices are not at the double point†, so that we expect two lines through  $P$  lying on  $V_3^6$ . We shall call these lines axes.

We can also obtain some information as to these axes by projecting a rational normal quintic from a point on to a space [4]. If we project from  $X$  (see the end of § 90) any axes of the projected curve must arise from secant planes through  $X$ ; so that we get a finite number of axes, any one of which must pass through one of  $P, Q, R$ . Further, since any line in the secant plane through  $X$  meets the conic in that plane in two points there are two chords of the projected curve passing through any point of an axis.

The chords of  $C$  meeting any axis form a quartic ruled surface in [4] on which the axis is a double line—the projection of a normal quartic ruled surface in [5] from a point in the plane of one of its directrix conics.

99. *The geometry of the axes.* We have two axes through each of the points  $P, Q, R$ , giving six in all; call them  $p, p', q, q', r, r'$ , where  $p$  and  $p'$  pass through  $P$ . They lie not only on  $V_3^6$  but also on  $F_2^7$ .

On any axis there are two points such that there is only one chord of  $C$  passing through each of them; if the axis is the projection of a conic  $\Gamma$  from a point  $X$  in a secant plane in [5], these two points arise from the two tangents of  $\Gamma$  which pass through  $X$ . We may call these “pinch-points.”

There is a cubic ruled surface  $\phi$  containing  $C$ ; the directrix of  $\phi$  is the trisecant of  $C$ , while every generator of  $\phi$  meets  $C$  in one point. The  $\infty^2$  conics of  $\phi$  give the  $\infty^2$  quadrisecant planes of  $C$ ‡. No two quadrisecant

which lie on the ruled surface of order 12 formed by those chords of  $C$  which lie on  $\Omega$ , it appears that the two curves have fifteen intersections. But these really consist of the three intersections  $P, Q, R$  and the six others each counted twice. See the Note at the end of the volume.

\* Salmon, *Geometry of Three Dimensions* (Dublin, 1914), p. 359, and above § 63.

† We can regard the quartic curve as the intersection of the two quadrics

$$2xy + z^2 - t^2 = 0 \quad \text{and} \quad y^2 - m^2z^2 + n^2t^2 = 0.$$

‡ Marletta, *loc. cit.* pp. 101, 102. For the cubic ruled surface see § 48 above.



planes of  $C$  can intersect except on  $\phi$ . The  $\infty^1$  quadrisecant planes of  $C$  which pass through any point of  $\phi$  lie on a quadric point-cone, and all belong to one system of its planes.

Consider the  $V_3^3$  formed by the planes of the directrix conics of a normal quartic ruled surface in [5]; project it on to a space [4] from a point  $O$  of itself,  $O$  lying in the plane of a directrix conic  $\Gamma$ . We obtain a quadric point-cone; one system of planes of this cone is the projection of the planes of the directrix conics; the other system of planes is the projection of those solids which contain pairs of generators of the quartic surface and pass through  $O$ , meeting the plane of  $\Gamma$  in lines through  $O^*$ .

This result shews that the quadrisecant planes passing through the points of an axis all intersect in the same point, forming one system of planes of a quadric point-cone.

Any axis determines an involution on  $C$ ; denote, for example, by  $(p)$  the involution determined by  $p$ . Since any two involutions on  $C$  have a common pair of points the involutions  $(p)$  and  $(r)$  give a chord joining their common pair; this chord meeting both  $p$  and  $r$ . But any chord of  $C$  is met by only *one* chord beside those passing through the two points in which it meets  $C$ . Hence the chord must meet at least one of  $p$  and  $r$  in a pinch-point.

The quadrics in [4] containing  $C$  are linearly dependent from four quadrics. Thus we can make such a quadric contain  $p$  by assigning two points of  $p$ ; there is a pencil of quadrics passing through  $C$  and  $p$ . Moreover, such a quadric, besides passing through  $Q$  and  $R$ , meets each of  $q, q', r, r'$  in fixed points†. Hence we can find a quadric of the pencil containing  $q$ . This quadric will contain the point of  $r$  which lies on the chord common to  $(q)$  and  $(r)$ , so that if this point is different from the intersection of  $r$  with the chord common to  $(p)$  and  $(r)$  the quadric will contain  $r$  entirely. There is a similar statement concerning  $r'$ . But it is impossible for the quadric to contain  $p, q$  and both the axes  $r$  and  $r'$ ; for then we should have chords of  $C$  lying on it and forming a ruled surface of order at least  $4 \times 4 = 16$ , which is impossible. We must therefore conclude that the quadric containing  $C, p, q$  contains one of  $r$  and  $r'$ , say  $r$ , while the chord common to  $(p)$  and  $(r')$  and the chord common to  $(q)$  and  $(r')$  meet  $r'$  in the same point. We may denote the quadric by  $pqr$ .

Denote by  $(pq)r'$  the fact that the chord common to  $(p)$  and  $(r')$  and the chord common to  $(q)$  and  $(r')$  meet  $r'$  in the same point.

\* Through each point of  $V_3^3$  there passes a line meeting all its generating planes. See the footnote to § 9 of Segre's paper, "Sulle varietà normali a tre dimensioni," *Atti Torino*, 21 (1885), 95. Also § 81 above.

† It meets  $q$  for instance in the point where it is met by the chord common to  $(p)$  and  $(q)$ . It does not, however, meet  $p'$  in a fixed point, because the chord common to  $(p)$  and  $(p')$  is  $QR$ .

There are four quadrics

$$pqr, \quad pq'r', \quad p'qr', \quad p'q'r,$$

which contain  $C$  and three axes, and we have

$$\begin{array}{cccc} (pq) r', & (pq') r, & (p'q) r, & (p'q') r', \\ (qr) p', & (qr') p, & (q'r) p, & (q'r') p', \\ (rp) q', & (rp') q, & (r'p) q, & (r'p') q', \end{array}$$

the twelve elements of this array being determined by the first one.

Now from the three statements

$$(pq) r', \quad (qr') p, \quad (r'p) q,$$

we see that the involutions  $(p)$ ,  $(q)$ ,  $(r')$  have a common chord (we cannot have three co-planar chords common to these pairs of involutions); this meets the three axes  $p$ ,  $q$ ,  $r'$  which must therefore lie in a solid since they have two transversals. We have four such solids which we may denote by

$$[pqr'], \quad [pq'r'], \quad [p'qr], \quad [p'q'r'].$$

Let us denote the four common chords of the sets of three involutions by

$$\overline{pqr'}, \quad \overline{pq'r'}, \quad \overline{p'qr}, \quad \overline{p'q'r'}.$$

The first of these, for example, is a transversal of the axes  $p$ ,  $q$ ,  $r'$ .

The points in which  $\overline{pqr'}$  meets the axes  $p$ ,  $q$ ,  $r'$  are pinch-points on at least two of them; since the six axes are on the same footing it is probable that all the three points of intersection will be pinch-points.

We are thus led to the supposition that the unique quadrisecant plane through  $\overline{pqr'}$  does not contain any other chord, but touches  $C$  at both ends of the chord  $\overline{pqr'}$ . Thus we find four pairs of intersecting tangents of  $C$ .

The existence of these pairs of intersecting tangents can be seen in another way, for the tangents form a ruled surface which is the projection, from a point, of the ruled surface formed by the tangents of a normal curve in [5]. Now any surface whatever in [5] has a finite number of "apparent double points"\*; the number will be reduced by the existence of a cuspidal curve on the surface, but we can still expect a certain number, and on projection these give pairs of intersecting tangents of  $C$ .

The number of pairs of intersecting tangents of  $C$  can be found directly.

Take any point  $Q$  of  $C$ . The quadrisecant planes of  $C$  which pass through  $Q$  are the planes of the  $\infty^1$  directrix conics of the cubic ruled surface  $\phi$  which pass through  $Q$ . Since any one of these conics is determined by one point of any generator of  $\phi$  they cut out on  $C$  an ordinary involution of sets of three points, and such an involution has four double points. Hence there are four quadrisecant planes which pass through  $Q$  and contain tangents of  $C$ .

\* The only surface in [5] without any apparent double points is the surface of Veronese. See Severi, *Palermo Rendiconti*, 15 (1901), 41, 42.

Now let us establish a correspondence between the points  $P$  and  $Q$  of  $C$ ; two points corresponding when the tangent at  $P$  meets a chord through  $Q$  other than  $QP$ . We have just seen that to any position of  $Q$  there correspond four positions of  $P$ . Also a tangent of  $C$  is met by one chord other than those passing through its point of contact\*, so that to any position of  $P$  there correspond two positions of  $Q$ . We have then, on the rational curve  $C$ , a (4, 2) correspondence between  $P$  and  $Q$ ; there are eight points  $P$  for which the two corresponding points  $Q$  coincide†. Hence there are eight tangents of  $C$  which are met by other tangents, and these divide into four pairs of intersecting tangents.

The tangents of  $C$  therefore form a (developable) ruled surface whose cuspidal curve  $C$  is of order 5 and which has four double points.

The chords which join the points of contact of the pairs of intersecting tangents are none other than the four chords

$$\overline{pqr'}, \quad \overline{pq'r}, \quad \overline{p'qr}, \quad \overline{p'q'r'}.$$

The four planes which touch  $C$  twice can be taken in pairs in six ways; the six solids formed by the six pairs of chords of contact each contain one of the six axes.

The four chords of contact lie on  $F_2^7$ .

A bitangent plane of  $C$  meets  $V_3^6$  in the two tangents together with the chord of contact counted four times; it meets  $F_2^7$  in the chord of contact and the point of intersection of the two tangents.

There is one more remark which may be made concerning the configuration of the six axes.

It is easily shewn, in ordinary space, that in general there are four quadric cones passing through the curve of intersection of two quadrics. But if the two quadrics touch their intersection is a rational quartic curve with a double point; the four cones become the cone projecting the curve from its double point (this cone counting doubly) with two other cones. Now the plane of the two tangents to the quartic at its double point contains the vertices of these last two cones.

Hence the solid containing the tangents of  $C$  at  $Q$  and  $R$  contains the axes  $p$  and  $p'$ , with two other similar results. We thus have three solids which we may denote by

$$[pp't_Qt_R], \quad [qq't_Rt_P], \quad [rr't_Pt_Q].$$

100. Consider now the quintic ruled surface whose generators are represented by the points of a rational quintic  $C$  in [4],  $C$  lying on a quadric which contains also the axis  $p$ ‡.

\* For on projecting  $C$  from the tangent on to a plane we obtain a cubic with a double point.

† If we have an  $(r, s)$  correspondence between points  $P$  and  $Q$  on a rational curve then there are  $2r(s - 1)$  positions of  $P$  for which two of the  $s$  points  $Q$  coincide (§ 13).

‡ This quadric is regarded as a prime section of  $\Omega$ .

The ruled surface formed by the chords of  $C$  which lie on the quadric here breaks up into a quartic ruled surface passing through  $C$ , and having  $p$  for a double line, and an octavic ruled surface having  $C$  for a double curve. The trisecant  $t$  lies on the first of these ruled surfaces and is double on the second.

Every line of  $\Omega$  lying in [4] meets the quartic\* in two points and therefore the octavic in four points. Hence we have a quintic ruled surface whose double curve consists of a conic and a quartic and whose bitangent developable consists of a quadric cone and a developable of the fourth class.

The quartic has a double point which lies on the conic, this being represented by the plane  $\omega$  of  $\Omega$  through  $t$ ; also the developable of the fourth class has a double tangent plane which touches the quadric cone, this being represented by the plane  $\rho$  of  $\Omega$  through  $t$ .

Through any point of  $C$  there pass one generator of the quartic ruled surface and two of the octavic; by considering the correspondence thus set up on  $C$  it is easily seen that the surfaces have two common generators. Hence the quartic curve and the conic have two further common points, while the developable of the fourth class and the quadric cone have two further common tangent planes. The common generators of the quartic and octavic ruled surfaces are  $\overline{pqr'}$  and  $\overline{pq'r}$ .

The prime section of  $\Omega$  on which  $C$  lies now contains the two chords  $\overline{pqr'}$  and  $\overline{pq'r}$ . Hence the curve of order 14 in which it meets  $F_2^7$  is made up as follows

$$C + p + \overline{pqr'} + \overline{pq'r} + 3t + c_3,$$

where  $c_3$  denotes a twisted cubic, which must be a double curve on the octavic ruled surface.

The section of this ruled surface by a solid is thus a curve of the eighth order lying on a quadric and meeting each generator in four points, the curve having nine double points†. This curve is rational; it represents the double quartic curve of the ruled surface and also the bitangent developable of the fourth class.

The section of the quartic ruled surface by a solid is a rational quartic curve with a double point on  $p$ .

**101. The surface with three double conics.** Suppose that we have a rational quintic ruled surface in a space [3] whose generators are represented by the points of a quintic  $C$  lying on the section of  $\Omega$  by a [4]; the axes  $p, q, r$  also lying on the quadric.

\* A solid through the line meets  $\Omega$  in a quadric with the line as generator and the quartic in a quartic curve, with a double point, lying on this quadric.

† Five arising from  $C$ , one from  $t$ , and three from  $c_3$ .

Then the ruled surface formed by the chords of  $C$  which lie on  $\Omega$  breaks up into three quartic ruled surfaces; each of these passes through  $C$  and has a double line, while any two of them have a common generator. A line on  $\Omega$  lying in [4] meets each of these surfaces in two points, and  $t$  lies on each surface.

Hence the quintic ruled surface in [3] has three double conics which all pass through one point, this being represented on  $\Omega$  by the plane  $\omega$  through  $t$ , while any two of the conics have a second intersection. The bitangent developable consists of three quadric cones all having a common tangent plane, this being represented on  $\Omega$  by the plane  $\rho$  through  $t$ , while any two of the cones have a second common tangent plane. The tritangent plane passes through the triple point.

The quadric  $\Omega$  now contains the three lines  $\overline{pqr'}$ ,  $\overline{pq'r}$ ,  $\overline{p'qr}$ ; these are, in fact, the common generators of the three pairs of quartic ruled surfaces.

The tangent planes at the different points of one of these lines, say  $\overline{pqr'}$ , to either of the quartic surfaces on which it is a generator, are the same; the two tangents of  $C$  at the ends of its chord  $\overline{pqr'}$  intersect and their plane touches both the quartic surfaces at all points of  $\overline{pqr'}$ . We can say that  $\overline{pqr'}$  is a torsal generator of both surfaces. The two quartic curves in which the surfaces are met by an arbitrary solid will touch at the intersection of the solid with  $\overline{pqr'}$ .

The curve of order 14 in which  $\Omega$  and  $F_2^7$  intersect is now

$$C + p + q + r + \overline{pqr'} + \overline{pq'r} + \overline{p'qr} + 3t.$$

The section by a solid of the composite ruled surface formed by the chords of  $C$  which lie on  $\Omega$  consists of three rational quartic curves lying on the same quadric; each curve has a double point and there are six points common to the three curves. Any two of the curves, besides having these six points in common, touch each other at another point.

102. *The surface with a double generator.* If  $C$  lies in a non-tangent prime  $S_4$  of  $\Omega$  and has a double point  $P$  its points represent the generators of a rational quintic ruled surface in [3]; the generators belong to a linear complex and there is a double generator  $G$  represented by  $P$ . Any line of  $S_4$  is met by five proper chords of  $C$ .

Any plane of  $\Omega$  meets  $S_4$  in a line  $l$  lying on  $\Omega$ ; this line is met by five proper chords of  $C$ , and the plane  $lP$  meets  $\Omega$  in  $l$  and another line passing through  $P$ . Hence we have a ruled surface whose double curve consists of the points of a double generator  $G$  together with a quintic, while the bitangent developable consists of the planes through a double generator  $G$  together with a developable of the fifth class.

The tangent prime of  $\Omega$  at  $P$  meets  $S_4$  in a [3] which meets  $C$  in four points at  $P$  and therefore in one other point  $Q$ . Thus  $G$  is met by one other generator  $g$ .

Through any point of  $C$  there pass three chords lying on  $\Omega$ ; except that through  $P$  we have the single chord  $PQ$  and through  $Q$  the chord  $QP$  and one other chord.

Take the plane  $\omega_1$  through  $PQ$ . A plane  $\rho$  which meets  $\omega_1$  in a line must meet  $S_4$  in a line intersecting  $PQ$ . This line is only met by three chords of  $C$  other than  $PQ$ , for if we project  $C$  from the line on to a plane of  $S_4$  we obtain a rational quintic with a triple point and three double points. The plane  $\omega_1$  represents the point  $Gg$ , so that any plane through this point meets the quintic double curve in only three other points.

Thus the quintic ruled surface has a double curve consisting of a double generator  $G$  and a quintic. These meet in a point which is a double point of the quintic, while through this point there passes one other generator.

The bitangent developable consists of a developable of the fifth class together with the planes through  $G$ . One of these planes is a double tangent plane of the developable; this is the plane  $Gg$  and is represented on  $\Omega$  by the plane  $\rho_1$  through  $PQ$ .

Consider again a rational quintic  $C$  in  $S_4$  with a double point  $P$ . Projecting from a general line of  $S_4$  on to a plane we obtain a quintic with six double points; hence the line is met by five chords. Projecting from a line meeting  $C$  we obtain a quartic with three double points, so that this line is met by two chords other than those which pass through its point of intersection with  $C$ . Projecting from a line through  $P$  we obtain a cubic with one double point, so that this line is met by one chord not passing through  $P$ . Hence the chords of  $C$  form a locus  $V_3^5$  of three dimensions and of the fifth order, on which  $C$  is a triple curve and  $P$  a quadruple point.

Further, if we take a chord of  $C$  passing through  $P$ , and project  $C$  on to a plane from a line meeting this chord, we see that the chord must be double on  $V_3^5$ . The chords of  $C$  through  $P$  form a cubic cone which is a double surface on  $V_3^5$ .

Consider now a quadric  $Q_3$  containing  $C$ . It meets the cubic cone just mentioned in  $C$  and also in one of its generators  $PQ$ . The chords of  $C$  which lie on  $Q_3$  form a ruled surface of order ten, the intersection of  $Q_3$  and  $V_3^5$ , on which  $C$  is a triple curve and  $PQ$  is a double generator. The section of this ruled surface by a solid is a curve of order ten with five triple points and one double point; it is the intersection of a quadric and a quintic surface, so that it meets every generator of the quadric in five points. If it is projected from a point of the quadric on to a plane we obtain a curve of order ten with five triple points, one double point and two quintuple points, which is therefore a rational curve.

The quintic double curve of the ruled surface is therefore also a rational curve. If we project it from a point of itself on to a plane we obtain a rational quartic with three double points. Hence there are two trisecants of the quintic curve passing through any point of it, and the ruled surface is formed by these trisecants.

Any line on  $\Omega$  meets the ruled surface formed by the chords of  $C$  lying on  $\Omega$  in five points, but if the line passes through  $P$  it only has one other intersection with the surface because  $P$  is a quadruple point thereon. Now any plane  $\rho$  on  $\Omega$  which passes through  $P$  meets  $S_4$  in such a line, so that any plane passing through the double generator  $G$  of the quintic ruled surface meets the double quintic curve in only one point not on  $G$ . Hence the quintic curve meets  $G$  in two points besides the double point; this shews at once that the quintic curve is rational.

Similarly, there are two planes of the developable, other than its double tangent plane, which pass through  $G$ .

103. If we project  $C$  from a point of itself on to an  $S_3$  we obtain a rational quartic with a double point; there are two quadric cones containing this quartic whose vertices are not on the curve, so that we expect to find two lines through any point of  $C$  which lie on  $V_3^5$  and are not chords of  $C$ . These lines will form a ruled surface with  $C$  for a double curve.

The existence of these lines is also seen by projecting a normal quintic in [5] from a point  $O$  on one of its chords. There are  $\infty^1$  secant planes\* through such a point, one for each involution on the curve which contains the extremities of the chord as a pair. Thus we have a finite number of secant planes joining  $O$  to any point of the curve, and in  $S_4$  a finite number of lines through any point of  $C$ . The chords of  $C$  meeting such a line form a cubic ruled surface, the projection of a quartic ruled surface in [5] from a point  $O$  on it.

The quadrics of  $S_4$  containing  $C$  are linearly dependent from five quadrics. It would thus seem that there is a quadric containing  $C$  and one of the lines, and so the ruled surface formed by the chords of  $C$  lying on the quadric breaks up into a cubic ruled surface passing through  $C$  and a septic ruled surface having  $C$  for a double curve. It would, however, be wrong to conclude from this that we have found a new species of quintic ruled surface belonging to the type II (B); for it can easily be shewn that a quadric in  $S_4$  which contains a cubic ruled surface is necessarily a point-cone.

*Surfaces with a directrix line which is not a generator*

104. Consider a ruled surface whose generators are represented by the points of a curve  $C$  lying in a tangent prime  $T$  of  $\Omega$ .  $T$  touches  $\Omega$  in a point  $O$ ;  $O$  represents a line  $R$  which is a directrix of the ruled surface.

\* Cf. § 90.

If  $C$  is of the type III (A) it meets every plane  $\omega$  through  $O$  in four points and every plane  $\rho$  through  $O$  in one point. An arbitrary plane  $\rho$  meets  $T$  in a line and the plane  $\omega$  through this line is the plane joining it to  $O$ . This contains four points and therefore six chords of  $C$ , so that the double curve of the ruled surface is the line  $R$  counted six times. There are four generators passing through every point of  $R$  and there is one generator lying in every plane through  $R$ .

An arbitrary plane  $\omega$  meets  $T$  in a line and the plane  $\rho$  through this line joins it to  $O$ . This contains only one point of  $C$ ; there will be six chords of  $C$  meeting the line so that in general the bitangent developable is non-degenerate and of the sixth class.

Similarly in III (B) we have a surface whose double curve is non-degenerate and of the sixth order, while the bitangent developable consists of the planes through  $R$  counted six times. Any plane through  $R$  contains four generators, while through any point of  $R$  there passes one generator.

Remembering that  $C$  has a trisecant we see that in III (A) there is a tritangent plane and in III (B) there is a triple point. Also the bitangent developable in III (A) and the double curve in III (B) are seen to be elliptic precisely as in § 96.

**105.** We have seen in § 99 that we can have a quadric point-cone containing  $C$  and an axis; we may then regard this cone as a section of  $\Omega$  by  $T$ .

Thus in III (A) the bitangent developable may break up into a quadric cone and a developable of the fourth class, this latter having a double tangent plane which touches the former, and there being also two other common tangent planes. The quadrisecant planes of  $C$  meeting the axis are the  $\omega$ -planes of the cone.

Also in III (B) the double curve may break up into a conic and a quartic; the quartic has a double point lying on the conic and the curves have also two other intersections. The quadrisecant planes of  $C$  meeting the axis are the  $\rho$ -planes of the cone.

**106.** We shall now shew how a rational quintic curve  $C$  can be found, in [4], so that it lies, together with three axes, on a quadric point-cone.

Take two planes  $\pi, \pi'$  in a space  $S_4$  intersecting in a point  $O$ . Then taking three arbitrary points  $A, B, C$  in  $\pi$  and three arbitrary points  $A', B', C'$  in  $\pi'$  there is thus defined a collineation between the planes; the points  $A, B, C, O$  of  $\pi$  corresponding to the points  $A', B', C', O$  of  $\pi'$ . If a conic in  $\pi$  is drawn through  $O$  to have  $ABC$  as a self-conjugate triangle (there is a pencil of such conics), then we have correspondingly in  $\pi'$  a conic through  $O$  with  $A'B'C'$  as a self-conjugate triangle.



Through any point of  $S_4$  there passes a plane pencil of lines incident to both  $\pi$  and  $\pi'$ ; the plane of this pencil is the intersection of the solids which join the point to  $\pi$  and  $\pi'$  and meets these planes in two lines passing through  $O$ . These lines will not, in general, correspond to one another in the collineation.

The three lines  $AA'$ ,  $BB'$ ,  $CC'$  have a common transversal meeting them in  $P$ ,  $Q$ ,  $R$ . Now take  $S$ , any point of the line  $PQR$ ; the plane of the pencil of lines through  $S$  will meet  $\pi$  in a line  $l$  through  $O$  and  $\pi'$  in a line  $l'$  through  $O$ . The three planes  $\pi$ ,  $\pi'$ ,  $OPQR$  determine a quadric point-cone with vertex  $O$ , and the four planes  $OAA'P$ ,  $OBB'Q$ ,  $OCC'R$ ,  $OU'S$  are planes of the opposite system of this cone. Hence the two pencils  $O\{ABC\}$  and  $O\{A'B'C'l\}$  are homographic, so that  $l$  and  $l'$  are corresponding lines in the collineation between  $\pi$  and  $\pi'$ .

The pencil of lines through  $S$  incident to  $\pi$  and  $\pi'$  meets  $l$  and  $l'$  in two ranges in perspective. For a general position of  $S$  on the line  $PQR$  there will be two points of  $l$  which are in perspective from  $S$  with their corresponding points on  $l'$ , and these include  $O$ . Hence there is one line, other than  $SO$ , through  $S$  which joins a pair of corresponding points of  $\pi$  and  $\pi'$ .

The lines  $BC$ ,  $B'C'$ ,  $QR$  lie in the solid determined by  $BB'Q$  and  $CC'R$ ; this solid contains  $P$  on the line  $QR$ . Hence there is a line  $PDD'$  meeting  $BC$  in  $D$  and  $B'C'$  in  $D'$ . The plane  $PDD'AA'$  meets  $\pi$  in the line  $AD$  and  $\pi'$  in the line  $A'D'$ , so that  $AD$ ,  $A'D'$  must intersect in  $O$ . Then the lines  $OAD$ ,  $OA'D'$  are corresponding lines and the points  $D$ ,  $D'$  are corresponding points. We have therefore two lines  $PAA'$  and  $PDD'$  other than  $PO$  which join  $P$  to a pair of corresponding points of  $\pi$  and  $\pi'$ ; there must then be an infinite number of such lines through  $P$ .

Through each of the points  $P$ ,  $Q$ ,  $R$  there passes a pencil of lines incident to  $\pi$  and  $\pi'$  in corresponding points; every line through one of these three points which meets both  $\pi$  and  $\pi'$  does so in a pair of corresponding points.

If  $X$  is the harmonic conjugate of  $O$  in regard to  $A$  and  $D$ , any conic through  $O$  having  $ABC$  as a self-conjugate triangle must pass through  $X$ . Similarly we have  $X'$  in  $\pi'$ , and  $XX'$  passes through  $P$ . We also have  $QYY'$  and  $RZZ'$  in the same way.

Returning to the point  $S$ , let  $SUU'$  be the line through it which joins corresponding points  $U$  and  $U'$  of  $\pi$  and  $\pi'$ . There is a definite conic passing through  $O$  and  $U$  for which  $ABC$  is a self-conjugate triangle; to this corresponds the conic through  $O$  and  $U'$  for which  $A'B'C'$  is a self-conjugate triangle. The points of these two conics are in (1, 1) correspondence with a united point  $O$ ; hence the lines joining corresponding points of the two conics form a cubic ruled surface in  $S_4$ . The four lines  $XX'$ ,  $YY'$ ,  $ZZ'$ ,  $UU'$  are generators of this surface, so that  $P$ ,  $Q$ ,  $R$ ,  $S$  are on the surface. Hence  $PQRS$  is its directrix line.

Through eight points of general position on this surface there passes a rational quintic  $C$  with the directrix for its trisecant\*.

Take then points  $a, b, c, d$  on the conic  $OXYZ$  such that  $bc, ad$  pass through  $A$ ;  $ca, bd$  pass through  $B$ ;  $ab, cd$  pass through  $C$ . Similarly, take points  $e, f, g, h$  on the conic  $OU'X'Y'Z'$  such that  $fg, eh$  pass through  $A'$ ;  $ge, fh$  pass through  $B'$ ;  $ef, gh$  pass through  $C'$ . Then there is a rational quintic passing through  $a, b, c, d, e, f, g, h$  with the directrix for its trisecant.

The lines  $PAA', QBB', RCC'$  must lie on the  $V_3$ 's formed by the chords of the curve, as each meets it in a triple point and two double points. Hence these are three axes of the curve, which must meet its trisecant in  $P, Q, R$ .

There is a quadric point-cone whose vertex is  $O$  and which contains the planes  $OAA'P, OBB'Q, OCC'R$ , as already mentioned above. The planes  $\pi, \pi', OPQR$  belong to the opposite system. The cone therefore meets the quintic curve in at least eleven points, and so contains it entirely. It is the cone projecting the cubic ruled surface from  $O$ .

Hence we have constructed in  $S_4$  a rational quintic which lies, together with three of its axes and trisecant, on a quadric point-cone.

This shews that for surfaces of the type III (A) we can have a bitangent developable consisting of three quadric cones; all the cones having a common tangent plane, while any two have a second common tangent plane.

Also in surfaces of the type III (B) we can have a double curve consisting of three conics; all the conics have a point in common, while any two of them have a second intersection.

107. In the type III (C) the curve  $C$  meets every plane  $\omega$  through  $O$  in three points, every plane  $\rho$  through  $O$  in two points, and has a double point,  $O$  lying on a chord of  $C$ .

An arbitrary plane  $\rho$  meets  $T$  in a line; the plane  $\omega$  through this line is the plane joining it to  $O$ , which contains three points and therefore three chords of  $C$ . The plane joining the line to the double point  $P$  meets  $\Omega$  in a second line through  $P$ , and there will be two further chords of  $C$  meeting the line.

Hence the double curve consists of the directrix  $R$  counted three times, a double generator  $G$ , and a conic.

$R$  and  $G$  intersect.  $R$  and the conic also intersect; their point of intersection is represented on  $\Omega$  by the  $\omega$ -plane which contains the chord of  $C$  passing through  $O$ . The conic meets  $G$  also.

\* Marletta, *loc. cit.* p. 102. No five of the points must be on the same conic and no two on the same generator.

The tangent prime of  $\Omega$  at  $P$  meets  $T$  in a solid passing through  $O$  and meeting  $C$  in four points at  $P$ . Thus it meets the curve in one further point  $Q$ , and the plane  $OPQ$  is a  $\omega$ -plane. Hence  $G$  is met by one other generator  $g$  only, and this passes through the point  $RG$ .

Through any point of  $R$  there pass three generators.

An arbitrary plane  $\omega$  meets  $T$  in a line; the plane  $\rho$  through this line is the plane joining it to  $O$ , which contains two points and therefore one chord of  $C$ .

Hence the bitangent developable consists of the two pencils of planes through  $R$  and  $G$ , together with a developable of the fourth class.

There is a plane of this developable passing through  $R$ ; it is represented on  $\Omega$  by the  $\rho$ -plane which contains the chord of  $C$  passing through  $O$ . The developable has the plane  $Gg$  as a double tangent plane; this is represented by the plane  $\rho_1$  through  $PQ$ . Any plane  $\omega$  which meets  $\rho_1$  in a line meets  $T$  in a line intersecting  $PQ$ ; this line is met by three chords of  $C$  other than  $PQ^*$ , but one of these lies in the  $\rho$ -plane which joins the line to  $O$ . Correspondingly, through any point of the plane  $Gg$  there pass only two planes of the developable besides  $Gg$ .

**108.** The ruled surface formed by the chords of  $C$  which lie on a quadric containing it is, in general, of order 10 with  $C$  for a triple curve, and has a double generator\*. But in surfaces of the type III (C) this surface will clearly break up into two distinct parts, one formed by those chords which lie in  $\omega$ -planes through  $O$  and the other formed by those chords which lie in  $\rho$ -planes through  $O$ .

The chords of  $C$  which lie in the  $\rho$ -planes through  $O$  join the pairs of points of an involution on  $C$  and therefore form a cubic ruled surface†. There is a generator of this surface passing through  $P$ , and the plane  $\omega$  through this generator represents the point of intersection of  $G$  with the double conic. Thus the ruled surface formed by the chords of  $C$  which lie on  $\Omega$  breaks up into a cubic ruled surface and a ruled surface of the seventh order, this latter having  $C$  for a double curve and having also a double generator‡. The section of this composite ruled surface by a solid consists of a twisted cubic and a septic having six double points. The two curves lie on a quadric and form its curve of intersection with a surface of the fifth order; the cubic meets all generators of one system in one point and all of the other system in two points, so that the septic meets all generators of one system in four points and all of the other system in three points. If then the septic is projected on to a plane we obtain a curve with six

\* Cf. § 102.

† The pair of points of  $C$  on its two branches at  $P$  is a pair of the involution, so that we have a cubic ruled surface—the projection of a quartic ruled surface in [5] from a point of itself.

‡ Cf. § 103.

double points, one triple point and one quadruple point, and this curve is of genus  $15 - 6 - 3 - 6 = 0$ , so that it is rational.

The cubic passes through five of the double points of the septic and meets it in one other point. This other point is on the chord of  $C$  which passes through  $O$ ; this chord is a common generator of the two ruled surfaces.

**109.** In surfaces of the type III (D) the double curve consists of a directrix  $R$ , a double generator  $G$  and a quartic. This quartic is rational, having a double point on  $G$ , this being the point of intersection of  $G$  with the only other generator which meets it. The quartic meets  $R$ . The bitangent developable consists of the planes through  $R$  counted three times, the planes through  $G$ , and a quadric cone; one tangent plane of this cone passing through  $R$  and another through  $G$ .

Through each point of  $R$  there pass two generators; there is one point of  $R$  at which the plane of the two generators contains  $R$ , it is represented on  $\Omega$  by the  $\omega$ -plane containing the chord of  $C$  which passes through  $O$ , and is the intersection of  $R$  with the double conic. Each plane through  $R$  contains three generators.

**110.** In the type III (E)  $C$  meets every plane  $\omega$  through  $O$  in three points and every plane  $\rho$  through  $O$  in two points, while two chords of  $C$  pass through  $O$ . We have a rational quintic ruled surface in [3] with a directrix line  $R$ ; through any point of  $R$  there pass three generators, while any plane through  $R$  contains two generators. There are two and only two planes through  $R$  such that the two generators in either plane intersect in a point of  $R$ .

An arbitrary plane  $\rho$  meets  $T$  in a line; the plane  $\omega$  through this line is the plane joining it to  $O$ , which contains three points and therefore three chords of  $C$ . There are three other chords of  $C$  meeting the line, so that the double curve of the ruled surface consists of the directrix  $R$  counted three times together with a twisted cubic.  $R$  is a chord of the cubic. The two points in which  $R$  meets the cubic have, in fact, already been noticed; they are the two points of  $R$  in which pairs of generators meet and at the same time lie in a plane with  $R$ . The tangents of the cubic at these two points lie in the planes of the pairs of generators.

An arbitrary plane  $\omega$  meets  $T$  in a line, and the plane  $\rho$  through this line is the plane joining it to  $O$ , which contains two points and therefore one chord of  $C$ . There will be five other chords of  $C$  meeting the line. Hence the bitangent developable consists of the planes through  $R$  together with a developable of the fifth class. There are two planes of this developable passing through  $R$  represented by the  $\rho$ -planes of  $\Omega$  containing the two chords of  $C$  which pass through  $O$ ; it has also a tritangent plane, this being represented on  $\Omega$  by the plane  $\rho$  through the trisecant of  $C$ .

111. In the type III (F) we have a rational quintic ruled surface in [3] with a directrix line  $R$ ; through any point of  $R$  there pass two generators, while any plane through  $R$  contains three generators. The double curve consists of  $R$  and a quintic; this quintic has a triple point and meets  $R$  in two points. The bitangent developable consists of the planes through  $R$ , counted three times, together with a developable of the third class, two planes of this latter developable passing through  $R$ .

112. Consider now the genus of this quintic curve; we know that it must be rational, having a triple point.

The trisecant  $t$  of  $C$  is joined to  $O$  by a  $\rho$ -plane; it therefore meets every  $\omega$ -plane through  $O$  in one point. The chords of  $C$  form the locus  $V_3^6$  with the double surface  $F_2^7$ ; a  $\omega$ -plane through  $O$  meets  $F_2^7$  in two points of  $C$ , one other point on the chord joining these two points, three points at the intersection of the plane with  $t$ , and also in  $O$ ; while a  $\rho$ -plane through  $O$  meets  $F_2^7$  in three points of  $C$ , three other points on the chords joining the pairs of these, and also in  $O^*$ .

The intersection of  $F_2^7$  with the quadric point-cone, vertex  $O$ , on which  $C$  lies, consists of  $C$ ,  $t$  counted three times, and a sextic curve with a double point at  $O$  and meeting every chord of  $C$  in one point†. It is clear that this sextic meets every  $\omega$ -plane through  $O$  in one point and every  $\rho$ -plane through  $O$  in three points other than  $O$ ; it is thus a rational curve.

Here again the ruled surface formed by the chords of  $C$  which lie on  $\Omega$  breaks up. The  $\omega$ -planes through  $O$  meet  $C$  in pairs of an involution, so that the chords in these planes form a quartic ruled surface with a double point at  $O$ , the projection of a normal quartic ruled surface in [5] from a point. The section of this ruled surface by any solid lying in  $T$  is a rational quartic lying on a quadric, meeting all generators of one system in three points and all of the other system in one point.

The chords of  $C$  which lie in the  $\rho$ -planes through  $O$  form a ruled surface of order 8 with  $C$  as a double curve and  $t$  as a triple line. By considering the intersections of the planes of  $\Omega$  through  $O$  with the surface  $F_2^7$  it is easily seen that there is no other multiple curve on this ruled surface. The section by a solid gives an octavic curve having five double points and one triple point; it lies on a quadric, meeting all the generators of one system in three points and all of the other system in five points. Hence the genus of this curve is

$$21 - 5 - 3 - 3 - 10 = 0,$$

so that it is a rational curve. Thus the quintic double curve of the ruled surface in [3] is a rational curve.

The chords of  $C$  which lie in the  $\omega$ -planes through  $O$  correspond to the points of the line  $R$  regarded as part of the double curve and to the planes

\* Cf. § 95.

† Cf. § 96.

of the developable of the third class regarded as part of the bitangent developable; the chords of  $C$  which lie in  $\rho$ -planes through  $O$  correspond to the points of the quintic curve and to the triple pencil of planes through  $R$ ; this last must be regarded as a rational family of planes.

The ruled surface is formed by the chords of the quintic curve which meet  $R$ ; clearly two such chords pass through any point of the curve and three lie in any plane through  $R$ . We thus take a rational quintic with a triple point and one of its chords, and we at once obtain a surface of this type.

There are similar results for the surface of the type III (E).

*Surfaces with a directrix line which is also a generator*

113. In surfaces belonging to the type IV we have a directrix line  $R$  which is also a generator, this being represented by the point  $O$  of  $\Omega$ ;  $C$  lies in the tangent prime of  $\Omega$  at  $O$  and passes through  $O$ .

In IV (A)  $C$  meets every  $\omega$ -plane through  $O$  in three points and every  $\rho$ -plane through  $O$  in one point other than  $O$ . Through every point of  $R$  there pass three generators other than  $R$ , while every plane through  $R$  contains one generator other than  $R$ .

An arbitrary plane  $\rho$  meets  $T$  in a line, and the plane  $\omega$  through this line is the plane joining it to  $O$ . This contains four points and therefore six chords of  $C$ , and no other chords meet the line. Hence the double curve is the line  $R$  counted six times. An arbitrary plane  $\omega$  meets  $T$  in a line, and the plane  $\rho$  through this line joins it to  $O$ . This contains two points and therefore one chord of  $C$ ; there will be five other chords meeting the line. Hence the bitangent developable consists of the planes through  $R$  together with a developable of the fifth class. Two planes of this latter pass through  $R$ , and it has a tritangent plane represented on  $\Omega$  by the plane  $\rho$  passing through the trisecant of  $C$ . The  $\omega$ -plane through the tangent of  $C$  at  $O$  meets  $C$  in two other points; the two  $\rho$ -planes containing the chords of  $C$  which join these points to  $O$  represent the two planes of the quintic developable which pass through  $R$ .

In IV (B) we have another type of surface;  $R$  is both a directrix and a generator, through any point of  $R$  there passes one generator other than  $R$ , while any plane through  $R$  contains three generators other than  $R$ . The double curve consists of  $R$  and a quintic with a triple point,  $R$  being a chord of the quintic; the bitangent developable consists of the planes through  $R$  counted six times.

114. In IV (C)  $C$  meets every plane of  $\Omega$  through  $O$  in two points other than  $O$  and has a double point  $P$ ;  $P$  will represent a double generator  $G$  of the ruled surface. Through every point of  $R$  there pass two generators other than  $R$ , while every plane through  $R$  contains two generators other than  $R$ .

Any plane of  $\Omega$  meets  $T$  in a line; the plane of the opposite system through this line passes through  $O$ , containing three points and therefore three chords of  $C$ . The plane joining the line to  $P$  meets  $\Omega$  in a second line through  $P$ , while there are two other chords of  $C$  meeting the line. The double curve therefore consists of  $G$ ,  $R$  counted three times, and a conic.  $R$  and  $G$  intersect. The  $\rho$ -plane through the tangent of  $C$  at  $O$  meets  $C$  in another point, and the  $\omega$ -plane, which contains the line joining this point to  $O$ , represents a point of intersection of  $R$  and the conic.  $G$  and the conic also intersect. Similarly the bitangent developable consists of the planes through  $G$ , the planes through  $R$  counted three times, and the tangent planes of a quadric cone, one tangent plane of this cone passing through  $R$  and another through  $G$ .

The ruled surface formed by the chords of  $C$  which lie on  $\Omega$  here breaks up into a quartic cone with vertex  $O$  passing through  $C$  and having the double generator  $OP$ , together with two cubic ruled surfaces both passing through  $C$ .

115. In IV (D) the trisecant of  $C$  passes through  $O$ ,  $C$  meeting every plane of  $\Omega$  through  $O$  in two points other than  $O$ . Again, we have a surface for which the line  $R$  is both a directrix and a generator; every plane through  $R$  contains two generators other than  $R$ , while through every point of  $R$  there pass two generators other than  $R$ . The difference between this surface and one of the type IV (C) lies in the fact that there is a plane through  $R$  which contains two generators meeting in a point of  $R$ , while in the former type of surface we only have the plane through  $R$  and a double generator.

Any plane of  $\Omega$  meets  $T$  in a line; the plane of the opposite system through this line joins it to  $O$ , containing three points and therefore three chords of  $C$ . There are three other chords of  $C$  meeting the line. Hence the double curve consists of the line  $R$  counted three times together with a twisted cubic.  $R$  is a chord of the cubic. One of the common points of  $R$  and the cubic is represented on  $\Omega$  by the  $\omega$ -plane through the trisecant of  $C$ . The other is represented by the  $\omega$ -plane which contains that chord of  $C$  which joins  $O$  to the remaining intersection of  $C$  with the  $\rho$ -plane containing its tangent at  $O$ . Similarly the bitangent developable consists of the planes through  $R$  counted three times together with a developable of the third class, two of whose planes pass through  $R$ .

116. If  $C$  lies in a tangent prime  $T$  of  $\Omega$  and has a double point at  $O$  we have a rational quintic ruled surface in [3] with a directrix line  $R$  which is also a double generator.

For the type V (A)  $C$  meets every  $\omega$ -plane through  $O$  in two points and every  $\rho$ -plane through  $O$  in one point other than  $O$ . An arbitrary plane  $\rho$  meets  $T$  in a line; the plane  $\omega$  through this line joins it to  $O$ , and meets

$C$  in the double point  $O$  and two points  $A$  and  $B$ . The chords  $OA$ ,  $OB$ ,  $AB$  are the only chords of  $C$  meeting the line, since  $OA$  and  $OB$  are double on the locus  $V_3^5$  formed by the chords of  $C^*$ . Hence the double curve is the line  $R$  counted six times. An arbitrary plane  $\omega$  meets  $T$  in a line; the plane  $\rho$  through this line joins it to  $O$  and contains one other point of  $C$ , so that there will be three chords of  $C$  meeting the line which do not lie in this  $\rho$ -plane. Hence the bitangent developable consists of  $R$  counted three times and a developable of the third class. There are two planes of this developable passing through  $R$ . The  $\omega$ -plane through either tangent of  $C$  at  $O$  meets  $C$  in one other point; the  $\rho$ -plane through the chord of  $C$  joining this point to  $O$  represents a plane of the developable passing through  $R$ .

Through any point of  $R$  there pass two generators other than  $R$ , while any plane through  $R$  contains one generator other than  $R$ .

For a surface of the type V (B) we have a directrix  $R$  which is also a double generator; through any point of  $R$  there passes one generator other than  $R$ , while any plane through  $R$  contains two generators other than  $R$ . The double curve consists of the triple line  $R$  together with a twisted cubic having  $R$  for a chord; the bitangent developable consists of the planes through  $R$  counted six times.

The ruled surface, formed by the chords of  $C$  which lie on  $\Omega$ , breaks up into the cubic cone which projects  $C$  from  $O$ , counted twice, together with a quartic ruled surface passing through  $C$ .

### *Surfaces whose generators belong to a linear congruence*

117. Suppose now that  $C$  lies on the quadric  $Q$  in which  $\Omega$  is met by a solid  $S_3$ , the two tangent primes of  $\Omega$  through  $S_3$  touching it in  $O$  and  $O'$ . Then  $O$  and  $O'$  represent lines  $R$  and  $R'$ , in [3], which are both directrices of the surface.

If  $C$  belongs to the type VI (A) it meets all generators of  $Q$  of one system in four points and all of the other system in one point. We can suppose that the generators of the first system lie in the  $\omega$ -planes through  $O$  and the  $\rho$ -planes through  $O'$ ; those of the other system lying in the  $\omega$ -planes through  $O'$  and the  $\rho$ -planes through  $O$ . Then through every point of  $R$  there pass four generators of the ruled surface which lie in a plane through  $R'$ , while conversely every plane through  $R'$  contains four generators which meet in a point of  $R$ .

The rational quintic in  $S_3$  has six apparent double points. An arbitrary plane  $\rho$  meets  $S_3$  in a point of  $Q$ ; the six chords of  $C$  through this point all coincide with the generator of  $Q$  which is quadriseccant to  $C$ , the projection of  $C$  from this point on to a plane in  $S_3$  being a quintic with a quadruple point. The plane  $\omega$  through this generator passes through  $O$ ,



so that the double curve of the surface consists of the line  $R$  counted six times. Similarly the bitangent developable consists of the line  $R'$  counted six times.

118. If  $C$  belongs to the type VI (B) it meets all generators of  $Q$  of one system in three points and all of the other system in two points, having two double points. These represent two double generators  $G$  and  $H$  of the surface. Any point of  $R$  is the intersection of three generators lying in a plane through  $R'$ , while conversely any plane through  $R'$  contains three generators meeting in a point of  $R$ . Through any point of  $R'$  there pass two generators lying in a plane through  $R$ , while any plane through  $R$  contains two generators meeting in a point of  $R'$ .

An arbitrary plane  $\rho$  meets  $S_3$  in a point of  $Q$ ; the six chords of  $C$  through this point consist of the lines to the two double points, the generator of  $Q$  meeting  $C$  in two points, together with the trisecant generator counted three times. Hence the double curve consists of  $G$ ,  $H$ ,  $R'$  together with  $R$  counted three times. The bitangent developable consists of the planes through  $G$ ,  $H$  and  $R$  together with those through  $R'$  counted three times.

119. Now suppose that  $C$  lies on the quadric cone in which  $\Omega$  is met by a solid touching it at a point  $V$ .

In the type VII (A)  $C$  passes through  $V$  and meets every generator of the cone in two points other than  $V$ . The surface has a directrix line  $R$  which is also a generator. Since each generator of the cone is the intersection of two planes of  $\Omega$ , of opposite systems, both lying in the tangent prime at  $V$ , through each point of  $R$  there pass two other generators which lie in a plane with  $R$ , while each plane through  $R$  contains two other generators meeting in a point of  $R$ .  $C$  has two double points, so that the surface has two double generators  $G$  and  $H$ .

Any plane of  $\Omega$  meets  $S_3$  in a point of the cone. The projection of  $C$  from this point on to a plane of  $S_3$  gives a rational quintic with two double points, and a triple point at which two branches touch each other. This latter is equivalent to four double points, being formed by the union of two double points and a tacnode. The six chords of  $C$  which can be drawn from any point on the cone consist of the generator of the cone, counted four times, together with the lines to the two double points.

The double curve of the surface consists of  $G$  and  $H$  with  $R$  counted four times, while the bitangent developable consists of  $G$  and  $H$  considered as pencils of planes together with the planes through  $R$  counted four times.

This surface is really a degeneration of the type VI (B) when the two directrices  $R$  and  $R'$  of that type coincide. The plane sections will be rational quintic curves with double points on  $G$  and  $H$  and triple points (at which two branches have a common tangent) on  $R$ .

120. In the type VII (B)  $C$  has a triple point at  $V$  and meets every generator of the cone in one point other than  $V$ . The surface has a directrix line  $R$  which is also a triple generator; through any point of  $R$  there passes one other generator, while any plane through  $R$  contains one other generator. Clearly the double curve consists of the points of  $R$  counted six times, the bitangent developable of the planes through  $R$  counted six times.

121. We have now found twenty-four different kinds of rational quintic ruled surfaces in [3]; these are exhibited in tabular form on p. 304.

## SECTION II

### RATIONAL QUINTIC RULED SURFACES CONSIDERED AS PROJECTIONS OF NORMAL SURFACES IN HIGHER SPACE

#### *The general surface in [6]*

122. The rational quintic ruled surface is normal in [6]; there are two distinct kinds of surfaces, one with a directrix line  $\lambda$ , and the most general one with a directrix conic  $\Gamma^*$ .

The rational quintic ruled surfaces  $f$  of ordinary space  $\Sigma$  can all be obtained by projection from the normal surfaces  $F$  in [6]. The projection will be from a plane  $\omega$  which must not meet  $F$ ; the solids joining  $\omega$  to the points of  $F$  meet  $\Sigma$  in the points of  $f$ , while the [4]'s joining  $\omega$  to the generators of  $F$  meet  $\Sigma$  in the generators of  $f$ .

Let us now consider the general surface  $F$  with a directrix conic  $\Gamma$ . A prime through  $\omega$  meets  $F$  in a rational normal quintic, six of whose chords meet  $\omega$ ; so that there are six solids in this prime which pass through  $\omega$  and meet  $F$  in two points. Projecting on to  $\Sigma$  we see that in any plane there are six points in which two generators of  $f$  intersect, so that the double curve is a sextic.

123. Consider now the five-dimensional locus  $M_5$  formed by the chords of  $F$ .

This locus  $M_5$  contains any solid  $K$  which contains a directrix cubic  $\Delta$  of  $F$ ; for through any point of  $K$  there passes a chord of  $\Delta$ . Also  $M_5$  contains any solid  $K'$  which contains a pair of generators  $g, g'$  of  $F$ ; for through any point of  $K'$  there passes a transversal of  $g$  and  $g'$ .

Conversely, through any point of  $M_5$  there passes a chord of  $F$ ; this meets  $F$  in two points through which† there passes a directrix cubic  $\Delta$ , while the points themselves lie on two generators  $g$  and  $g'$ . Hence every

\* § 43.

† § 45.

point of  $F$  lies in a solid  $K$  and also in a solid  $K'$ , so that  $M_5$  can be generated either by the  $\infty^2$  solids  $K$  or by the  $\infty^2$  solids  $K'$ .

If any solid  $K$  is taken, a prime through it will meet  $F$  in the directrix cubic of  $K$  and in two generators; these two generators lie in a solid  $K'$  which meets the plane of  $\Gamma$  in the line joining the points where the two generators meet  $\Gamma$ . Taking three different solids  $K$ , each of them is the base of a doubly-infinite system of primes; and we can establish a projectivity between the primes of these three systems, two primes of different systems corresponding when they join the respective solids  $K$  to the same line in the plane of  $\Gamma$ . Then corresponding primes of the three systems meet in a solid  $K'$ , and all the solids  $K'$  are given in this way.

Hence  $M_5$ , being generated by the solids  $K'$ , is generated by three doubly-infinite systems of primes projectively related to each other. This proves that  $M_5$  is a *cubic* primal  $M_5^3$ ; for if we take the section by an arbitrary solid we obtain a surface generated by three projectively related "stars" of planes, the vertices of the stars being the points in which the arbitrary solid is met by the three solids  $K$ . We have thus the well-known generation of the *cubic* surface.

When the cubic surface in [3] is generated by three projective stars of planes there are six sets of three corresponding planes of the stars which have a line in common instead of a point only; these lines lie on the cubic surface and form half of a double-six. Hence, given an arbitrary solid in the [6] containing  $F$ , there are six solids  $K'$  meeting it in lines. In particular, we may take this arbitrary solid to contain the plane  $\omega$  from which we are projecting  $F$  on to  $\Sigma$ . Then, through a given point of  $\Sigma$  there pass six planes, each of which contains a pair of generators of  $f$ ; so that the bitangent developable of  $f$  is of the sixth class. The projected surface  $f$  has one tritangent plane; this being the intersection of  $\Sigma$  with the prime containing  $\omega$  and  $\Gamma$ .

The chords of  $F$  meet  $\omega$  in the points of a cubic curve  $c_3$ , this being the intersection of  $\omega$  with  $M_5^3$ . Since no two chords of  $F$  can intersect (except on  $F$  itself or at a point in the plane of  $\Gamma$ ) this curve has no double point and is therefore elliptic. Hence the double curve and bitangent developable of  $f$  are both elliptic; the points of the double curve and the planes of the bitangent developable both being in (1, 1) correspondence with the points of  $c_3$ .

124. Algebraically, suppose that the surface is generated by a (1, 1) correspondence between its directrix conic  $\Gamma$  in the plane  $x_0 = x_1 = x_2 = x_3 = 0$  and one of its directrix cubics  $\Delta$  in the solid  $x_4 = x_5 = x_6 = 0$ . Then we may take the coordinates of corresponding points to be

$$(0, 0, 0, 0, \theta^2, \theta, 1) \text{ and } (\theta^3, \theta^2, \theta, 1, 0, 0, 0),$$

so that any point of  $F$  has coordinates

$$(\theta^3, \theta^2, \theta, 1, \lambda\theta^2, \lambda\theta, \lambda),$$

and the equations of the surface are

$$\frac{x_0}{x_1} = \frac{x_1}{x_2} = \frac{x_2}{x_3} = \frac{x_4}{x_5} = \frac{x_5}{x_6},$$

the generators being given by  $\theta = \text{const.}$

The chords of  $F$  are  $\infty^4$  in aggregate and form a locus  $M_5$  whose equation can at once be written down. For, taking any two points

$$(\theta^3, \theta^2, \theta, 1, \lambda\theta^2, \lambda\theta, \lambda) \text{ and } (\phi^3, \phi^2, \phi, 1, \mu\phi^2, \mu\phi, \mu)$$

on  $F$ , the coordinates of a point on the chord joining them are

$$(\theta^3 + \kappa\phi^3, \theta^2 + \kappa\phi^2, \theta + \kappa\phi, 1 + \kappa, \lambda\theta^2 + \kappa\mu\phi^2, \lambda\theta + \kappa\mu\phi, \lambda + \kappa\mu),$$

and these may be taken as the coordinates of any point of  $M_5$ . The coordinates are thus expressed in terms of five parameters. They satisfy the relations

$$x_0 - (\theta + \phi)x_1 + \theta\phi x_2 = 0,$$

$$x_1 - (\theta + \phi)x_2 + \theta\phi x_3 = 0,$$

$$x_4 - (\theta + \phi)x_5 + \theta\phi x_6 = 0,$$

so that the equation of  $M_5$  is

$$\begin{vmatrix} x_0 & x_4 & x_2 \\ x_1 & x_2 & x_3 \\ x_4 & x_5 & x_6 \end{vmatrix} = 0,$$

a primal  $M_5^3$  of the third order.

Since the equation of  $M_5^3$  is given by equating a determinant to zero it follows at once that it can be generated by systems of spaces in two different ways; it contains two doubly-infinite systems of solids.

The solid whose equations are

$$ax_0 + \beta x_1 + \gamma x_2 = 0,$$

$$ax_1 + \beta x_2 + \gamma x_3 = 0,$$

$$ax_4 + \beta x_5 + \gamma x_6 = 0,$$

lies on  $M_5^3$  for all values of  $a : \beta : \gamma$ . This solid meets  $F$  in points for which

$$a\theta^2 + \beta\theta + \gamma = 0,$$

i.e. in two generators.

Conversely, the solid containing the two generators given by  $\theta = \theta_1$  and  $\theta = \theta_2$  is determined by the four points

$$(\theta_1^3, \theta_1^2, \theta_1, 1, 0, 0, 0), \quad (0, 0, 0, 0, \theta_1^2, \theta_1, 1),$$

$$(\theta_2^3, \theta_2^2, \theta_2, 1, 0, 0, 0), \quad (0, 0, 0, 0, \theta_2^2, \theta_2, 1),$$

so that its equations are

$$x_0 - (\theta_1 + \theta_2)x_1 + \theta_1\theta_2 x_2 = 0,$$

$$x_1 - (\theta_1 + \theta_2)x_2 + \theta_1\theta_2 x_3 = 0,$$

$$x_4 - (\theta_1 + \theta_2)x_5 + \theta_1\theta_2 x_6 = 0,$$

all other primes through the solid being linear combinations of these three. Thus the solid lies on  $M_5^3$ ;  $M_5^3$  can be generated by the solids  $K'$ .

We also have the conjugate generation of  $M_5^3$ ; the solid whose equations are

$$ax_0 + bx_1 + cx_4 = 0,$$

$$ax_1 + bx_2 + cx_5 = 0,$$

$$ax_2 + bx_3 + cx_6 = 0,$$

lying on  $M_5^3$  for all values of  $a : b : c$ . This solid meets  $F$  in points for which  $\theta$  and  $\lambda$  are connected by the relation

$$a\theta + b + c\lambda = 0.$$

Since this is linear in  $\lambda$  it represents a directrix curve meeting each generator  $\theta = \text{const.}$  in one point; in fact it is a directrix cubic  $\Delta$  given by

$$\{c\theta^3, c\theta^2, c\theta, c, -\theta^2(a\theta + b), -\theta(a\theta + b), -(a\theta + b)\}.$$

We have thus the  $\infty^2$  curves  $\Delta$  on  $F$ , and  $M_5^3$  can be generated by the solids  $K$  containing these curves.

All these curves  $\Delta$  can be obtained by means of prime sections through any two fixed generators; for the prime, whose equation is

$$a\{x_0 - (\theta_1 + \theta_2)x_1 + \theta_1\theta_2x_2\} + b\{x_1 - (\theta_1 + \theta_2)x_2 + \theta_1\theta_2x_3\} + c\{x_4 - (\theta_1 + \theta_2)x_5 + \theta_1\theta_2x_6\} = 0,$$

contains the curve and the pair of generators given by  $\theta = \theta_1$  and  $\theta = \theta_2$ .

If we consider the six primes

$$\begin{aligned} \Sigma_1 &\equiv ax_0 + \beta x_1 + \gamma x_2 = 0, & \Sigma_2 &\equiv ax_1 + \beta x_2 + \gamma x_3 = 0, & \Sigma_3 &\equiv ax_4 + \beta x_5 + \gamma x_6 = 0, \\ S_1 &\equiv ax_0 + bx_1 + cx_4 = 0, & S_2 &\equiv ax_1 + bx_2 + cx_5 = 0, & S_3 &\equiv ax_2 + bx_3 + cx_6 = 0, \end{aligned}$$

then  $a\Sigma_1 + b\Sigma_2 + c\Sigma_3 \equiv \alpha S_1 + \beta S_2 + \gamma S_3,$

so that the six primes have a line in common. Thus a solid  $K$  and a solid  $K'$  have a line of intersection.

Since all the first minors of the determinant

$$\begin{vmatrix} x_0 & x_1 & x_2 \\ x_1 & x_2 & x_3 \\ x_4 & x_5 & x_6 \end{vmatrix}$$

vanish at a point of  $F$ ,  $F$  must be a double surface on  $M_5^3$ . Since they also vanish at a point of the plane of  $\Gamma$  this plane must be a double plane on  $M_5^3$ .

It follows, as a consequence of the two methods of generating  $F$ , that an arbitrary solid of [6] is met in lines by six solids  $K$  and also by six solids  $K'$ , and that these lines

$$\begin{matrix} k_1, & k_2, & k_3, & k_4, & k_5, & k_6, \\ k_1', & k_2', & k_3', & k_4', & k_5', & k_6', \end{matrix}$$

form a double-six on the cubic surface which is the section of  $M_5^3$  by the arbitrary solid.

The quintic ruled surface  $F$  in [6] which has a directrix line has the equations

$$\frac{x_0}{x_1} = \frac{x_1}{x_2} = \frac{x_2}{x_3} = \frac{x_3}{x_4} = \frac{x_5}{x_6},$$

and its chords form the cubic primal

$$\begin{vmatrix} x_0 & x_1 & x_2 \\ x_1 & x_2 & x_3 \\ x_2 & x_3 & x_4 \end{vmatrix} = 0.$$

125. Suppose now that we take two fixed generators,  $g$  and  $g'$ , of  $F$ . Then the  $\infty^4$  directrix quartics of  $F$  are obtained\* by primes through  $g$  and also by primes through  $g'$ , and the system of  $\infty^4$  primes through  $g$  is thus related projectively to the system of  $\infty^4$  primes through  $g'$ . If we take a pencil of primes through  $g$  we obtain on  $F$  a pencil of directrix

\* § 45. This applies also to the normal surface  $F$  with a directrix line  $\lambda$ .

quartics with three common points; these three points being in fact the intersections (other than  $g$  itself) of  $F$  with the [4] which is the base of the pencil of primes. Corresponding to this [4] we have a [4] through  $g'$  containing the same three points of  $F$ . *The trisecant planes of  $F$  are thus the intersections of corresponding [4]'s of the projectivity.* The system of  $\infty^4$  primes through  $g$  contains  $\infty^6$  pencils of primes, the bases of these pencils being the [4]'s through  $g$ ; these, with the corresponding [4]'s through  $g'$ , give the  $\infty^6$  trisecant planes of  $F$ . The surface  $F$  itself is generated by the intersections of corresponding planes of the projectivity; there are  $\infty^4$  planes through  $g$  and  $\infty^4$  corresponding planes through  $g'$ , and, since two conditions are necessary in order that two planes in [6] should have a common point, there will be  $\infty^2$  planes through  $g$  which meet their corresponding planes through  $g'$ ; these  $\infty^2$  points are the points of the surface  $F$ .

126. We now enquire how many trisecant planes of  $F$  there are which meet an arbitrary plane  $\omega$  of [6] in lines; we expect that there will be a finite number\*.

The generator  $g$  and the plane  $\omega$  determine a [4]  $S_1$ ; corresponding to this we have a [4]  $S_1'$  through  $g'$  meeting  $S_1$  in a trisecant plane of  $F$ , and this trisecant plane, lying in the [4]  $S_1$  with  $\omega$ , meets  $\omega$  in a point  $P_1$ . Similarly, the generator  $g'$  and the plane  $\omega$  determine a [4]  $S_2'$ ; corresponding to this we have a [4]  $S_2$  through  $g$  meeting  $S_2'$  in a trisecant plane of  $F$ , this trisecant plane meeting  $\omega$  in a point  $P_2$ .

Suppose now that there is a trisecant plane  $\pi$  of  $F$  which meets  $\omega$  in a line. There is then a [5] through  $S_1$  which contains  $\pi$ , so that there must also be a [5] through  $S_1'$  containing  $\pi$ . Hence  $\pi$  meets  $S_1'$  in a line, and this line will have to meet the line of intersection of  $\pi$  with  $\omega$ , and therefore must pass through  $P_1$ , the point of intersection of  $S_1'$  and  $\omega$ . Hence  $\pi$  must pass through  $P_1$ , and similarly  $\pi$  must pass through  $P_2$ . If then there is a trisecant plane meeting  $\omega$  in a line, this line must be  $P_1P_2$ .

There are three† chords of  $F$  meeting  $P_1P_2$ , and these do in fact lie in a plane. For consider one of these chords meeting  $F$  in  $B$  and  $C$ . We have a [4] containing  $g$ ,  $P_1P_2$  and  $BC$ . Now the plane  $gP_2$  is the intersection of  $S_1$  and  $S_2$ , so that the corresponding plane through  $g'$  is the intersection of  $S_1'$  and  $S_2'$ , i.e. the plane  $g'P_1$ . Hence the [4] through  $g'$  which corresponds to the [4] through  $g$  containing  $P_2$  and  $BC$  must contain  $P_1$  and  $BC$ , and therefore the plane  $P_1P_2BC$ . We thus have corresponding [4]'s through  $g$  and  $g'$  meeting in the plane  $P_1P_2BC$ , which is therefore a trisecant plane of  $F$ . If  $A$  is its third point of intersection with  $F$  the three chords of  $F$  which meet  $P_1P_2$  are  $BC$ ,  $CA$ ,  $AB$ .

There is then *one trisecant plane of  $F$  which meets  $\omega$  in a line.*

\* § 53.

† § 123.

127. Any prime through  $\omega$  contains six chords of  $F$ , so that those chords of  $F$  which meet  $\omega$  meet  $F$  in the points of a curve  $C_{12}$  of order 12; this projects from  $\omega$  into the double curve of  $f$ .  $C_{12}$  has three double points at  $A, B, C$ , and, since the [4] through  $\omega$  and any generator meets  $F$  in three further points,  $C_{12}$  must meet each generator of  $F$  in three points. It cannot have any other double points, so that it is a curve  $C_{12}^4$  of genus\* 4.

The (3, 1) correspondence between  $C_{12}^4$  and  $\Gamma$  shews† that there are twelve generators of  $F$  touching  $C_{12}^4$ ; hence there are twelve generators of  $f$  touching the double curve.

Also the (2, 1) correspondence between  $C_{12}^4$  and  $c_3$ , the elliptic cubic curve in  $\omega$ , shews† that there are six tangents of  $C_{12}^4$  meeting  $\omega$ . Hence the tangents of  $F$  form a locus  $M_4^6$  of the sixth order‡.

128. *Rational quintic ruled surfaces in [4].* Before proceeding to investigate the surfaces  $f$  of [3] it will be convenient to interpolate here a few remarks upon the rational quintic ruled surface in [4]; this is obtained by projecting the general surface  $F$  in [6] from a line  $l$  which does not meet it. Some of the properties that we shall obtain are given by Severi||, but it is instructive to deduce them directly by projection.

We have seen that there are three points  $a, b, c$  on  $l$  through which there pass chords  $aa_1a_2, bb_1b_2, cc_1c_2$  of  $F$ . The  $S_4$  determined by these three chords must meet  $F$  in a directrix quartic, and on projecting from  $l$  on to the space [4] this becomes a plane quartic curve with three double points.

There is a solid  $K$  containing a directrix cubic of  $F$  which passes through  $a_1$  and  $a_2$ ;  $K$  containing the chord  $aa_1a_2$ . On projection this cubic becomes a plane cubic curve with a double point, this being at the same point as one of the double points of the quartic. We thus obtain three rational plane cubic curves on the surface in [4]; any two of these cubics intersect in the point of intersection of their planes, this being the projection of the point of intersection of the corresponding twisted cubics on  $F$ .

\* § 17.

† By Zeuthen's formula, § 16.

‡ The order of  $M_4$  is obtainable at once by elementary methods. For suppose  $F$  to be generated by a (1, 1) correspondence between  $\Gamma$  and one of the directrix cubics  $\Delta$ . The tangents of  $\Gamma$  and  $\Delta$  are also in (1, 1) correspondence, and the tangent solids of  $F$  are determined by the pairs of corresponding tangents. But the  $\infty^1$  solids determined by corresponding generators of two ruled surfaces, of orders 4 and 2, whose generators are in (1, 1) correspondence, form a locus of order  $4 + 2 = 6$ .

Similarly, the tangents of a rational ruled surface of order  $n$  form an  $M_4^{2n-4}$ .

|| See the footnote on p. 49 of his paper, "Intorno ai punti doppi impropri di una superficie generale dello spazio a quattro dimensioni e a' suoi punti tripli apparenti," *Palermo Rendiconti*, 15 (1901) 33.

Thus the general rational quintic ruled surface in [4] has three double points. It has a plane quartic on it; this lies in the plane of the three double points and has double points itself at these points. There are, further, three plane cubics, each of which has a double point at one of the double points of the surface. Further still, there is a directrix conic.

Of course we can obtain other surfaces in [4] by specialising the position of  $l$  in regard to  $F$ . If  $l$  lies in a solid  $K$  the projected surface has a triple line and is generated by a (1, 3) correspondence between a line and a conic. If  $l$  lies in a solid  $K'$  the projected surface has a double generator and contains  $\infty^1$  plane cubics with double points on this generator. If  $l$  is an axis of a directrix quartic  $E$  the projected surface has a double conic. It is generated by two conics in (1, 2) correspondence with a united point. We can also project the surface with a directrix line.

Similarly, ruled surfaces in [5] are obtained by projecting  $F$  from a point of [6]; a general point of [6] does not lie on  $M_5^3$ , so that the general rational quintic ruled surface in [5] has no double points. But if the point of projection does lie on a chord of  $F$  we obtain a surface in [5] with one double point and having a plane cubic on it with a node there, while if the point of projection lies in the plane of  $\Gamma$  we obtain a surface in [5] with a double line, generated by a (1, 2) correspondence between a line and a cubic curve.

*The surfaces in [3] derived by projection from the  
general surface in [6]*

129. We now proceed to obtain the rational quintic ruled surfaces of [3] by projection from the two normal surfaces in [6], and also to give methods for generating them. Of the twenty-four types which we have enumerated seventeen are obtained from the general surface  $F$ , the other seven arise from the surface  $F'$  with a directrix line.

We have already shewn how to obtain the most general surface  $f$ . Since  $F$  can be generated by placing its directrix conic  $\Gamma$  in (1, 1) correspondence with any one of its directrix cubics  $\Delta$ , the most general surface  $f$  is generated by a conic and a twisted cubic in (1, 1) correspondence.

130. Suppose that we take an ordinary involution  $I$  of pairs of generators on  $F$ . If we take any directrix quartic  $E$  on  $F$ ,  $I$  will determine an involution on  $E$ ; the chords of  $E$  joining the pairs of the involution form a cubic ruled surface lying in the  $S_4$  determined by  $E$ . This has a directrix line  $l$ , and a general plane  $\omega$  will not contain this line\*; but let us choose  $\omega$  to pass through such a line.

The plane  $\omega$  will meet  $M_5^3$  in  $l$  and a conic  $\mathfrak{D}$ . The chords of  $F$  which meet  $l$  will meet  $F$  in the points of the quartic  $E$ , while those which meet

\* The lines of [6] are  $\infty^{10}$  in aggregate and include  $\infty^6$  lines  $l$ .



$\mathfrak{D}$  will meet  $F$  in the points of an octavic  $C_8$ , meeting each generator of  $F$  in two points. The involution on  $E$  has two double points, so that there are two tangents of  $E$  which meet  $l$ ; hence there must be four tangents of  $C_8$  meeting  $\mathfrak{D}$ , the (2, 1) correspondence between  $C_8$  and  $\mathfrak{D}$  having four double points. Applying Zeuthen's formula to this correspondence we at once find that  $C_8$  is an elliptic curve, and then applying Segre's formula for the genus of a curve on a ruled surface we see that  $C_8$  has one double point  $A$ . There are two chords of  $F$  passing through  $A$ , meeting  $C_8$  again in  $B$  and  $C$ , and as these cannot be double points on  $C_8$  they must be intersections of  $C_8$  with  $E$ . The two chords of  $F$  which pass through the points common to  $l$  and  $\mathfrak{D}$  are chords both of  $C_8$  and  $E$ , so that we have four other intersections of these curves. There are, in fact, precisely six intersections\*.

Projecting from  $\omega$  on to the solid  $\Sigma$  we obtain a surface  $f$  with a double curve consisting of a conic (the projection of  $E$ ) and a quartic (the projection of  $C_8$ ); the quartic is rational since its points are in (1, 1) correspondence with those of  $\mathfrak{D}$ ; it has a double point (the projection of  $A, B$  and  $C$ ) lying on the conic and meets the conic in two other points†.

$F$  can be generated by placing  $\Gamma$  and  $E$  in (1, 1) correspondence with a united point. Hence  $f$  can be generated by two conics  $\Gamma_1$  and  $\Gamma_2$  in (1, 2) correspondence with a united point  $P$ . To  $P$ , regarded as a point of  $\Gamma_1$ , there correspond two points  $P, P'$  of  $\Gamma_2$  one of which is  $P$ ; to  $P$  regarded as a point of  $\Gamma_2$  corresponds the point  $P$  of  $\Gamma_1$ .

$\Gamma_1$  meets the plane of  $\Gamma_2$  in a second point  $Q$ ; to  $Q$  there correspond two points  $Q', Q''$  of  $\Gamma_2$ . The pairs of points of  $\Gamma_2$  which correspond to the points of  $\Gamma_1$  are the pairs of an involution; their joins all pass through  $O$ , the intersection of  $PP'$  and  $Q'Q''$ . The planes of the pairs of generators which intersect in the points of the double conic  $\Gamma_1$  are therefore formed by the points of  $\Gamma_1$  and the corresponding lines of a plane pencil in (1, 1) correspondence with  $\Gamma_1$ ; there is one united element, the point  $P$  of  $\Gamma_1$  lying on the line  $PP'$  of the pencil which corresponds to it. Thus‡ these planes touch a quadric cone  $E_2$ . This quadric cone is part of the bitangent developable of the surface.

The plane of  $\Gamma_2$  is a tritangent plane of the ruled surface since it contains the three generators  $PP', QQ', QQ''$ . It is a tangent plane of  $E_2$  as joining  $Q$  to the line  $OQ'Q''$ . The vertex of the cone  $E_2$  is  $O$ .

131. If any directrix quartic  $E$  is taken on  $F$  all the axes  $l$  arising from this lie on the cubic locus formed by the chords of  $E$ , and no two involutions on  $E$  can have the same axis||.

\* See the formula for  $i$  given in the footnote to § 97.

† Cf. § 92.

‡ § 22.

|| For the configuration of the axes of  $E$  see Segre, "Sulle varietà cubiche dello spazio a quattro dimensioni," *Memorie Torino* (2), 39 (1889), 3; in particular § 43.

Any two directrix quartics  $E$ ,  $E'$  intersect in three points; the plane of these three points is the plane of intersection of the two [4]'s which contain the curves. If then an axis of  $E$  coincided with an axis of  $E'$  it would lie in a trisecant plane of either curve, which is impossible.

If a plane  $\omega$  contains two axes of the same curve  $E$  it must be contained in the  $S_4$  to which  $E$  belongs; it then occupies a special position. The type of surface  $f$  arising by projection from such a plane will be subsequently considered.

Thus to obtain a surface  $f$  with two double conics we try to find a plane  $\omega$  containing an axis  $l_1$  of a directrix quartic  $E$  and an axis  $l_2$  of a directrix quartic  $E'$ . This we can certainly do; we have merely to take a common chord of  $E$  and  $E'$  and an axis of each curve passing through any point of this chord\*. The plane  $\omega$  meets  $M_5^3$  in the lines  $l_1$  and  $l_2$  together with a third line  $l_3$ , and the chords of  $F$  which pass through the points of  $l_3$  meet it in a third directrix quartic  $E''$ . The curves  $E'$  and  $E''$  have a common chord passing through the intersection of  $l_2$  and  $l_3$  and have also a third common intersection  $A$ , and we have similar points  $B$  and  $C$  arising from the other two pairs of curves. The plane  $ABC$  meets  $\omega$  in a line; the chords  $BC$ ,  $CA$ ,  $AB$  all meeting  $\omega$ .

Projecting from  $\omega$  on to a solid  $\Sigma$  we obtain a rational quintic ruled surface  $f$  with three double conics. The conics all pass through one point, while any two of them have a second intersection.

The surface  $F$  can be generated by a (1, 1) correspondence between  $E$  and  $E'$  with three united points. Thus to generate a surface  $f$  with two (and therefore three) double conics we take two conics  $\Gamma_1$  and  $\Gamma_2$  in  $\Sigma$  with two common points  $P$ ,  $Q$  and place them in (2, 2) correspondence. To the point  $P$  regarded as a point of  $\Gamma_1$  there correspond two points of  $\Gamma_2$  which both coincide with  $P$ , while to the point  $P$  regarded as a point of  $\Gamma_2$  there correspond two points of  $\Gamma_1$  which both coincide with  $P$ . To the point  $Q$  regarded as a point of  $\Gamma_1$  there correspond two points of  $\Gamma_2$ , one of which coincides with  $Q$ , while to the point  $Q$  regarded as a point of  $\Gamma_2$  there correspond two points of  $\Gamma_1$ , one of which coincides with  $Q$ . The lines joining corresponding points of  $\Gamma_1$  and  $\Gamma_2$  generate a rational quintic ruled surface with two double conics  $\Gamma_1$  and  $\Gamma_2$ .

132. We have seen that the prime  $\omega\Gamma$  contains three generators of  $F$ ; in general there will not be a solid passing through  $\omega$  and meeting each of these generators. But we may clearly choose  $\omega$  so that this happens, and then the surface  $f$ , instead of being of the type I, is specialised and is of the type II (A).

\* There are two axes of each curve passing through any point of the chord (Segre, *loc. cit.*).

Take any three points of  $\Gamma$  and the three generators through them. These determine a prime containing  $\Gamma$ ; take points  $X, Y, Z$  one on each of the three generators and a line  $l$  lying in the plane  $XYZ$ . Then if we take any plane  $\omega$  passing through  $l$  and lying in the prime and project from it on to  $\Sigma$  we obtain a surface  $f$  of the type II (A), having three generators passing through a point and lying in a plane.

When a surface  $f$  is generated by a conic and a twisted cubic in (1, 1) correspondence the plane of the conic is the tritangent plane. But, if we specialise the correspondence so that the lines which join the three points in which the cubic meets the plane of the conic to their corresponding points are concurrent, the three generators in the tritangent plane meet in the triple point.

This type of surface can be further specialised so as to have one or two double conics.

If, in the generation given in § 130, the correspondence between  $\Gamma_1$  and  $\Gamma_2$  is so specialised that  $PO$  passes through  $Q$  we obtain a surface with a double conic  $\Gamma_1$  which is of the type II (A).

In the generation at the end of § 131 the point  $Q$  is a triple point of the ruled surface; one generator through  $Q$  lies in the plane of  $\Gamma_1$  and a second in the plane of  $\Gamma_2$ . As a point approaches  $Q$  along one conic one of its corresponding points on the other conic must also approach  $Q$ ; the limiting position of the line joining the two points is the third generator through  $Q$ . If the correspondence is specialised so that these three generators lie in a plane we have a surface belonging to the type II (A).

133. A solid  $K'$  through two generators does not meet  $F$  again. If we choose  $\omega$  to pass through a line  $l$  of  $K'$  which does not meet either generator, and then project on to  $\Sigma$ , we obtain a surface  $f$  with a double generator  $G$ .

$\omega$  meets  $M_5^3$  in a line  $l$  and a conic  $\vartheta$ ; the chords of  $F$  which meet  $\omega$  will meet  $F$  in two generators and a curve  $C_{10}$  of order 10 meeting each generator in three points;  $C_{10}$  meets every solid  $K'$  in six points and every solid  $K$  in four points.

There are six tangents of  $C_{10}$  meeting  $\vartheta$ , and applying Zeuthen's formula to the (2, 1) correspondence between  $C_{10}$  and  $\vartheta$  we find that  $C_{10}$  is a hyperelliptic curve of genus 2; the  $g_2^1$  on  $C_{10}$  consists of the pairs of points which correspond to the points of  $\vartheta$ . Applying Segre's formula we see that  $C_{10}$  must have one double point  $A$ .

There are two chords of  $C_{10}$  passing through  $A$  and meeting  $\omega$ ; these meet  $C_{10}$  again in points  $B$  and  $C$  which must be intersections of  $C_{10}$  with the two generators of  $F$  that we originally selected. The chord  $BC$  meets  $l$ . On projecting we have a surface  $f$  with a double generator and a double quintic curve having a double point on the double generator. The quintic is rational since its points are in (1, 1) correspondence with those of  $\vartheta$ .

Further, the quintic meets the double generator in two other points, these arising from the chords of  $F$  which pass through the intersections of  $l$  and  $\vartheta$ . The generators of  $f$  are trisecants of the quintic\*. The bitangent developable consists of the planes through  $G$ , and a developable of the fifth class whose planes form a rational family in (1, 1) correspondence with the points of  $\vartheta$ . This developable has one double tangent plane and two ordinary tangent planes passing through  $G$ .

The prime  $\varpi\Gamma$  necessarily contains the two chosen generators of  $F$ , assuming that the planes of  $\varpi$  and  $\Gamma$  do not meet.

Hence to generate this surface  $f$  in  $\Sigma$  we take a conic and a twisted cubic in (1, 1) correspondence, but we so specialise the correspondence that two of the points in which the cubic meets the plane of the conic have for corresponding points those two points of the conic which lie on the line joining them.

134. Suppose now that  $\varpi$  is chosen to meet the plane of  $\Gamma$  in a point  $O$ . Then the projected surface  $f$  has a directrix line  $R$ , the intersection of  $\Sigma$  with the [4] containing  $\varpi$  and  $\Gamma$ . A prime through the [4]  $\varpi\Gamma$  contains three generators of  $F$ , so that a plane through  $R$  contains three generators of  $f$ . A solid passing through  $\varpi$  and lying in the [4]  $\varpi\Gamma$  meets two generators of  $F$ , so that through any point of  $R$  there pass two generators of  $f$ . This plainly indicates the types III (D) and III (F).

Suppose that there is a point  $X$  on  $R$  such that the two generators  $g_1$  and  $g_2$  of  $f$  which intersect there are co-planar with  $R$ .

The two generators of  $F$  which give rise to  $g_1$  and  $g_2$  must clearly lie in a prime containing  $\varpi$  and  $\Gamma$ . Such primes form a pencil, and each meets  $F$  in  $\Gamma$  and three generators; they thus cut a  $g_3^1$  on any directrix curve of  $F$ , the two generators in question passing through two points of a set of this  $g_3^1$ .

Further, the two spaces [4] which project the generators from  $\varpi$  must meet the [4]  $\varpi\Gamma$  in the same solid and therefore must meet the plane of  $\Gamma$  in the same line through  $O$ . The lines through  $O$  meet  $\Gamma$  in the sets of a  $g_2^1$  and the pairs of generators so determined give a  $g_2^1$  on any directrix curve.

Now, if we are given a  $g_2^1$  and a  $g_3^1$  on a rational curve, there are two † sets of the  $g_2^1$  belonging to sets of the  $g_3^1$ . Hence there are, in general, two pairs of generators of  $F$  such as we require.

\* Cf. § 102.

† The  $g_2^1$  may be regarded as a (1, 1) correspondence, and the  $g_3^1$  as a (2, 2) correspondence; both these correspondences are symmetrical, and are, in fact, involutions. Hence the number of common pairs of corresponding points is 2—half the number given by Brill's formula (see § 15). More generally, the number of pairs of points common to a  $g_m^1$  and a  $g_n^1$  on a curve of genus  $p$  is  $(m-1)(n-1)-p$ .

Hence, in general, there will be two points  $X$  on  $R$  such that the two generators of  $f$  which intersect there are co-planar with  $R$ , and then  $f$  is of the type III (F).

It may, however, happen that one of the two pairs of generators of  $F$  is such that there is a [4] through  $\omega$  containing them. The surface  $f$  has then a double generator and is of the type III (D).

If both pairs of generators are special in this way it appears that there is a directrix cubic  $\Delta$  of  $F$  for which the solid  $K$  meets  $\omega$  in a line;  $f$  is then a type of surface which will be subsequently considered.

For let  $gg'$  and  $hh'$  be the two pairs of generators; the solids  $gg'$  and  $hh'$  meeting  $\omega$  in the lines  $x$  and  $y$ . Take a point  $P$  on  $g$  and the transversal through  $P$  of  $gg'x$ , meeting  $g'$  in  $Q$  and  $x$  in  $R$ . The two points  $P$  and  $Q$  determine a directrix cubic  $\Delta$  meeting  $h$  and  $h'$  in  $S$  and  $T$ . Through  $S$  there passes a transversal of  $hh'y$ , meeting  $y$  in  $U$  and  $h'$  in  $T'$ .




Fig. 5.

Then the ranges  $(T)$  and  $(T')$  on  $h'$  are homographic, and have two common corresponding points. One of these is clearly on  $\Gamma$ ; the other gives a curve  $\Delta$  for which the solid  $K$  meets  $\omega$  in a line.

135. The generation of the surfaces  $f$  of the preceding article is at once obtained, since  $F$  can be generated by a (1, 1) correspondence between  $\Gamma$  and any of its directrix cubics. We take in  $\Sigma$  a line and a twisted cubic in (1, 2) correspondence; the joins of corresponding points give a quintic ruled surface with the line as a double directrix.

The pairs of points of the cubic which correspond to the points of the line form an ordinary rational involution; the chords joining the pairs therefore form a regulus. Hence there will be two of these chords meeting the line in points  $A$  and  $B$ . Through  $A$  passes a chord  $Aaa'$ , the pair of points  $aa'$  corresponding to a point  $\alpha$  of the line; through  $B$  passes a chord  $Bbb'$ , the pair of points  $bb'$  corresponding to a point  $\beta$  of the line. In general, the surface is of the type III (F); but if either  $\alpha$  coincides with  $A$  or  $\beta$  with

*B* we have a surface of the type III (D) with a double generator. If  $\alpha$  coincides with *A* and  $\beta$  coincides with *B* the surface *f* degenerates into a type with two directrices.

If a (1, 1) correspondence has been established between the points of a line and the lines of a regulus then the planes containing corresponding elements form a developable of the third class\*. If one of the two lines of the regulus meets the line in the point which corresponds to it the planes touch a quadric cone, while if both lines of the regulus which meet the line do so in their corresponding points the planes all pass through a line. This confirms the result found for the bitangent developables of the surfaces III (F), III (D) and VI (B).

**136.** When  $\omega$  has been chosen to meet the plane of  $\Gamma$  in a point the [4]  $\omega\Gamma$  will not, in general, contain a generator of *F*, but we may clearly choose  $\omega$  so that it does. Then the projected surface *f* will have a directrix line *R* which is also a generator; any plane through *R* contains two other generators, while through any point of *R* there pass two other generators.

It is easily seen that, in general, there is one point of *R* at which these two other generators are co-planar with *R*; *f* is then of the type IV (D). We have to consider two linear series on a directrix curve of *F*; one  $g_2^1$  and a second  $g_2^1$  which arises from a  $g_3^1$  having a fixed point. These have one pair of points in common.

It may, however, happen that the common pair of these two linear series gives two generators which lie in the same [4] with  $\omega$ ; we may clearly choose  $\omega$  so that this happens. Then *f* has a double generator and is of the type IV (C).

**137.** We again consider *F* as generated by a (1, 1) correspondence between  $\Gamma$  and one of its directrix cubics  $\Delta$ . The [4]  $\omega\Gamma$  contains a generator *g*, and, when we project on to  $\Sigma$ ,  $\Gamma$  and  $\Delta$  become a line and a twisted cubic which intersect in a point *P*, the projection of the point  $g\Delta$ . Thus to generate *f* we take a line and twisted cubic meeting in a point *P* and establish a (1, 2) correspondence between them. To any point of the line correspond two points of the cubic, while to any point of the cubic corresponds one point of the line, *P* not being special in any way. We have a surface *f* of the type IV (D).

The pairs of points of the cubic which correspond to the points of the line give rise to lines forming a regulus meeting the line in two points *P* and *Q*. The planes of the pairs of generators issuing from the points of the line thus form a developable of the third class which is part of the bitangent developable of *f*.

\* See § 23 above.

Through  $Q$  there passes a chord  $Qaa'$  of the cubic;  $a$  and  $a'$  forming the pair of points corresponding to some point  $\alpha$  of the line. If then  $\alpha$  coincides with  $Q$  we have a surface  $f$  with a double generator of the type IV (C); this has a quadric cone as part of its bitangent developable.

138. Further, when  $\omega$  meets the plane of  $\Gamma$  in a point, it may happen that the [4]  $\omega\Gamma$  contains two generators of  $F$ . Then  $f$  has a directrix line  $R$  which is also a double generator; any plane through  $R$  contains one other generator, while through any point of  $R$  there pass two other generators. Hence  $f$  is of the type V (A).

To generate  $f$  we take a twisted cubic and one of its chords and place them in (2, 1) correspondence without united points. Those planes of the bitangent developable which do not pass through the chord form a developable of the third class.

139. We have seen that of the  $\infty^2$  solids  $K$  there is a singly infinite set meeting a general plane  $\omega$  in points but that there is none meeting it in a line. If, however, we choose  $\omega$  to meet a solid  $K$  in a line the projected surface  $f$  has a directrix line  $R$ , the intersection of  $\Sigma$  with the [4]  $\omega K$ . Since any prime through the [4]  $\omega K$  meets  $F$  in a directrix cubic  $\Delta$  and two generators we see that any plane through  $R$  contains two generators of  $f$ , and since any solid passing through  $\omega$  and lying in the [4]  $\omega K$  meets  $F$  in three points there will be three generators of  $f$  passing through any point of  $R$ . This indicates the types III (C) and III (E).

Suppose that there is a plane  $x$  through  $R$  such that the pair of generators which it contains intersect in a point on  $R$ .

The two spaces [4] joining  $\omega$  to the two generators of  $F$  from which these arise must meet the [4]  $\omega K$  in the same solid, and must therefore meet  $K$  in the same plane. This plane will contain the line of intersection of  $\omega$  and  $K$ , so that the two generators of  $F$  meet  $\Delta$  in a pair of points belonging to a set of a  $g_3^1$  cut out on  $\Delta$  by planes of  $K$  passing through a line. Also the two generators of  $F$  must lie in a prime through  $\omega$  and  $K$ ; such primes form a pencil, each meeting  $F$  in  $\Delta$  and two generators, giving thus a  $g_2^1$  on  $\Delta$ . But the  $g_2^1$  has two sets belonging to sets of the  $g_3^1$ . Hence, in general, there are two planes  $x$  through  $R$  such as we are seeking, and  $f$  is of the type III (E).

It may happen that one of these two pairs of generators of  $F$  is such that there is a [4] through  $\omega$  containing them. Then  $f$  has a double generator and is of the type III (C).

If both pairs of generators are special in this way it appears that  $\omega$  must meet the plane of  $\Gamma$  in a point; the surface  $f$  then belongs to a type which will be subsequently considered.

For let one pair of generators be  $g$  and  $g'$ , the other pair  $h$  and  $h'$ . The solid  $gg'$  will meet  $\omega$  in a line  $x$ ; the solid  $hh'$  will meet  $\omega$  in a line  $y$ . Let  $G, G', H, H'$  be the points of intersection of  $\Gamma$  with the four generators. Then  $GG'$  and  $HH'$  intersect, and this point of intersection is the point common to the two solids  $gg'$  and  $hh'$  and is therefore also the point of intersection of the lines  $x, y$ .

**140.** Since  $F$  is generated by a (1, 1) correspondence between  $\Delta$  and  $\Gamma$ ,  $f$  can be generated by a (1, 3) correspondence between a line and a conic; to any point of the line  $R$  correspond three points of the conic, while to any point of the conic corresponds one point of the line  $R$ .

We have an involution of sets of three points on the conic, and the sets of three joins of all such triads are known to touch a conic  $\Gamma_0^*$ . There are two tangents to  $\Gamma_0$  from the point in which the line  $R$  meets its plane; these give the two pairs of generators<sup>c</sup> of  $f$  which lie in planes through  $R$  and intersect in points of  $R$ . The surface is, in general, of the type III (E).

The point of intersection of  $R$  with the plane of the conic gives three corresponding points of the conic. If two of these happen to be collinear with the point on  $R$  we have a surface  $f$  of the type III (C) with a double generator.

**141.** When  $\omega$  is chosen to meet a solid  $K$  in a line the [4]  $\omega K$  will not, in general, contain a generator of  $F$ , but we may clearly choose  $\omega$  so that it does. Then, on projecting, the surface  $f$  has a directrix  $R$  which is also a generator. Any plane through  $R$  contains one other generator, while through any point of  $R$  there pass three generators. Hence  $F$  is of the type IV (A).

To generate  $f$  we again take a line  $R$  and a conic in (1, 3) correspondence, but here they have a point of intersection. This point does not specialise the correspondence in any way.

The point of intersection of  $R$  and the conic is the projection from  $\omega$  on to  $\Sigma$  of the point of intersection of  $\Gamma$  with the generator of  $F$  which lies in the [4]  $\omega K$ .

**142.** We now investigate a type of surface  $f$  to which we have already twice referred. Suppose that  $\omega$  is chosen to meet a solid  $K$  in a line and also to meet the plane of  $\Gamma$  in a point  $O$ . Then the projected surface  $f$  has two directrices;  $R$ , the intersection of  $\Sigma$  with the [4]  $\omega K$ , and  $R'$ , the intersection of  $\Sigma$  with the [4]  $\omega \Gamma$ .

Any prime through  $\omega$  and  $K$  meets  $F$  in  $\Delta$  and two generators which must meet  $\Gamma$  in a pair of points collinear with  $O$ . Also any solid through

\* Cf. Baker, *Principles of Geometry*, 2 (Cambridge, 1922), 137.



$\omega$  lying in the [4]  $\omega\Gamma$  meets  $\Gamma$  in two points collinear with  $O$ ; there is then a prime containing these two generators and  $\omega$ , which necessarily contains  $K$  also. Hence any plane through  $R$  contains two generators of  $f$  meeting in a point of  $R'$ , while through any point of  $R'$  there pass two generators of  $f$  lying in a plane with  $R$ . Similarly, any plane through  $R'$  contains three generators of  $f$  meeting in a point of  $R$ , while through any point of  $R$  there pass three generators lying in a plane with  $R'$ .

Now let us enquire whether  $f$  has any double generators. This is the same question as whether there exist spaces  $S_4$  through  $\omega$  which contain two generators of  $F$ . Such an  $S_4$  would have to meet the plane of  $\Gamma$  in a line through  $O$  and the solid  $K$  in a plane. Conversely, if we have a pair of generators of  $F$  such that the chords of  $\Gamma$  and  $\Delta$  determined by the pair both meet  $\omega$ , there is an  $S_4$  through  $\omega$  containing the pair of generators. Now the chords of  $\Gamma$  through  $O$  determine an involution on  $\Gamma$  and thus also a  $g_2^1$  on  $\Delta$ ; while the planes of  $K$  passing through its line of intersection with  $\omega$  determine a  $g_3^1$  on  $\Delta$ . Hence there are two pairs of generators of  $F$  such as we require, and  $f$  has two double generators.  $f$  is of the type VI (B).

The (1, 1) correspondence between  $\Delta$  and  $\Gamma$  clearly gives rise to a (2, 3) correspondence between  $R$  and  $R'$ , and  $f$  can be generated by means of this correspondence. But this is not the most general (2, 3) correspondence between two lines; it must be specialised in order to give the two double generators.

143. We can give another method of generating this surface of the type VI (B) which leads at once to a generation for a surface of the type VII (A).

$\omega$  and a generator of  $F$  determine a space  $S_4$ ; any prime through this meets  $F$  in the generator and in a directrix quartic  $E$ ;  $E$  meets  $\Gamma$  in one point and  $\Delta$  in two points. The [4] containing  $\omega$  and  $\Delta$  will meet the prime containing  $\omega$  and  $E$  in a solid which contains  $\omega$  and the two intersections of  $E$  and  $\Delta$ . Hence the line joining these two points must meet  $\omega$ . The  $S_4$  containing  $E$  meets  $\omega$  in a line which is met by three chords of  $E$ , one of these being already accounted for.

When we project,  $E$  gives rise to a plane quartic with three double points; it passes through the point in which  $R'$  meets its plane and one of its double points lies at the point in which  $R$  meets its plane.

Now  $F$  can be generated by means of a (1, 1) correspondence between  $\Gamma$  and  $E$  with a united point. Hence, to generate  $f$ , we take a plane quartic with three double points and a line  $R'$  meeting it in a point  $P$ . To any point of  $R'$  there correspond two points of the quartic, while to any point of the quartic corresponds one point of  $R'$ ,  $P$  being a united point.

But it is necessary that the two points which correspond to any point on  $R'$  should be collinear with one of the double points. For, returning to the surface  $F$  in [6], we have seen that the prime containing  $\omega$  and two generators  $g$  and  $g'$ , which meet  $\Gamma$  in a pair of points collinear with  $O$ , also contains  $K$ . It therefore meets  $E$  in four points; one on  $g$ , one on  $g'$ , and the two on  $\Delta$  already mentioned. This shews on projection that, given a point of  $R'$ , the two corresponding points on the quartic are collinear with the double point on  $R$ .

The two other double points of the quartic determine two double generators of  $f$ .

The line joining the points in which  $R$  and  $R'$  meet the plane of the quartic is the generator of  $f$  which lies in this plane.

This method of generation at once suggests the following for a surface of the type VII (A).

Take a plane quartic with three double points  $A, B, C$  and a line  $R$ , not in the plane of the quartic, passing through  $A$ . Establish a (1, 2) correspondence between  $R$  and the quartic, with a united point, such that to any point of  $R$  there correspond two points of the quartic collinear with  $A$ : this is at once secured by referring the range of points on  $R$  to the pencil ( $A$ ) of lines through  $A$  in the plane of the quartic. The point  $A$ , regarded as a point of  $R$ , must give rise to two points of the quartic, one of which is  $A$  itself: this is secured by making the point  $A$  on the range  $R$  correspond to one of the tangents of the quartic at  $A$  considered as a line of the pencil ( $A$ ). There will be a second point on  $R$  which corresponds to  $A$ ; this being that point of the range which corresponds to the other tangent of the curve at  $A$  considered as a line of the pencil ( $A$ ). Then we have a ruled surface  $f$  with a directrix line which is also a generator; through each point of  $R$  there pass two generators lying in a plane with  $R$ , while each plane through  $R$  contains two generators intersecting in a point of  $R$ . There are two double generators passing through the points  $B$  and  $C$ . This is the surface of the type VII (A).

144. We can examine in closer detail the way in which  $\omega$  may be chosen so as to give a surface of the type VII (A).

Take any two generators  $g$  and  $g'$  of  $F$  meeting  $\Gamma$  in the points  $G$  and  $G'$ , and any point  $O$  on the line  $GG'$ ; also a point  $X$  on  $F$ . Then there are  $\infty^1$  directrix cubics passing through  $X$  and their solids  $K$  meet the solid  $gg'$  in lines forming a regulus.  $GG'$  belongs to this regulus; this line arises from the degenerate cubic consisting of  $\Gamma$  and the generator  $g_0$  through  $X$ . Any line through  $O$  lying in the solid  $gg'$  will meet the regulus in a second point  $O'$ , the line of the regulus through  $O'$  being a chord of a directrix cubic  $\Delta$ . There are planes through  $OO'$  meeting the solid  $K$  containing

$\Delta$  in lines, and on projection from such a plane we obtain a surface  $f$  of the type VI (B).

But suppose that we take a tangent, at  $O$ , of the quadric surface on which the regulus lies; and then a plane  $\omega$  through this tangent, containing a line passing through  $O$ , and lying in the solid  $g_0\Gamma$ . On projecting we have a surface  $f$  with a directrix  $R$  which is also a generator.

Through  $\omega$  and two generators of  $F$  which meet  $\Gamma$  in two points collinear with  $O$  there passes a prime, which necessarily contains  $\Gamma$  and  $g_0$ ; so that through any point of  $R$  there pass two other generators which lie in a plane with  $R$ , and  $f$  is of the type VII (A).

145.  $F$  has  $\infty^4$  quartic directrices, and  $\omega$  will meet the [4] containing such a directrix in a point. There are, however,  $\infty^2$  quartic directrices such that the [4]'s containing them meet  $\omega$  in lines; in fact, any line of  $\omega$  lies in such a [4], determined by the three chords of  $F$  which meet the line. But, in general, there is no [4] so arising which contains  $\omega$  entirely; this is evident when we remember that all directrix quartics can be obtained as residuals of prime sections of  $F$  through any fixed generator.

Let us suppose, however, that  $\omega$  is chosen to lie in a space  $S_4$  containing a directrix quartic  $E$ . Then the projected surface  $f$  has a directrix line  $R$ , the intersection of  $\Sigma$  with  $S_4$ . Since a prime through  $S_4$  meets  $F$  in  $E$  and one generator there will be one generator of  $f$  contained in any plane passing through  $R$ , and since any solid lying in  $S_4$  and passing through  $\omega$  meets  $E$  in four points there are four generators of  $f$  passing through each point of  $R$ . Hence  $f$  is of the type III (A).

The cubic locus in  $S_4$  formed by the chords of  $E$  is none other than the section of  $M_5^3$  by  $S_4$ ; in general,  $\omega$  meets this locus in an ordinary non-degenerate elliptic cubic curve, and the bitangent developable of  $f$  is non-degenerate and of the sixth class, having a tritangent plane. The planes of the developable form an elliptic family, since they are in (1, 1) correspondence with the points of the cubic curve in  $\omega$ . The tritangent plane is, of course, the intersection of  $\Sigma$  with the prime  $\omega\Gamma$ .

But  $\omega$  may be chosen to pass through an axis  $l$  of  $E$ ; the chords of  $E$  through the points of  $l$  determining an involution on  $E$  and thus an involution  $I$  of pairs of generators of  $F$ .  $\omega$  meets  $M_5^3$  in  $l$  and a conic  $\mathfrak{S}$ . We have already seen (§ 130) that the bitangent planes of  $f$  which arise from  $I$  touch a quadric cone. The residual developable is of the fourth class, with two ordinary tangent planes and one double tangent plane which touch the cone also. This last plane clearly arises from the prime  $\omega\Gamma$ ; the other two are determined by the chords of  $F$  which pass through the two points of intersection of  $l$  and  $\mathfrak{S}$ .

Further,  $\omega$  may be chosen to contain two, and therefore three, axes of  $E$ , say  $l_1, l_2, l_3$ . Then the bitangent developable of  $f$  consists of three

quadric cones. The prime  $\omega\Gamma$  meets  $\Sigma$  in a plane which touches each cone. Also the chord of  $F$  which passes through the intersection of  $l_2$  and  $l_3$ , say, meets a pair of generators which give on projection a common tangent plane of the two cones which arise from  $l_2$  and  $l_3$ ; thus any two of the three cones have a second common tangent plane.

**146.**  $F$  can be generated by placing  $\Gamma$  and  $E$  in (1, 1) correspondence with a united point. Hence to generate this type of surface  $f$  we take a line  $R$  and a conic which meets it, and place them in (1, 4) correspondence with their intersection as a united point. Any plane through  $R$  contains one generator, while through every point of  $R$  there pass four generators. The plane of the conic contains three generators all passing through the point in which  $R$  meets it; it is thus a tritangent plane.

But if the plane  $\omega$  contains an axis of  $E$  then, corresponding to each point of  $R$ , we have four points of the conic as before; but two of these four points form a pair of a certain involution, so that the line joining them passes through a fixed point  $O$ . The bitangent developable of  $f$  breaks up into two parts.

The pencil of lines through  $O$  in the plane of the conic is related to the range of points on  $R$ , and the planes joining the rays of the pencil to the corresponding points on  $R$  touch a quadric cone with vertex  $O$ , which touches the plane of the conic. Hence the bitangent developable breaks up into this cone and a developable of the fourth class having the plane of the conic for a double tangent plane.

The joins of the sets of an involution of sets of four points on a conic are known to touch a curve of the third class. In this degenerate case the curve will consist of the point  $O$  and a conic, and two tangents of this last conic will pass through  $O$ . This shews that the quadric cone and the developable of the fourth class have two other common tangent planes.

Now suppose that the plane  $\omega$  contains three axes of  $E$ . Any solid through  $\omega$  meets the cubic locus of chords of  $E$  in a four-nodal cubic surface, the nodes being the four points in which the solid meets  $E$ . The three axes are the three lines on this surface other than the six edges of the tetrahedron formed by the nodes, and it is a well-known property of the four-nodal cubic surface that each of these three lines meets a pair of opposite edges of the tetrahedron. If, then, any point  $A$  is taken on  $E$  the solid  $\omega A$  meets  $E$  again in three points  $B, C, D$  such that  $AB$  and  $CD$  meet one axis,  $AC$  and  $BD$  meet another, while  $AD$  and  $BC$  meet the third.

Thus in the correspondence between  $R$  and the conic if  $a, b, c, d$  are the four points of the conic which correspond to some point of  $R$  we have  $ab$  and  $cd$  belonging to one fixed involution,  $ac$  and  $bd$  belonging to a second and  $ad$  and  $bc$  belonging to a third. We thus have three points  $O_1, O_2, O_3$  forming a self-conjugate triangle in regard to the conic; these being the

three points through which the joins of pairs of the three involutions respectively pass. The envelope of the third class formed by the joins of the sets of four points here degenerates into the three points  $O_1, O_2, O_3$ .

Thus the bitangent developable of  $f$  consists of three quadric cones with vertices  $O_1, O_2, O_3$  all touching the plane  $O_1O_2O_3$ . Any two of the cones have a second common tangent plane.

This completes the determination of those surfaces  $f$  which are projections of the surface  $F$  with a directrix conic.

*The surfaces in [3] derived by projection from the surface in [6] with a directrix line*

147. Take now a rational quintic ruled surface  $F$  in [6] with a directrix line  $\lambda$ . Any chord of  $F$  lies in a solid through  $\lambda$  containing two generators of  $F$ , so that the locus  $\dot{M}_5$  of the chords of  $F$  is a five-dimensional cone having a line-vertex and  $\infty^2$  generating solids. If we take a directrix quartic  $E$  of  $F$  the [4] in which it lies meets  $M_5$  in the cubic locus formed by the chords of  $E$ ; we have thus a cubic line-cone  $M_5^3$ .

No two generating solids of the line-cone can intersect, unless on the three-dimensional locus formed by the planes through  $\lambda$  containing the generators of  $F$ . Since a general plane  $\omega$  does not meet this locus it will meet  $M_5^3$  in a cubic curve without a double point.

Take now a general plane  $\omega$  of [6] not meeting  $F$  and project from  $\omega$  on to a solid  $\Sigma$ . We obtain a rational quintic ruled surface  $f$  with a directrix line  $R$ , the intersection of  $\Sigma$  with the [4]  $\omega\lambda$ . Since a prime through  $\omega$  and  $\lambda$  meets  $F$  in  $\lambda$  and four generators there are four generators of the surface  $f$  which lie in any plane through  $R$ , and since a solid passing through  $\omega$  and lying in the [4]  $\omega\lambda$  meets  $\lambda$  in a point there is one generator of  $f$  passing through any point of  $R$ . Hence  $f$  is of the type III (B).

It is seen as in § 127 that the chords of  $F$  which meet  $\omega$  meet  $F$  in the points of a curve  $C_{12}$  of order 12 and genus 4, having three double points whose plane meets  $\omega$  in a line,  $C_{12}$  meeting each generator of  $F$  in three points. The double curve of  $f$  is an elliptic sextic curve with a triple point.

The locus  $M_4^6$  formed by the tangents of  $F$  is here again of the sixth order, but it is now a line-cone with  $\lambda$  for vertex and has  $\infty^1$  generating solids all passing through  $\lambda$ . It is of the sixth order, because the [4] which contains any directrix quartic  $E$  meets  $M_4^6$  in the ruled surface formed by the tangents of  $E$ , and this is known to be of the sixth order\*.

This type of surface  $F$  contains  $\infty^4$  directrix quartics  $E$ ; any one of these is determined by four points of  $F$ , and any two intersect in three

\* Cf. the footnote to § 127.

points. They can all be obtained as residuals of prime sections of  $F$  through any fixed generator. Each curve  $E$  determines a space [4] containing it. In general, such a space [4] will meet  $\omega$  in a point, but there are  $\infty^2$  of them meeting  $\omega$  in lines, one passing through each line of  $\omega$ . For a general position of  $\omega$  no [4] will contain it entirely.

Now  $\omega$  may clearly be chosen to pass through an axis  $l$  of one of these curves  $E$ , meeting  $M_5^3$  in  $l$  and a conic  $\vartheta$ . The chords of  $F$  meeting  $l$  meet  $F$  on  $E$  itself, those meeting  $\vartheta$  meet  $F$  in an elliptic curve of the eighth order, with one double point, which meets each generator in two points. On projecting we have a surface  $f$  of the type III (B) whose double curve consists of a conic and a rational quartic; this quartic has a double point lying on the conic and meets the conic in two other points.

Further,  $\omega$  may be chosen to pass through an axis  $l_1$  of a quartic  $E$  and an axis  $l_2$  of a quartic  $E'$ . It meets  $M_5^3$  in three lines  $l_1, l_2, l_3$ , and  $l_3$  is an axis of a quartic  $E''$ . The surface  $f$  has a double curve consisting of three conics with a common point, any two of the conics intersecting in one other point\*.

148.  $F$  can be generated by means of a (1, 1) correspondence between  $\lambda$  and any of its directrix quartics  $E$ . Hence to generate the most general surface  $f$  belonging to the type III (B) we take a line and a rational skew quartic and place them in (1, 1) correspondence.

If, however,  $\omega$  contains an axis of  $E$  then  $E$  becomes on projection a double conic of  $f$ , so that to generate this surface we take a line and a conic in (2, 1) correspondence.

The construction of the quintic surface with a directrix line and two double conics can be deduced from that for the quintic ruled surface with two double conics without a directrix line† simply by making the (2, 2) correspondence between  $\Gamma_1$  and  $\Gamma_2$  express the condition that the generators should all meet a line. Or we can deduce a construction from that of the dual surface; we take a line lying in a tangent plane of a quadric cone and place the planes through the line and the tangent planes of the cone in (1, 4) correspondence. The four planes of the cone which correspond to any plane through the line form pairs of three involutions.

149. Suppose now that  $\omega$  is chosen to lie in a space [4] with one of the directrix quartics  $E$ . Then the projected surface  $f$  has two directrices;  $R$ , the intersection of  $\Sigma$  with the [4]  $\omega\lambda$ , and  $R'$ , the intersection of  $\Sigma$  with the [4]  $\omega E$ . A prime through  $\omega$  and  $\lambda$  meets  $F$  in  $\lambda$  and four generators, and these must meet  $E$  in four points lying in a solid through  $\omega$ . Also, if a solid through  $\omega$  meets  $E$  in four points the four generators of  $F$  passing through these points all meet  $\lambda$ . Hence any plane through  $R$

\* Cf. § 131.

† § 131.

contains four generators of  $f$  which meet in a point of  $R'$ , while through any point of  $R'$  there pass four generators of  $f$  lying in a plane with  $R$ . Similarly any plane through  $R'$  contains one generator of  $f$  which meets  $R$ , while through any point of  $R$  there passes one generator of  $f$  meeting  $R'$ . Hence  $f$  is of the type VI (A).

Since  $F$  can be generated by a (1, 1) correspondence between  $\lambda$  and  $E$ , the surface  $f$  can be generated by a (1, 4) correspondence between  $R'$  and  $R$ .

150. Any prime meets  $F$  in a rational normal quintic curve, one point of which lies on  $\lambda$ ; the chords of the curve passing through this point form a cone of the fourth order. This shews that the  $\infty^1$  planes joining  $\lambda$  to the generators of  $F$  form a locus  $M_3^4$  of three dimensions and the fourth order. In general this has no point in common with a plane  $\omega$ . But if  $\omega$  does meet a plane of  $M_3^4$  we have on projection a surface  $f$  with a directrix line  $R$  which is also a generator.

Since a prime through  $\omega$  and  $\lambda$  now meets  $F$  in three variable generators there are three generators of  $f$  which lie in a plane through  $R$  other than  $R$  itself. Also through any point of  $R$  there passes one generator of  $f$  other than  $R$ . Hence  $f$  is of the type IV (B). To generate  $f$  we take a line and a rational quartic in  $\Sigma$  which meet, and place them in (1, 1) correspondence without a united point.

We may choose  $\omega$  to meet two planes of  $M_3^4$ . Then the surface  $f$  has a directrix  $R$  which is at the same time a double generator. Any plane through  $R$  contains two generators of  $f$  other than  $R$ , while through any point of  $R$  there passes one generator other than  $R$ . Hence  $f$  is of the type V (B). To generate  $f$  we take in  $\Sigma$  a rational quartic and a line meeting it in two points, and place them in (1, 1) correspondence without any united points.

Further, we may choose  $\omega$  to meet three planes of  $M_3^4$ . Then  $f$  has a directrix  $R$  which is also a triple generator. Any plane through  $R$  contains one other generator, while through any point of  $R$  there passes one other generator. Hence  $f$  is of the type VII (B). To generate  $f$  we take in  $\Sigma$  a rational quartic and one of its trisecants, placing them in (1, 1) correspondence without any united points.

## SECTION III

## QUINTIC RULED SURFACES WHICH ARE NOT RATIONAL

*Elliptic quintic ruled surfaces considered as curves on  $\Omega$* 

151. *Elliptic quintic curves on  $\Omega$ .* The elliptic curve of the fifth order is normal in [4]\*. Also we can have on a quadric in [3] an elliptic quintic curve meeting all the generators of one system in three points and all of the other system in two points, the curve having a double point; it is the intersection of the quadric with a cubic surface passing through one of its generators and touching it in a point. Similarly we can have an elliptic quintic curve lying on a quadric cone; the curve has a double point and passes through the vertex, meeting every generator of the cone in two points other than the vertex.

We can thus write down the following types of elliptic quintic curves  $C$  which lie on  $\Omega$ , numbering them consecutively with the types of rational quintic curves.

VIII. The normal elliptic quintic curve  $C$  lying in a [4] which does not touch  $\Omega$ .

IX.  $C$  lies in a tangent prime  $T$  touching  $\Omega$  in a point  $O$  but does not pass through  $O$ .

(A)  $C$  meets every plane  $\omega$  through  $O$  in three points and every plane  $\rho$  through  $O$  in two points, a chord of  $C$  passing through  $O$ .

(B)  $C$  meets every plane  $\omega$  through  $O$  in two points and every plane  $\rho$  through  $O$  in three points, a chord of  $C$  passing through  $O$ .

X.  $C$  lies in the tangent prime  $T$  touching  $\Omega$  in a point  $O$  and passes through  $O$ , meeting every plane of  $\Omega$  in two points other than  $O$ .

XI.  $C$  lies on the quadric  $Q$  in which  $\Omega$  is met by a space  $S_3$  through which pass two tangent primes.  $C$  has a double point, meeting all the generators of one system of  $Q$  in three points and all of the other system in two points.

XII.  $C$  lies on the quadric cone in which  $\Omega$  is met by a space  $S_3$  which itself touches  $\Omega$  in a point  $V$ .  $C$  has a double point, passes through  $V$ , and meets every generator of the cone in two points other than  $V$ .

152. *The general type of surface.* Suppose that we have a normal elliptic quintic  $C$  in [4]. The projection of  $C$  from any line on to a plane is a quintic curve with five double points, so that any line is met by five chords of  $C$ . Hence the chords of  $C$  form a locus  $V_3^5$  of three dimensions and the fifth order.



Hence if  $C$  belongs to the type VIII any plane of  $\Omega$  meets the [4] containing  $C$  in a line which is met by five chords of  $C$ . Hence we have a ruled surface in [3] whose double curve is of the fifth order and whose bitangent developable is of the fifth class.

If  $C$  is projected on to a plane from a line which meets it we obtain a quartic with two double points, so that the line is met by two chords other than those passing through its intersection with  $C$ . Hence  $C$  is triple on  $V_3^5$ . If  $C$  is projected from one of its chords on to a plane we obtain an elliptic cubic without a double point; thus there will not be any double surface on  $V_3^5$ .

The intersection of  $\Omega$  with  $V_3^5$  is a surface  $F_2^{10}$  on which  $C$  is a triple curve; this is the ruled surface formed by the chords of  $C$  which lie on  $\Omega$ , three such chords passing through any point of  $C$ . The section of this ruled surface by a solid is a curve of order 10 with five triple points; the curve lying on a quadric and meeting every generator in five points. Projecting from a point of this quadric on to a plane we obtain a curve of order 10 with five triple points and two quintuple points which is therefore of genus

$$36 - 15 - 20 = 1.$$

But this curve is representative of the double curve and bitangent developable of the surface in [3]; hence the double curve is an elliptic quintic curve, while the planes of the bitangent developable form an elliptic family.

If we take an elliptic quintic curve in ordinary space then, on projecting it from a point of itself on to a plane, we obtain a quartic with two double points; hence there are two trisecants of the curve passing through any point of it, assuming that the curve itself has not a double point. The surface is, in fact, generated by the trisecants of an elliptic quintic curve without a double point\*.

153. Suppose now that  $C$  belongs to the type IX (A). The point  $O$  represents a line  $R$  which is a directrix of the ruled surface.

An arbitrary plane  $\rho$  meets  $T$  in a line which is met by five chords of  $C$ . The plane  $\omega$  through this line joins it to  $O$  and contains three points and therefore three chords of  $C$ . There are two other chords meeting the line. Hence the double curve of the corresponding ruled surface consists of the line  $R$  counted three times and a conic.  $R$  and the conic intersect, their point of intersection being represented on  $\Omega$  by the  $\omega$ -plane through the chord of  $C$  which passes through  $O$ .

An arbitrary plane  $\omega$  meets  $T$  in a line, and the plane  $\rho$  through this line contains two points and therefore one chord of  $C$ . Four other chords

\* See the reference to Zeuthen in § 89.

of  $C$  meet the line. Hence the bitangent developable consists of the planes through  $R$  and a developable of class four. There is a plane of this developable passing through  $R$ ; it is represented on  $\Omega$  by the  $\rho$ -plane which contains the chord of  $C$  passing through  $O$ .

**154.** The ruled surface formed by the chords of  $C$  which lie on  $\Omega$  must clearly break up into two parts; one will consist of the chords joined to  $O$  by  $\rho$ -planes and the other of the chords joined to  $O$  by  $\omega$ -planes.

The chords of  $C$  which lie in  $\rho$ -planes join the pairs of an ordinary rational involution on  $C$ ; hence they form a cubic ruled surface\*. The plane  $\rho$  which passes through the directrix of this surface represents the plane of the double conic. Then the chords of  $C$  which lie in  $\omega$ -planes must form a ruled surface of order seven with  $C$  for a double curve.

We now take the section of this composite ruled surface by a solid lying in  $T$ ; the solid meets  $\Omega$  in a quadric, and we have on this quadric a cubic curve meeting every generator of one system in two points and every generator of the other system in one point, together with a septic curve meeting every generator of the first system in three points and every generator of the other system in four points. This septic has five double points through which the cubic passes. The cubic is rational but the septic is elliptic, for on projecting it from a point of the quadric on to a plane we obtain a plane curve of order seven with one quadruple point, one triple point and five double points, and therefore of genus

$$15 - 6 - 3 - 5 = 1.$$

Two points of  $C$  which lie in a  $\rho$ -plane through  $O$  represent two generators of the ruled surface whose plane passes through  $R$  and whose point of intersection lies on the double conic; the conic is a rational curve and the pencil of planes through  $R$  is a rational family. But  $R$  regarded as a triple line must be considered an elliptic curve, while the planes of the developable of the fourth class form an elliptic family.

If we apply a general result for the number of intersections of two curves lying on the same quadric† we find that the cubic and septic intersect in eleven points; ten of these are accounted for by the five double points on the septic, the other arises from the chord of  $C$  passing through  $O$ .

**155.** If  $C$  is of the type IX (B) we have a ruled surface with a directrix line  $R$ ; through every point of  $R$  there pass two generators, while every plane through  $R$  contains three generators. The double curve consists of

\* See the footnote to § 19.

† If there are two curves on a quadric of which one meets all generators of one system in  $x$  points and all of the other system in  $y$  points, while for the second curve the corresponding numbers are  $x'$  and  $y'$ , the number of intersections of the two curves is  $xy' + x'y$ .

the line  $R$  and an elliptic quartic curve which meets  $R$ ; the bitangent developable consists of the planes through  $R$  counted three times together with the tangent planes of a quadric cone, one tangent plane of this cone passing through  $R$ .

This surface can be generated by taking an elliptic quartic curve and a line  $R$  meeting it in a point; the generators of the surface are those chords of the curve which meet  $R$ .

**156.** Suppose now that  $C$  is of the type X. Then the ruled surface has a directrix line  $R$  which is itself a generator; through each point of  $R$  there pass two generators other than  $R$ , while each plane through  $R$  contains two generators other than  $R$ .

Any plane of  $\Omega$  meets  $T$  in a line and the plane of the opposite system through this line joins it to  $O$ ; this contains three points and therefore three chords of  $C$ , so that two other chords will meet the line. Hence the double curve consists of  $R$  counted three times and a conic, while the bitangent developable will consist of the planes through  $R$  counted three times together with the tangent planes of a quadric cone. The conic will meet  $R$  and a tangent plane of the cone will pass through  $R$ .

The ruled surface formed by the chords of  $C$  which lie on  $\Omega$  here breaks up into two cubic ruled surfaces and the elliptic quartic cone projecting  $C$  from  $O$ .

**157.** Now let  $C$  be of the type XI. There are two tangent primes of  $\Omega$  through  $S_3$ , which touch it in points  $O$  and  $O'$ . These represent two lines  $R$  and  $R'$  which are both directrices of the ruled surface.  $C$  has a double point  $P$  which represents a double generator  $G$ .

Suppose that the generators of  $Q$  which are trisecants of  $C$  lie in the  $\omega$ -planes through  $O$  and the  $\rho$ -planes through  $O'$ . Then the generators of the other system will lie in the  $\rho$ -planes through  $O$  and the  $\omega$ -planes through  $O'$ . Through any point of  $R$  there pass three generators of the surface which lie in a plane through  $R'$ ; any plane through  $R'$  contains three generators meeting in a point of  $R$ . Any plane through  $R$  contains two generators meeting in a point of  $R'$ , while through any point of  $R'$  there pass two generators lying in a plane through  $R$ .

Any plane of  $\Omega$  meets  $S_3$  in a point of  $Q$ .  $C$  has five apparent double points, but the five chords which can be drawn from a point of  $Q$  consist of the trisecant generator through the point counted three times, the other generator through the point, and the line to the double point.

The double curve of the ruled surface consists of  $R'$  and  $G$  together with  $R$  counted three times; the bitangent developable consists of the planes through  $R$  and  $G$  together with those through  $R'$  counted three times.

**158.** Finally, suppose that  $C$  is of the type XII. Then the ruled surface has a directrix  $R$  which is itself a generator; through any point of  $R$  there pass two generators other than  $R$  which lie in a plane with  $R$ ; any plane through  $R$  contains two generators which intersect in a point of  $R$ .

$C$  has five apparent double points. But if  $C$  is projected from a point of the cone on to a plane we obtain an elliptic quintic with a double point, and a triple point at which two branches touch. Hence the five chords of  $C$  which pass through a point of the cone consist of the line to the double point  $P$ , and the generator counted four times.

The double curve of the ruled surface thus consists of the double generator  $G$ , together with the directrix  $R$  counted four times; the bitangent developable consists of the planes through  $G$ , and the planes through  $R$  counted four times.

A plane section of the surface is an elliptic quintic curve with a double point on  $G$  and a triple point, at which two branches touch, on  $R$ .

**159.** We have now obtained six different types of elliptic quintic ruled surfaces in [3]; these are exhibited in tabular form on p. 305.

*Elliptic quintic ruled surfaces in [3] considered as projections of normal surfaces in higher space*

**160.** A ruled surface of the fifth order which belongs to a space [6] is rational, its prime sections being rational normal quintic curves.

Suppose now that we have a ruled surface of the fifth order belonging to a space [5]. If the surface is elliptic a prime through one of its generators meets it again in an elliptic quartic curve, unless the surface is a cone. The elliptic quartic curve lies in a [3]; through this [3] there passes a pencil of primes. The surface must be contained in one of these primes, since otherwise each prime of the pencil would contain a generator of the surface, which would then be rational\*. Hence *an elliptic quintic ruled surface which is not a cone belongs to a space of dimension 3 or 4.*

Suppose now that we have an elliptic quintic ruled surface in [3]; a quadric which contains one of its generators  $g$  will meet it again in a curve  $C_9$  of order 9. This curve meets each generator of the surface in two points, and those two points where it meets  $g$  are points of contact of the quadric and the ruled surface. The quadric can be chosen so that it does not touch the ruled surface elsewhere, so that the only double points of  $C_9$  will be on the double curve of the ruled surface. Hence  $C_9$  will be of genus 5.

\* The generators would be in (1, 1) correspondence with the primes of a pencil and therefore with the points of a line.

† § 17.

Now a curve  $C_9$  of genus 5 in [3] is the projection\* of a normal curve of order 9 and genus 5 in [4]. The generators of the ruled surface in [3] join pairs of points of an elliptic involution on  $C_9^5$ ; these are the projections of the joins of pairs of points of an elliptic involution on the normal curve, and these joins are generators of an elliptic ruled surface in [4]. Since we have on this ruled surface a curve of order 9 and genus 5 meeting each generator in two points the surface must be of order † 5; the elliptic quintic ruled surface in [3] is then the projection of this elliptic quintic ruled surface in [4]. Hence we have the result: *the elliptic quintic ruled surface is normal in [4]; and any elliptic quintic ruled surface in [3] can be obtained as the projection of a normal surface in [4].*

We now consider the normal elliptic quintic ruled surfaces  $F$  of [4]. We find that there are two types of surface, on one of which no two generators intersect, while on the other the generators intersect each other in pairs. By projection we obtain from the first surface two of the types of elliptic quintic ruled surfaces which we have just obtained in [3]; the other four types of surfaces in [3] are obtained by projection from the normal surface with a double line.

161. Suppose then that we have, in [4], an elliptic quintic ruled surface  $F$ , of which no two generators lie in a plane. The section of  $F$  by a solid will be a directrix (i.e. a curve meeting each generator of the surface in one point) together with, perhaps, a certain number of generators. But since the directrix must be an elliptic curve it cannot be a line or a conic, so that no three generators of  $F$  can lie in the same solid.

Any two generators of  $F$  determine a solid which must meet  $F$  further in a plane cubic curve without a double point; we have  $\infty^2$  solids determined by the pairs of generators of  $F$ , while through the plane of any cubic curve on  $F$  there pass  $\infty^1$  solids each meeting  $F$  again in two generators. We have therefore  $\infty^1$  cubic directrices on  $F$ ; these will be the directrices of minimum order and there will be a finite number of them passing through any point of  $F$ .

Every cubic curve lies in a solid with any fixed generator  $g$  of  $F$ ; to obtain then the cubic curves passing through any point  $P$  of  $F$  we have to take those solids which contain the plane  $gP$  and meet  $F$  in another generator. The number of such solids is clearly the number of remaining intersections of the plane  $gP$  with  $F$ . Now any plane  $\alpha$  through  $g$  meets  $F$  in three points not on  $g$ ; for a solid through  $\alpha$  meets  $F$  again in an elliptic quartic curve which meets  $\alpha$  in four points, one of these four points is on  $g$  and the other three give isolated intersections of  $\alpha$  with  $F$ . Hence the plane  $gP$  meets  $F$  in two further points, so that through  $P$  there pass two of the cubic curves.

\* § 8.

† By the formula of § 17.

We have then the following result: *the normal elliptic quintic ruled surface  $F$  without double points has, as directrices of minimum order,  $\infty^1$  plane elliptic cubic curves; through each point of  $F$  there pass two of these curves.*

Any two of the cubic curves are put into (1, 1) correspondence by the generators of  $F$ ; this correspondence must have a united point\*, so that any two of the cubic curves intersect. We notice that, in the (1, 1) correspondence between any two of the cubic curves, three collinear points of one curve do not correspond to three collinear points of the other; for if they did we should have three generators of  $F$  lying in the same solid.

The planes of the cubic curves form a locus  $W_3$  of three dimensions. The section of this locus by a solid consists of the trisecants of the elliptic quintic curve in which the solid meets  $F$ , and is therefore an elliptic quintic ruled surface †. Hence  $W_3$  is a locus  $W_3^5$  of the fifth order. It is elliptic when considered as a family of  $\infty^1$  planes; the planes of  $W_3^5$  are in (1, 1) correspondence with the generators of  $F$ , since, if any cubic curve of  $F$  is taken, there is one plane of  $W_3^5$  passing through each point of it. The point of intersection of two planes of  $W_3^5$  is the intersection of the two cubic curves which they contain and therefore lies on  $F$ ;  $F$  is a double surface on  $W_3^5$ .

**162.** Now project  $F$  from a point  $O$  on to a [3]  $\Sigma$ . The lines joining  $O$  to the points of  $F$  meet  $\Sigma$  in the points of an elliptic quintic ruled surface  $f$ , while the planes joining  $O$  to the generators of  $F$  meet  $\Sigma$  in the generators of  $f$ .

A solid through  $O$  meets  $F$  in an elliptic quintic curve; this has five apparent double points, or five of its chords pass through  $O$ . Hence the double curve of  $f$  is of the fifth order, a plane section of  $f$  being an elliptic quintic curve with five double points.

The class of the bitangent developable of  $f$  is the number of solids which contain a given line through  $O$  and also two generators of  $F$ . Such a solid meets  $F$  further in a cubic curve, whose plane must then meet the line through  $O$ . Hence the class of the bitangent developable of  $f$  is equal to the order of  $W_3^5$ ; we have a bitangent developable of the fifth class.

We have a surface in [3] of the type VIII.

Since  $F$  is generated by two elliptic cubic curves in (1, 1) correspondence with a united point,  $f$  is also generated by two elliptic cubic curves (whose planes both lie in  $\Sigma$ ) in (1, 1) correspondence with a united point.

**163.** This surface  $f$  is the most general type of elliptic quintic ruled surface in [3] and has been obtained by projecting  $F$  from a point  $O$  of general position in [4]. But suppose now that for  $O$  is taken a point of  $W_3^5$ .

\* § 19.

† § 152.

Then the plane  $\omega$  of  $W_3^5$  in which  $O$  lies meets  $\Sigma$  in a line  $R$  which is a directrix of  $f$ . Since any line of  $\omega$  passing through  $O$  meets  $F$  in three points there are three generators of  $f$  passing through each point of  $R$ , and, since any solid through  $\omega$  meets  $F$  in two generators, there are two generators of  $f$  lying in any plane through  $R$ . Hence we have a surface  $f$  of the type IX (A).

We can generate  $F$  by establishing a (1, 1) correspondence between the cubic curve in  $\omega$  and another cubic directrix, there being a united point. Hence  $f$  is generated by means of a (1, 3) correspondence, with a united point, between a line  $R$  and an elliptic cubic curve.

These two types of surfaces are the only ones which can be obtained by projection from the normal surface  $F$  which we have been considering.

164. We suppose now that we have, in [4], a normal elliptic quintic ruled surface  $F$ , two of whose generators intersect. A solid containing the plane of these two generators meets  $F$  further in an elliptic cubic curve; we thus obtain  $\infty^1$  directrix cubic curves. Now the plane of any one of these cubic curves  $\Gamma$  meets the plane of the two intersecting generators in a line; this line meets  $\Gamma$  in three points, one on each of the generators. Thus there is a point  $X$  in which the plane of the two generators meets  $\Gamma$  and which does not lie on either of them. Then there is a solid containing the plane of the two generators and also the generator through  $X$ ; the residual intersection with  $F$  will be a directrix curve of order 2, and, since the surface is elliptic, this will be a *double line*  $\lambda$ . Hence, if there are two generators of  $F$  which intersect, all the generators of  $F$  intersect one another in pairs, and the locus of the points of intersection is a line  $\lambda$ .

The solid which we have just considered, containing a pair of intersecting generators and the generator through  $X$ , meets  $F$  in these three generators together with  $\lambda$ ; hence the point  $X$  is the only point of  $F$  which lies in the plane of the two intersecting generators and not on either of the generators themselves. But each of the  $\infty^1$  directrix cubics of  $F$  meets this plane in a point which does not lie on either generator; hence all the cubic curves pass through  $X$ . And, since the same argument may be applied to any pair of intersecting generators, the plane of every pair of intersecting generators passes through  $X$ .

We have then, in [4], *a normal elliptic quintic ruled surface  $F$  with a double line  $\lambda$ ; this surface contains  $\infty^1$  plane elliptic cubic curves all passing through the same point  $X$ , while through a general point of  $F$  there passes one of the cubic curves.*

The surface is generated by a line  $\lambda$  and an elliptic cubic curve in (1, 2) correspondence.

We may of course also generate the surface by means of a (1, 1) correspondence between two of its directrix cubics,  $X$  being a united point.

Now if three collinear points are taken on one of the cubics the three generators all meet  $\lambda$  and therefore lie in the same solid; they therefore meet any other one of the directrix cubics in three collinear points, so that the correspondence between any two of the cubic curves is part of a projectivity between their planes. This kind of  $(1, 1)$  correspondence between two elliptic cubics will be called *special*. This explains clearly the difference between the two types of normal surface  $F$ . If we take, in [4], two elliptic cubic curves with an intersection  $X$  and place them in  $(1, 1)$  correspondence, then the lines joining pairs of corresponding points generate an elliptic quintic ruled surface  $F$ . In general,  $F$  is of the former type and has no double point, but if the correspondence is special the surface is of the type which we are now considering and has a double line  $\lambda$ . The pairs of intersecting generators meet either of the cubic curves in pairs of points collinear with  $X$ .

The plane containing any cubic curve of  $F$  meets the plane containing any pair of intersecting generators in a line through  $X$ ; the planes containing two cubic curves do not meet, except in  $X$ , and the planes containing two pairs of intersecting generators do not meet, except in  $X$ . Hence we have the two systems of planes of a quadric point-cone with vertex  $X$ . There is a generator through  $X$  meeting  $\lambda$ , these forming together a degenerate cubic curve of  $F$ ; the plane  $\pi_0$  containing them is met in a line through  $X$  by any plane containing a pair of intersecting generators.

The planes joining  $\lambda$  to the generators of  $F$  form an elliptic line-cone; this cone is of the third order, a prime section being the cubic cone of lines projecting an elliptic quintic curve from a double point. The plane  $\pi_0$  belongs to this cone; in fact the plane  $\pi_0$  and the surface  $F$  form together the complete intersection of this cubic line-cone with the quadric point-cone.

**165.** Let us now project  $F$  from a point  $O$ , of general position in [4], on to a solid  $\Sigma$ . The plane  $O\lambda$  meets  $\Sigma$  in a line  $R$  which is a directrix of the projected surface  $f$ ; through each point of  $R$  there pass two generators of  $f$ . Also, since any solid through the plane  $O\lambda$  meets  $F$  in three generators, any plane through  $R$  contains three generators of  $f$ . Thus  $f$  is of the type IX (B).

To generate  $f$  we take, in  $\Sigma$ , a line  $R$  and an elliptic cubic, and place them in  $(1, 2)$  correspondence.

**166.** Now let us specialise the position of  $O$ , so that it lies on the cubic line-cone containing  $F$ . Then  $O$  lies in a plane containing  $\lambda$  and a generator, so that this plane meets  $\Sigma$  in a line  $R$  which is a directrix and also a generator of  $f$ . Through any point of  $R$  there pass two generators of  $f$  other than  $R$  itself. Any solid through  $O$  and  $\lambda$  meets  $F$  in three generators, one of which is the fixed generator lying in the plane  $O\lambda$ . Hence any plane



through  $R$  contains two generators of  $F$  other than  $R$  itself. Thus  $f$  is of the type X.

To generate  $f$  we take a line  $R$  and an elliptic cubic curve;  $R$  and the cubic intersect and we place them in (1, 2) correspondence without a united point.

Now let us take  $O$  to lie on the quadric point-cone containing  $F$ ; then it lies in a plane  $\pi$  containing a directrix cubic of  $F$  and also in a plane  $\pi'$  containing a pair of intersecting generators of  $F$ .

The projected surface  $f$  will have two directrices;  $R$ , the intersection of  $\Sigma$  with the plane joining  $O$  to the double line, and  $R'$ , the intersection of  $\Sigma$  with  $\pi$ . We have also a double generator, the intersection of  $\Sigma$  with  $\pi'$ .

A prime through the plane joining  $O$  to the double line meets  $\pi$  in a line passing through  $O$  and  $F$  in three generators which meet this line. Thus a plane through  $R$  contains three generators of  $f$  which meet in a point of  $R'$ . Also a prime through  $\pi$  meets the double line in a point and contains the two generators of  $F$  which intersect there. Thus a plane through  $R'$  contains two generators of  $f$  which meet in a point of  $R$ .

This surface  $f$  is of the type XI. It can be generated by means of a (2, 3) correspondence between  $R'$  and  $R$ , but this is not the most general (2, 3) correspondence between two lines; it must be specialised to give the double generator.

Now let us choose  $O$  in the plane  $\pi_0$ . The plane  $\pi_0$  meets  $\Sigma$  in a line  $R$  which is a directrix and also a generator of  $f$ . Also the line  $OX$  meets the double line of  $F$ , and the plane of the two generators of  $F$  intersecting in this point passes through  $OX$ . Hence  $f$  has a double generator.

Through any point of the double line of  $F$  there pass two generators; and the plane of these two generators, which passes through  $X$ , meets  $\pi_0$  in a line. Hence through any point of  $R$  there pass two generators of  $f$ , other than  $R$  which is itself a generator, and the plane of these two generators contains  $R$ . This surface  $f$  is of the type XII.

To generate this last type of surface we take a line and an elliptic cubic which have a point of intersection and place them in (1, 2) correspondence without a united point. But the two points of the cubic curve which correspond to a point of the line must be collinear with the point in which the line meets the curve. The double generator passes through this point and lies in the plane of the cubic.

### *The quintic ruled surface of genus 2*

167. If we have a quintic ruled surface of genus 2 in a space [3], its generators will be represented by the points of a quintic curve  $C$  of genus 2 lying on the quadric  $\Omega$  in [5]. Now a quintic curve of genus 2 is necessarily contained in a space  $S_3^*$ , so that we have two possibilities.

XIII.  $C$  lies on the quadric  $Q$  in which  $\Omega$  is met by a space  $S_3$  through which pass two tangent primes.  $C$  meets every generator of  $Q$  of one system in three points and every generator of the other system in two points.

XIV.  $C$  lies on a quadric cone in which  $\Omega$  is met by an  $S_3$  which touches it in a point  $V$ .  $C$  passes through  $V$  and meets every generator of the cone in two points other than  $V$ .

If there are two tangent primes through  $S_3$  touching  $\Omega$  in  $O$  and  $O'$  then these two points represent lines  $R$  and  $R'$  of the original space [3] which are both directrices of the ruled surface. Each generator of  $Q$  is the intersection of a plane of  $\Omega$  through  $O$  with a plane of the opposite system through  $O'$ . Suppose that the generators of  $Q$  which are trisecants of  $C$  lie in the  $\varpi$ -planes through  $O$  and the  $\rho$ -planes through  $O'$ ; the other generators will lie in the  $\varpi$ -planes through  $O'$  and the  $\rho$ -planes through  $O$ . Through every point of  $R$  there pass three generators lying in a plane through  $R'$ , while through any point of  $R'$  there pass two generators lying in a plane through  $R$ .

The curve  $C$  has four apparent double points, four of its chords passing through any point of  $S_3$ . But if this point lies on  $Q$  three of the chords coalesce in the trisecant generator through the point,  $C$  projecting into a plane quintic curve with a triple point and a double point. The double curve of the ruled surface consists of the line  $R'$  together with the line  $R$  counted three times.

Similarly the bitangent developable consists of the planes through  $R'$  together with those through  $R$  counted three times.

This is the surface generated by the most general (2, 3) correspondence between two lines  $R$  and  $R'$ .

If  $S_3$  touches  $\Omega$  in a point  $V$  then this represents a line  $R$  which is a directrix and also a generator of the ruled surface. Through each generator on the cone in which  $S_3$  meets  $\Omega$  there passes a plane of each system of  $\Omega$ , and the two planes lie in the tangent prime of  $\Omega$  at  $V$ . Hence through any point of  $R$  there pass two generators of the ruled surface which lie in a plane with  $R$ .

The curve  $C$  has four apparent double points, but the four chords of  $C$  which pass through any point of the cone all coincide with the generator through the point; the projection of the curve from this point on to a plane of  $S_3$  is a quintic curve with a triple point at which two of the branches touch each other.

The double curve of the ruled surface is the line  $R$  counted four times, while the bitangent developable consists of the planes through  $R$  counted four times.

168. Since a quintic curve of genus 2 is necessarily contained in [3] a quintic ruled surface whose prime sections are of genus 2 is necessarily

contained in [4]. But those surfaces belonging to [4] are cones\*, so that the surface is normal in [3] and the question of projection from higher space does not arise.

The surface cannot contain a directrix line which is not a multiple line, nor a conic, nor a cubic.

Consider the surface with two directrices  $R$  and  $R'$ . A plane through a generator meets the surface again in a quartic curve with one double point, this double point being on  $R$ . The quartic meets the generator again on  $R'$  and also in the point of contact of the plane with the surface. We thus obtain  $\infty^2$  plane quartic curves of genus 2 on the surface.

Through two general points of the surface there pass three of these curves, for the line joining the two points meets the surface again in three further points, and there are planes through the first two points and the generator through any one of these last three points. Two of the quartics intersect in three points on the line of intersection of their planes.

**169.** We can generate the surface by means of a (1, 2) correspondence between a line  $R'$  and a plane quartic with one double point  $P$ . The quartic must pass through  $P'$ , the point in which  $R'$  meets its plane, and  $P'$  must be a united point for the correspondence.

The points of  $R'$  will give rise to the pairs of the  $g_2^1$  on the curve, and these are collinear with  $P$  †. The points of the range on  $R'$  are related projectively to the rays of the pencil through  $P$ ; the point  $P'$  of the range corresponding to the ray  $PP'$  of the pencil. The planes joining the corresponding elements will therefore all pass through a line  $R$  which passes through  $P$  ‡. Through any point of  $R'$  there pass two generators of the surface which lie in a plane through  $R$ , while any plane through  $R'$  contains three generators meeting in a point of  $R$ . The line  $PP'$  is a generator.

To obtain the surface with one directrix we have only to make  $P$  and  $P'$  coincide. We take a plane quartic with one double point  $P$  and a line  $R$  passing through  $P$ ; we then place the line and curve in (1, 2) correspondence,  $P$  being one of the two points on the quartic which corre-

\* If the surface were not a cone a [3] through a generator would meet it further in a quartic curve of genus 2. This is a plane curve, and we have an argument similar to that of § 160.

† There can only exist one  $g_2^1$  on a curve of genus 2; it is the canonical series of the curve, and for a plane quartic with one double point  $P$  it is cut out by the lines through  $P$ .

‡ If the point  $P'$  did not correspond to the ray  $PP'$  then the planes joining corresponding elements would touch a quadric cone.

spond to  $P$  regarded as a point of the line. This is at once secured by referring the range of points on  $R$  to the pencil of lines through  $P$ , so that the point  $P$  of the range corresponds to one of the tangents of the curve at  $P$ . The other tangent of the curve at  $P$  will correspond to some other point of  $R$ , and the line  $R$  is a generator as well as a directrix. Through any point of  $R$  there pass two generators (other than  $R$  itself) which lie in a plane through  $R$ . The generator of the surface which lies in the plane of the quartic curve is one of the tangents at the double point.

# CHAPTER IV

## SEXTIC RULED SURFACES

### SECTION I

#### RATIONAL SEXTIC RULED SURFACES

170. We shall in this chapter enumerate completely the different kinds of sextic ruled surfaces in [3]. So far as is known, no serious attempt has been made to solve this problem before, but if we employ the two powerful methods which we have already used in classifying the quintic ruled surfaces it should be possible to arrive at a solution.

In the first place we regard the generators of the ruled surface as the points of a sextic curve on a quadric  $\Omega$  in [5]; in the second place we regard the ruled surface as the projection of a normal ruled surface in higher space. We have up to the present pursued these two lines of investigation separately; we shall also pursue them separately for the ruled surfaces of the sixth order which are not rational, but since the ruled surfaces of the sixth order which are rational are so numerous we shall for these surfaces pursue the two lines of investigation concurrently, all the information concerning one particular type of surface being then collected together.

#### *The rational sextic curves which lie on quadrics*

171. All rational sextic curves can be obtained by projection from the normal curve in [6]. As we are only considering curves which lie on a quadric  $\Omega$  in [5] this normal curve does not enter directly into our work, but we can always resort to it in order to obtain properties of the actual curves that we are using.

We then divide the rational sextic curves  $C$  on  $\Omega$  into eight classes:

- I.  $C$  is contained in [5].
- II.  $C$  is contained in a [4] which does not touch  $\Omega$ .
- III.  $C$  lies in a tangent prime  $T$  of  $\Omega$  but does not pass through  $O$ , the point of contact of  $\Omega$  and  $T$ .
- IV.  $C$  lies in  $T$  and passes through  $O$ .
- V.  $C$  lies in  $T$  and has a double point at  $O$ .
- VI.  $C$  lies in  $T$  and has a triple point at  $O$ .
- VII.  $C$  lies in a [3] meeting  $\Omega$  in an ordinary quadric surface.
- VIII.  $C$  lies in a [3] meeting  $\Omega$  in a quadric cone.

We now give a more complete classification of these curves  $C$  as follows:

- I. (A)  $C$  is contained in [5] and has no double point.  
(B)  $C$  is contained in [5] and has a double point.

- II. (A)  $C$  is contained in [4] and has no singularity.  
 (B)  $C$  is contained in [4] and has a double point.  
 (C)  $C$  is contained in [4] and has two double points.  
 (D)  $C$  is contained in [4] and has a triple point.
- III. (A)  $C$  is  $\omega_5\rho_1^*$ .  
 (B)  $C$  is  $\omega_1\rho_5$ .  
 (C)  $C$  is  $\omega_4\rho_2$ , three chords of  $C$  passing through  $O$ .  
 (D)  $C$  is  $\omega_2\rho_4$ , three chords of  $C$  passing through  $O$ .  
 (E)  $C$  is  $\omega_4\rho_2$  with a double point, two chords of  $C$  passing through  $O$ .  
 (F)  $C$  is  $\omega_2\rho_4$  with a double point, two chords of  $C$  passing through  $O$ .  
 (G)  $C$  is  $\omega_4\rho_2$  with two double points, a chord of  $C$  passing through  $O$ .  
 (H)  $C$  is  $\omega_2\rho_4$  with two double points, a chord of  $C$  passing through  $O$ .  
 (I)  $C$  is  $\omega_3\rho_3$ , four chords of  $C$  passing through  $O$ .  
 (J)  $C$  is  $\omega_3\rho_3$  with a double point, three chords of  $C$  passing through  $O$ .  
 (K)  $C$  is  $\omega_3\rho_3$  with two double points, two chords of  $C$  passing through  $O$ .  
 (L)  $C$  is  $\omega_3\rho_3$  with a triple point, a chord of  $C$  passing through  $O$ .
- IV. (A)  $C$  is  $O\omega_4\rho_1$ .  
 (B)  $C$  is  $O\omega_1\rho_4$ .  
 (C)  $C$  is  $O\omega_3\rho_2$ , two trisecants of  $C$  passing through  $O$ .  
 (D)  $C$  is  $O\omega_2\rho_3$ , two trisecants of  $C$  passing through  $O$ .  
 (E)  $C$  is  $O\omega_3\rho_2$  with a double point, a trisecant of  $C$  passing through  $O$ .  
 (F)  $C$  is  $O\omega_2\rho_3$  with a double point, a trisecant of  $C$  passing through  $O$ .  
 (G)  $C$  is  $O\omega_3\rho_2$  with two double points.  
 (H)  $C$  is  $O\omega_2\rho_3$  with two double points.
- V. (A)  $C$  is  $O^2\omega_3\rho_1$ .  
 (B)  $C$  is  $O^2\omega_1\rho_3$ .  
 (C)  $C$  is  $O^2\omega_2\rho_2$ ; there being a line through  $O$  meeting  $C$  in two further points.  
 (D)  $C$  is  $O^2\omega_2\rho_2$  with another double point besides  $O$ .
- VI. (A)  $C$  is  $O^3\omega_2\rho_1$ .  
 (B)  $C$  is  $O^3\omega_1\rho_2$ .
- VII. (A)  $C$  is the intersection of a quadric and a quintic surface passing through four generators of the same system; it meets all generators of this system in five points and all of the other system in one point.  
 (B)  $C$  is the intersection of a quadric with a quartic surface touching it in three points and containing two generators of the same

\* This notation is self-explanatory: " $C$  is  $\omega_5\rho_1$ " means " $C$  meets every plane  $\omega$  of  $\Omega$  through  $O$  in five points and every plane  $\rho$  of  $\Omega$  through  $O$  in one point." Similarly, in V (D), " $C$  is  $O^2\omega_2\rho_2$ " means " $C$  has a double point at  $O$  and meets every plane of  $\Omega$  through  $O$  in two points other than  $O$ ."

system; it meets all generators of this system in four points and all of the other system in two points, having three double points.

- (C)  $C$  is the intersection of a quadric with a cubic surface touching it in four points; it meets every generator of the quadric in three points and has four double points.

VIII. (A)  $C$  is the intersection of a quadric cone with a cubic surface having a node at the vertex of the cone; it has a quadruple point at the vertex and meets each generator of the cone in one point other than the vertex.

- (B)  $C$  is the intersection of a quadric cone with a cubic surface passing through the vertex of the cone and touching the cone three times; it has a double point at the vertex and three other double points, meeting each generator of the cone in two points other than the vertex.

- (C)  $C$  is the intersection of a quadric cone with a cubic surface touching it in four points. It meets each generator of the cone in three points and has four double points.

We have thus divided the rational sextic curves  $C$  which lie on a quadric  $\Omega$  in [5] into thirty-eight classes, and these will form a basis for the classification of the rational sextic ruled surfaces in [3].

172. All these rational sextic curves can of course be obtained by projection from the rational normal sextic in [6]. To obtain, for example, the four classes in II we project the normal curve ( $a$ ) from an arbitrary line in [6], ( $b$ ) from a line meeting one of its chords, ( $c$ ) from a line meeting two of its chords, and ( $d$ ) from a line in one of its trisecant planes.

A prime of [6] meets the normal curve in six points; let us take three chords of the curve joining three pairs of these points. Any other chord of the curve meets the prime in a point through which there passes a plane meeting the first three chords\*. Projecting from a line in this plane we obtain in [4] a rational sextic with four concurrent chords. This proves the existence of the type III (I). Similarly, all the other curves  $C$  can be obtained by suitable projections†.

### *Surfaces without either a directrix line or a multiple generator*

173. Suppose that we have on  $\Omega$  a curve  $C$  of the type I (A). The chords of  $C$  form a three-dimensional locus, and, since the projection of  $C$  from a plane is a plane sextic with ten double points, there will be ten chords of  $C$  meeting an arbitrary plane. Thus the chords of  $C$  form a locus  $M_3^{10}$ .

\* Given three arbitrary lines in [5] there is just one plane through an arbitrary point of [5] meeting the three lines. If  $O$  is the point and  $a, b, c$  the lines the plane is the intersection of the three [4]'s  $Obc, Oca, Oab$ .

† For example, to obtain the curve of IV (C) and IV (D) which has two trisecants intersecting in a point of itself, we take two chords of the normal curve and a plane through a point of the curve meeting both these chords. Then projecting from a line in this plane we have the curve in [4] as required. And so on for other types.

$C$  is a quadruple curve on  $M_3^{10}$ , and there is also a double curve.

In particular a general plane of  $\Omega$  is met by ten chords of  $C$ ; hence  $C$  represents a rational sextic ruled surface in [3] whose double curve  $C_{10}$  is of order 10 and whose bitangent developable  $E_{10}$  is of class  $10^*$ .

Through any point of  $C$  there pass four chords which lie entirely upon  $\Omega$ ; these chords form a ruled surface  $R_2^{20}$  of order 20, the intersection of  $\Omega$  and  $M_3^{10}$ . They set up on  $C$  a (4, 4) correspondence of valency 2, so that † there are twenty-four points of  $C$  for which two of the four chords which lie on  $\Omega$  coincide. Also ‡ there are eight tangents of  $C$  lying on  $\Omega$ . If  $C'$  is a prime section of  $R_2^{20}$  we have a correspondence between  $C$  and  $C'$  for which

$$\alpha = 2, \quad \alpha' = 4, \quad p = 0, \quad \eta = 24, \quad \eta' = 8,$$

so that  $C'$  is of genus 3. Hence the double curve is  $C_{10}^3$  of genus 3 and the bitangent developable is  $E_{10}^3$  of genus  $3^*$ .

There are eight trisecant planes of  $C$  which lie entirely on  $\Omega$ ||; these are in fact four planes of each system. Hence  $C_{10}^3$  has four triple points and  $E_{10}^3$  four triple planes. These latter are tritangent planes of the ruled surface, each meeting it in three generators and a rational plane cubic.

Through each of the triple points of  $C_{10}^3$  there pass three lines meeting  $C_{10}^3$  in two further points; these are generators of the ruled surface.

This is the most general type of rational sextic ruled surface in [3].

174. Consider now, in [7], the most general rational sextic ruled surface  $F$ ; this surface contains  $\infty^1$  directrix cubic curves¶. It also contains  $\infty^3$  directrix quartic curves; these can all be obtained by means of primes through any two fixed generators¶. Taking a pencil of primes through the two generators we obtain a pencil of quartic curves; these curves have two common points—the remaining intersections of  $F$  with the [5] which is the base of the pencil of primes\*\*.

We have a system of  $\infty^3$  primes through the solid determined by any two fixed generators of  $F$ , and the different systems are projectively related to one another when we make those primes which contain the same directrix quartic of  $F$  correspond to one another. If we take three fixed pairs of generators of  $F$ , the [4]'s which contain the directrix quartics are determined as the intersections of corresponding primes of three projectively related triply infinite systems. A section by an arbitrary solid  $S$  gives three projectivities in  $S$ ; corresponding planes of these projectivities meet, in general, in a point, so that through a general point of  $S$ , and hence also through a general point of [7], there passes just one [4] which contains a directrix quartic of  $F$ .

\* § 33.  
|| § 35.

† § 13.  
¶ § 45.

‡ § 32.  
\*\* § 46.



Further, the chords of  $F$  are determined as the intersections of three corresponding [5]'s of the three projectively related systems. If we take a pencil of primes belonging to one system then we have corresponding pencils of primes belonging to the other two systems; the [5]'s which are the bases of the pencils meet in a line which is a chord of  $F$ . The  $\infty^4$  chords of  $F$  are all given in this way. Hence the points of intersection of an arbitrary solid  $S$  with chords of  $F$  are the points of concurrence of three corresponding lines in three projectivities between the planes of  $S$ . Such points lie on\* a curve  $\vartheta_6^3$  of order 6 and genus 3.

Hence the chords of  $F$  form a locus  $M_5^6$  of five dimensions and the sixth order; they meet an arbitrary solid  $S$  in a curve  $\vartheta_6^3$  of genus 3.

Suppose that we consider one of the  $\infty^5$  quintic curves of  $F$ ; it is contained in a space  $S_5$ . Then no chord of  $F$ , which is not also a chord of the quintic curve, can meet  $S_5$  except in a point of the quintic curve; for we cannot have a prime containing the quintic curve and also two generators. Hence the locus, of three dimensions, in which  $S_5$  meets the locus of five dimensions formed by the chords of  $F$ , is simply the locus formed by the chords of the quintic curve. Hence we again find † that the locus of chords of  $F$  is of the sixth order, and meets an arbitrary solid in a curve of genus 3.

This argument can also be applied to tangents of  $F$ ; the four-dimensional locus formed by the tangents of  $F$  meets  $S_5$  in the surface formed by the tangents of the quintic curve. Hence the tangents of  $F$  form an  $M_4^8$  of the eighth order; the section of this by an arbitrary [4] is a rational curve.

A directrix cubic  $\Delta$  of  $F$  is determined by a prime through any three fixed generators, so that the solids  $K$  containing the curves  $\Delta$  are determined as the intersections of four corresponding primes belonging to four projectively related pencils of primes. The four-dimensional locus formed by the solids  $K$  meets an arbitrary [4] in a curve which is the locus of the intersection of corresponding solids of four projectively related pencils of solids, and this is a rational quartic curve ‡. Hence the solids  $K$  form a locus  $M_4^4$  of the fourth order, and an arbitrary solid  $S$  will be met by four of them.

The [4] containing any directrix quartic of  $F$  is the base of a doubly infinite system of primes, each prime of the system containing two generators of  $F$ . If any directrix cubic  $\Delta$  is taken the directrix quartic will meet it in one point; so that the primes of the doubly infinite system meet the solid  $K$  containing  $\Delta$  in the star of planes which projects the chords of  $\Delta$  from a fixed point of itself. Now the star of planes which projects the chords of  $\Delta$  from any point of itself is projectively related to the star of planes which projects the chords of  $\Delta$  from any other point of itself, planes of the

\* Schur, *Math. Ann.* 18 (1881), 16.

† Cf. § 95.

‡ Veronese, *Math. Ann.* 19 (1882), 219.

two stars corresponding when they contain the same chord of  $\Delta$ . Hence the system of primes containing any directrix quartic of  $F$  is projectively related to the system of primes containing any other directrix quartic of  $F$ , primes of the two systems corresponding when they contain the same pair of generators of  $F$ .

Hence, taking any four fixed directrix quartics of  $F$ , we may say that the solids  $K'$  containing the pairs of generators of  $F$  are determined as the intersections of corresponding primes of four projectively related doubly infinite systems. If we consider the section by an arbitrary [4] we obtain four projectively related systems of solids, each system having a line as base. These are known to generate, by means of the intersections of corresponding solids, a sextic surface whose curve sections are of genus 3, and which has ten lines on it\*; there being ten sets of corresponding solids which have a line in common instead of a point.

The solids  $K'$  thus form a locus  $M_5^6$  of five dimensions and the sixth order; it is of course the same locus  $M_5^6$  as that formed by the chords of  $F$ . Of the  $\infty^3$  solids  $K'$  there are ten which meet any given [4] in lines.

Let us now project  $F$  from an arbitrary solid  $S$ , which does not meet it, on to a [3]  $\Sigma$ ; we obtain a surface  $f$ . The chords of  $F$  meet  $S$  in the points of a curve  $\mathfrak{D}_6^3$ . A prime through  $S$  meets  $F$  in a rational normal sextic curve, of which there are ten chords meeting  $S$ ; hence those chords of  $F$  which meet  $S$  meet  $F$  in the points of a curve  $C_{20}$  of order 20. Then, on projection,  $C_{20}$  becomes the double curve  $C_{10}^3$  of  $f$ ;  $C_{10}^3$  is of genus 3 since it is in (1, 1) correspondence with  $\mathfrak{D}_6^3$ .

There are eight tangents of  $F$  meeting  $S$ , so that the (1, 2) correspondence between  $\mathfrak{D}_6^3$  and  $C_{20}$  has eight branch points, and Zeuthen's formula shews that  $C_{20}$  is a curve  $C_{20}^9$  of genus 9. Since a [5] determined by  $S$  and a generator of  $F$  meets  $F$  again in four points† the curve  $C_{20}^9$  must meet each generator of  $F$  in four points; it must therefore have twelve double points‡. These will lie three in each of four planes meeting  $S$  in lines, and the double curve  $C_{10}^3$  of  $f$  will thus have four triple points. If we apply Zeuthen's formula to the (1, 4) correspondence between a directrix curve of  $F$  and  $C_{20}^9$  we find that there are twenty-four generators of  $F$  touching  $C_{20}^9$ ; thus there must be twenty-four generators of  $f$  touching its double curve.

The class of the bitangent developable of  $f$  is the number of primes which contain a given [4] through  $S$  and also two generators of  $F$ ; it is therefore the number of solids  $K'$  which meet a given [4] through  $S$  in lines. We have seen that this number is ten, so that the bitangent developable of  $f$  must be of class 10. It is of genus 3, since its planes are in (1, 1)

\* Veronese, *Math. Ann.* 19 (1882), 232. White, *Proc. Camb. Phil. Soc.* 21 (1923), 223.

† § 46.

‡ § 17.

correspondence with the points of  $\mathfrak{D}_6^3$ ; we have then a bitangent developable  $E_{10}^3$ . The number of tritangent planes of  $f$  is the number of primes through  $S$  which contain three generators of  $F$ ; such a prime meets  $F$  further in a directrix cubic  $\Delta$ . Then the solid  $K$  containing  $\Delta$  will meet  $S$ , and we have already seen that there are four such solids  $K$ . Hence  $f$  has four tritangent planes.

Since  $F$  is generated by a (1, 1) correspondence between two directrix cubics,  $f$  is generated by taking two twisted cubics in  $\Sigma$  and placing them in (1, 1) correspondence. We can take either or both of these cubics to be a rational plane cubic instead of a twisted cubic.

*Note 1.* Algebraically: take two cubics in [7]; let the coordinates of the points of one be  $(\theta^3, \theta^2, \theta, 1, 0, 0, 0)$  and the coordinates of the points of the other  $(0, 0, 0, 0, \theta^3, \theta^2, \theta, 1)$ . Then the general rational sextic ruled surface is given in terms of the parameters  $\lambda$  and  $\theta$  by  $(\theta^3, \theta^2, \theta, 1, \lambda\theta^3, \lambda\theta^2, \lambda\theta, \lambda)$ . Its equations are

$$\frac{x_0}{x_1} = \frac{x_1}{x_2} = \frac{x_2}{x_3} = \frac{x_4}{x_5} = \frac{x_5}{x_6} = \frac{x_6}{x_7},$$

and it has  $\infty^1$  directrix cubics  $\Delta$  given by  $\lambda = \text{const}$ . The  $\infty^1$  solids  $K$  containing these cubics generate the  $M_4^4$   $\frac{x_0}{x_4} = \frac{x_1}{x_5} = \frac{x_2}{x_6} = \frac{x_3}{x_7}$ , while the chords of  $F$  generate the  $M_5^6$

$$\left\| \begin{array}{cccc} x_0 & x_1 & x_4 & x_5 \\ x_1 & x_2 & x_5 & x_6 \\ x_2 & x_3 & x_6 & x_7 \end{array} \right\| = 0.$$

The equations of the solid  $K'$  determined by the generators  $\theta = \theta_1$  and  $\theta = \theta_2$  are

$$\begin{aligned} x_0 - (\theta_1 + \theta_2)x_1 + \theta_1\theta_2x_2 &= x_1 - (\theta_1 + \theta_2)x_2 + \theta_1\theta_2x_3 \\ &= x_4 - (\theta_1 + \theta_2)x_5 + \theta_1\theta_2x_6 = x_5 - (\theta_1 + \theta_2)x_6 + \theta_1\theta_2x_7 = 0, \end{aligned}$$

and the equations of the tangent solid along the generator  $\theta$  are

$$x_0 - 2\theta x_1 + \theta^2 x_2 = x_1 - 2\theta x_2 + \theta^2 x_3 = x_4 - 2\theta x_5 + \theta^2 x_6 = x_5 - 2\theta x_6 + \theta^2 x_7 = 0.$$

*Note 2.* It is, of course, essential to our work that we should investigate all the possible positions which the solid  $S$  may occupy in regard to  $F$ , and the curve  $\mathfrak{D}_6^3$  in which  $S$  meets  $M_5^6$  will break up in many different ways. When we have chosen  $S$  to occupy a certain position we shall state the orders and genera of the component curves of  $\mathfrak{D}_6^3$ , and also the number of their mutual intersections; we shall not give the proofs of these statements.

The way in which  $\mathfrak{D}_6^3$  can break up into separate curves can be seen by making use of the properties of the cubic surface. When  $\mathfrak{D}_6^3$  is generated by four projective stars of planes with vertices 1, 2, 3, 4 it lies on both the cubic surfaces  $F_3$  and  $F_4$ , where  $F_3$  is the cubic surface generated by the stars 4, 1, 2 and  $F_4$  the cubic surface generated by the stars 1, 2, 3. Now, if two corresponding lines of the stars 1 and 2 intersect, this intersection is on both  $F_3$  and  $F_4$ , but is not (in general) on  $\mathfrak{D}_6^3$ . Hence  $\mathfrak{D}_6^3$  is the residual intersection of two cubic surfaces with a cubic curve in common.

Suppose now that we represent one of the cubic surfaces,  $F_3$  say, in the usual way upon a plane. There are six sets of corresponding planes of the stars 1, 2, 3 which meet in lines  $a_1, a_2, a_3, a_4, a_5, a_6$  instead of only in points; these lines are represented by *fundamental points*  $A_1, A_2, A_3, A_4, A_5, A_6$  in the plane and form one-half of a double-six on  $F_3$ . The cubic curve generated by the stars 1 and 2 has these six lines as chords, it is represented on the plane by a quintic with nodes at the fundamental points  $A$ . Since the complete intersection of  $F_3$  with another cubic surface is represented on the plane by a curve of order 9 with triple points at the points  $A$ , the curve  $\mathfrak{D}_6^3$  must be represented by a quartic curve through the points  $A$ . Hence, in order to find the possible ways in which  $\mathfrak{D}_6^3$  may break up, we have simply to consider how a plane quartic, with six fixed points, may break up.

Suppose, for example, that  $\mathfrak{D}_6^3$  breaks up into a line and a quintic curve. The line may be one of the six lines  $a$ , or one of the six lines  $b$  belonging to the other half of the double-six, or one of the fifteen lines  $c$  not belonging to the double-six. If it is a line  $a$  we have in the plane a quartic with a double point at a point  $A$ ; hence, on  $F_3$ , a curve  $\mathfrak{D}_5^2$  with  $a$  as a chord. If it is a line  $b$  (represented by a conic through five points  $A$ ) we have in the plane a conic through one of the points  $A$ ; hence, on  $F_3$ , a curve  $\mathfrak{D}_5^0$  with  $b$  as its quadriseccant. If it is a line  $c$  (represented by a line joining two points  $A$ ) we have in the plane a cubic curve through four points  $A$ ; hence, on  $F_3$ , a curve  $\mathfrak{D}_5^1$  with  $c$  for a trisecant. And so on if  $\mathfrak{D}_6^3$  contains as part of itself a conic or cubic curve, or if it breaks up into more than two parts.

175. If we consider a rational normal sextic in [6], and take any  $g_2^1$  on it, the chords joining the pairs of points form a rational quintic ruled surface with a directrix conic  $\Gamma$ ,  $\Gamma$  not meeting the curve. If we then project from a point in the plane of  $\Gamma$  on to a [5] we obtain a rational sextic  $C$ , and a  $g_2^1$  giving chords of  $C$  which are generators of a rational quintic ruled surface with a double line. There are in [5]  $\infty^{20}$  quadric fourfolds; of these there are  $\infty^7$  containing  $C$ , and of these latter  $\infty^4$  contain the quintic ruled surface. Let us take one of these last quadrics to be  $\Omega$ .

The generators of the quintic surface are projected from the double line by planes forming a three-dimensional cubic line-cone; since it contains the quintic surface its residual intersection with  $\Omega$  must be a plane. Hence  $\Omega$  contains a plane joining the double line to a generator of the surface. Any general plane of  $\Omega$  of the same system as this meets the surface in two points, while a plane of the opposite system meets the surface in three points.

The chords of  $C$  which lie on  $\Omega$  form the quintic ruled surface together with a ruled surface  $R_2^{15}$ ; this latter meets planes of one system of  $\Omega$  in eight points and planes of the other system in seven points—every plane of  $\Omega$  being met by ten chords of  $C$ .  $C$  is a triple curve on  $R_2^{15}$ .

The quintic ruled surface gives a (1, 1) correspondence on  $C$ , while the surface  $R_2^{15}$  gives a (3, 3) correspondence, both these correspondences being

symmetrical. It follows, on using Brill's formula (cf. §§ 14, 15), that  $R_2^5$  and  $R_3^{15}$  have three common generators.

Of the eight tangents of  $C$  which lie on  $\Omega$  two belong to the quintic ruled surface and the remaining six to  $R_2^{15}$ . There are twelve points of  $C$  at which two generators of  $R_2^{15}$  coincide. Hence, if  $C'$  is a prime section of  $R_2^{15}$  we have, for the correspondence between  $C$  and  $C'$ ,

$$\alpha = 2, \quad \alpha' = 3, \quad p = 0, \quad \eta = 12, \quad \eta' = 6,$$

so that  $C'$  is an elliptic curve.

Suppose, for definiteness, that the plane of  $\Omega$  which contains the double line and one of the generators of the quintic surface is a  $\rho$ -plane. Then the points of  $C$  represent the generators of a ruled surface in [3] whose double curve is  $C_2 + C_3^1$  and bitangent developable  $E_3^0 + E_7^1$ . The fact that the two ruled surfaces on  $\Omega$  have three common generators shews that  $C_2$  and  $C_3^1$  have three common points, while  $E_3^0$  and  $E_7^1$  have three common planes.

Take any one of the four  $\rho$ -planes of  $\Omega$  which are trisecant to  $C$ , meeting  $C$  in  $P, Q, R$  say. Assume for the moment that no one of  $QR, RP, PQ$  is a generator of the quintic surface. Then through each of  $P, Q, R$  there pass generators of this surface which all meet the double line; and the plane  $PQR$ , together with the  $\rho$ -plane through the double line, determines a [4] meeting the quintic surface in the double line and four generators, which is impossible. Hence one of  $QR, RP, PQ$  must be a generator of the quintic surface, and similarly for each of the other trisecant  $\rho$ -planes. Hence  $E_7^1$  has four double planes which are also planes of  $E_3^0$ ; the two developables having three other planes in common.

Also it will be seen when this surface is obtained by projection that  $C_2$  passes through two double points of  $C_3^1$  and meets it in three further points;  $C_3^1$  has also two triple points.

We might, on the other hand, have taken the plane of  $\Omega$  which contains the double line and a generator of the quintic ruled surface to be a  $\omega$ -plane. We should then have obtained in [3] a ruled surface with a double curve  $C_3^0 + C_7^1$  and bitangent developable  $E_2 + E_8^1$ .  $C_7^1$  has four double points through which  $C_3^0$  passes, the curves having three other intersections.  $E_8^1$  has two double planes which are also planes of  $E_2$ , the two developables having three other planes in common;  $E_8^1$  has also two triple planes.

176. On the normal surface  $F$  in [7] there are  $\infty^3$  quartics  $E$ . An involution of pairs of generators of  $F$  gives rise to a  $g_2^1$  on each of these quartics, and thus to a cubic ruled surface through each of them. Let us then choose  $S$  to pass through the directrix line  $\lambda$  of a cubic ruled surface formed by chords of one of these quartics  $E$ . There are two tangents of  $E$  belonging to this cubic surface.

A prime through  $S$  meets  $F$  in a rational normal sextic, ten of whose chords meet  $S$ ; but two of these chords are generators of the cubic ruled surface. Hence the chords of  $F$  which meet  $S$  meet  $F$  in  $E$  and a curve  $C_{16}$  meeting each generator of  $F$  in three points.  $E$  and  $C_{16}$  have ten intersections\*.

$S$  meets  $M_5^6$  in  $\lambda$  and an elliptic quintic  $\vartheta_5^1$  of which  $\lambda$  is a trisecant. Through the three intersections of  $\lambda$  and  $\vartheta_5^1$  there pass chords of  $F$  which are chords both of  $E$  and  $C_{16}$ ; thus six of the intersections of  $C_{16}$  with  $E$  are accounted for. Let  $A$  be another intersection of these two curves. Through  $A$  there passes a chord  $AB$  of  $E$  meeting  $\lambda$  and a chord  $AC$  of  $C_{16}$  meeting  $\vartheta_5^1$ . Then the chord  $BC$  of  $F$  must also meet  $\vartheta_5^1$ , so that  $C_{16}$  passes through  $B$  and has a double point at  $C$ . There remains a further pair of intersections of  $E$  and  $C_{16}$ , these are associated in the same way with a double point on  $C_{16}$ . There are six tangents of  $C_{16}$  which meet  $\vartheta_5^1$ ; hence it is of genus 4. It must then have eight double points. Two of these are already accounted for; the six others lie three in each of two planes which meet  $S$  in lines.

Thus when we project from  $S$  on to  $\Sigma$  we obtain a ruled surface  $f$  whose double curve is  $C_2 + C_8^1$ ;  $C_2$  being the projection of  $E$  and  $C_8^1$  the projection of  $C_{16}$ .  $C_8^1$  has three simple points and two double points on  $C_2$ , and has also two triple points.

Since  $F$  can be generated by means of a (1, 1) correspondence between  $E$  and a directrix cubic  $\Delta$  with a united point, to generate  $f$  we take in  $\Sigma$  a conic and a twisted cubic in (1, 2) correspondence with a united point. The pairs of points of the cubic which correspond to the points of the conic form a  $g_2^1$ , so that their joins form a regulus; hence the planes of the pairs of generators which intersect in the points of the conic will form † a cubic developable  $E_3^0$  (since there is a united point). This is part of the bitangent developable of  $f$ . The other part is an  $E_7^1$ ; it is formed by the planes of pairs of generators which intersect in points of the double curve  $C_8^1$ .

177. A prime section of  $F$  is a rational normal sextic; an involution of pairs of generators of  $F$  gives a  $g_2^1$  on this sextic, and the chords joining the pairs of this  $g_2^1$  form a rational quintic ruled surface with a directrix conic  $\Gamma$ ,  $\Gamma$  not meeting the sextic. Choose then the solid  $S$  to pass through the plane of  $\Gamma$ .

A prime through  $S$  meets  $F$  in a rational normal sextic, ten of whose chords meet  $S$ ; but three of these belong to the quintic ruled surface. Thus the chords of  $F$  meeting  $S$  meet  $F$  in the prime section  $C_8$  and a curve  $C_{14}$  meeting each generator of  $F$  in three points.  $C_8$  and  $C_{14}$  meet in fourteen points.

\* Cf. the last footnote to § 97.

† § 22.

$S$  meets  $M_5^0$  in  $\Gamma$  and an elliptic quartic  $\mathfrak{D}_4^1$  having three intersections with  $\Gamma$ . Through these three points pass chords of  $F$  which are chords both of  $C_6$  and  $C_{14}$ ; this accounts for six intersections of  $C_6$  and  $C_{14}$ . The other eight fall into four pairs, and with each pair is associated a double point of  $C_{14}$ . Since  $C_{14}$  has six tangents which meet  $\mathfrak{D}_4^1$  it is of genus 4, and this shews that it has exactly four double points.

On projection from  $S$  on to  $\Sigma$  we obtain a ruled surface  $f$  whose double curve is  $C_3^0 + C_7^1$ ;  $C_3^0$  is the projection of  $C_6$ , while  $C_7^1$  is the projection of  $C_{14}$ .  $C_3^0$  passes through the four double points of  $C_7^1$  and also meets it in three other points.

Since  $F$  can be generated by a (1, 1) correspondence between  $C_6$  and a directrix cubic  $\Delta$ , there being three united points,  $f$  can be generated by two twisted cubics in (1, 2) correspondence with three united points. The pairs of points of the second cubic corresponding to the points of the first form a  $g_2^1$  and their joins form a regulus, hence the planes of the pairs of generators which pass through the points of the first cubic are (because of the united points) the tangent planes of a quadric cone  $E_2$ . This is part of the bitangent developable of the surface; the remaining part is an  $E_3^1$ .

178. On the rational normal sextic curve in [6] there are  $\infty^2$  involutions  $g_2^1$ . Each of these gives rise to a quintic ruled surface with a directrix conic; and these conics lie in  $\infty^2$  planes. Let us suppose that two of these planes intersect in a point not on either of their conics. Projecting from such a point on to a [5] we obtain a rational sextic  $C$  and two quintic ruled surfaces containing it, each with a double line. There is a quadric  $\Omega$  containing these two ruled surfaces.

The chords of  $C$  which lie on  $\Omega$  form two quintic ruled surfaces  $R_2^5$  and  $S_2^5$ , and a ruled surface  $R_2^{10}$  on which  $C$  is a double curve.  $R_2^5$  and  $S_2^5$  have one common generator;  $R_2^{10}$  has two generators in common with each of  $R_2^5$  and  $S_2^5$ . The prime sections of  $R_2^{10}$  are rational curves.

For either of the quintic surfaces there is a plane of  $\Omega$  passing through its double line and containing one of its generators.

First, suppose that both these planes are  $\rho$ -planes. Then we have in [3] a ruled surface whose double curve is  $C_2 + D_2 + C_6^0$  and bitangent developable  $E_3^0 + F_3^0 + E_4^0$ . Each of the four  $\rho$ -planes of  $\Omega$  which is trisecant to  $C$  must contain a generator of  $R_2^5$  and a generator of  $S_2^5$ . Hence there are four planes common to all the developables  $E_3^0$ ,  $F_3^0$  and  $E_4^0$ ;  $E_3^0$  and  $F_3^0$  have a further plane in common, while  $E_4^0$  has two further planes in common with each of them. It will be seen that  $C_2$  passes through two double points and two simple points of  $C_6^0$ , as also does  $D_2$ ;  $C_6^0$  has four double points, while  $C_2$  and  $D_2$  have one intersection.

If, on the other hand, both the planes of  $\Omega$  are  $\pi$ -planes we have a ruled surface in [3] whose double curve is  $C_3^0 + D_3^0 + C_4^0$  and bitangent developable  $E_2 + F_2 + E_6^0$ . There are four points common to all the curves  $C_3^0$ ,  $D_3^0$  and  $C_4^0$ ;  $C_3^0$  and  $D_3^0$  have one other intersection, while  $C_4^0$  meets each of them in two further points.  $E_6^0$  has four double planes; two of these are also planes of  $E_2$  and the other two of  $F_2$ ;  $E_6^0$  has two further planes in common with  $E_2$  and two with  $F_2$ , while  $E_2$  and  $F_2$  have a common plane.

But further: one of the planes of  $\Omega$  may be a  $\rho$ -plane, while the other is a  $\pi$ -plane. We then have in [3] a ruled surface whose double curve is  $C_2 + C_3^0 + C_5^0$  and bitangent developable  $E_3^0 + E_2 + E_5^0$ .

179. Take two directrix quartics  $E$  and  $E'$  on the normal surface  $F$  in [7]; they have two intersections. Through these intersections there pass two generators of  $F$ ; let us take two involutions of generators which both include this pair. Then we have two cubic ruled surfaces each with a directrix line; one line  $\lambda$  lies in the [4] containing  $E$ , while the other  $\lambda'$  lies in the [4] containing  $E'$ . Both  $\lambda$  and  $\lambda'$  meet the chord of  $F$  which joins the intersections of  $E$  and  $E'$ , and we choose the involutions so that they meet this chord in the same point\*.

Take a solid  $S$  containing  $\lambda$  and  $\lambda'$  and project from  $S$  on to  $\Sigma$ . A prime through  $S$  meets  $F$  in a rational normal sextic; there are ten chords of this curve meeting  $S$ , including two chords of  $E$  and two chords of  $E'$ ; thus the chords of  $F$  which meet  $S$  meet  $F$  in  $E$ ,  $E'$  and a curve  $C_{12}$  meeting every generator in two points.  $C_{12}$  meets each of  $E$  and  $E'$  in eight points.

$S$  meets  $M_5^0$  in  $\lambda$ ,  $\lambda'$  and a rational quartic  $\mathfrak{D}_4^0$  having  $\lambda$  and  $\lambda'$  as chords. Through the intersections of  $\lambda$  and  $\mathfrak{D}_4^0$  there pass lines which are chords both of  $E$  and  $C_{12}$ ; the other four intersections of  $E$  and  $C_{12}$  fall into two pairs with each of which is associated a double point of  $C_{12}$ . We can thus obtain four double points on  $C_{12}$ . There are four tangents of  $C_{12}$  meeting  $\mathfrak{D}_4^0$ , and the (1, 2) correspondence between  $\mathfrak{D}_4^0$  and  $C_{12}$  shews that  $C_{12}$  is elliptic, whereupon Segre's formula shews that it has exactly four double points.

Thus we obtain in  $\Sigma$  a ruled surface  $f$  with a double curve  $C_2 + D_2 + C_6^0$ .  $C_2$  and  $D_2$  have one intersection;  $C_6^0$  has four double points, two of which are on  $C_2$  and the other two on  $D_2$ , while it has two simple intersections with each of  $C_2$  and  $D_2$ .

Since  $F$  can be generated by placing  $E$  and  $E'$  in (1, 1) correspondence with two united points we can generate  $f$  by taking two conics  $C_2$  and  $D_2$  with one intersection, placing them in (2, 2) correspondence with a doubly united point. The lines joining the pairs of points of one conic which correspond to the points of the other touch a conic; hence the planes joining

\* See the footnotes to § 131.



the points of either conic to the pairs of points on the other which correspond to them form a developable of the third class which is part of the bitangent developable of the surface. We thus have a bitangent developable  $E_3^0 + F_3^0 + E_4^0$ .

**180.** Two chords of  $F$  determine a [3]; through this [3] there pass  $\infty^3$  primes meeting  $F$  in rational normal sextics through four fixed points. On any one of these curves we can regard the two chords as determining an involution, we thus have a quintic ruled surface containing the chords. We can choose the prime so that the directrix conic of the quintic surface meets these chords in assigned points.

Now take a directrix quartic, three of its chords  $l, m, n$  and their transversal, meeting them in  $L, M, N$ . We can regard  $l$  and  $m$  as determining an involution of pairs of generators on  $F$  and a quintic ruled surface whose directrix conic  $\Gamma$  passes through  $L$  and  $M$ ; similarly, we have another surface whose directrix conic  $\Gamma'$  passes through  $L$  and  $N$ , the two surfaces having a common generator through  $L$ . Let  $S$  be the solid containing the planes of  $\Gamma$  and  $\Gamma'$ .

A prime through  $S$  meets  $F$  in a rational sextic curve, ten of whose chords meet  $S$ ; of these ten chords three meet  $\Gamma$  and three meet  $\Gamma'$ . Those chords of  $F$  which meet  $S$  meet  $F$  in two prime sections and a curve  $C_8$  meeting every generator in two points and each prime section in eight points.

$S$  meets  $M_5^6$  in  $\Gamma, \Gamma'$  and a conic  $\vartheta_2$  meeting each of  $\Gamma$  and  $\Gamma'$  in two points. There are four tangents of  $C_8$  meeting  $\vartheta_2$ , so that  $C_8$  is elliptic and has no double points. The chords of  $F$  through the intersections of  $\Gamma$  and  $\vartheta_2$  are common chords of  $C_8$  and one of the prime sections; there are four other intersections of  $C_8$  with this section to be accounted for. Similarly we obtain four intersections of  $C_8$  with the other prime section. Also there are four points common to the two prime sections other than the two on the chord through  $L$ . These three sets of four points are associated with one another; they fall into four triads, each triad includes a point of each set and projects into a single point of  $\Sigma$ .

On projection we have in  $\Sigma$  a ruled surface  $f$  whose double curve is  $C_3^0 + D_3^0 + C_4^0$ ; there are four points common to all the curves  $C_3^0, D_3^0$  and  $C_4^0$ ;  $C_3^0$  and  $D_3^0$  have one further intersection, while  $C_4^0$  meets each of them in two further points.

To generate this surface  $f$  we take two twisted cubics  $C_3^0$  and  $D_3^0$  with five common points, placing them in (2, 2) correspondence with four ordinary united points and one doubly united point. The joins of the pairs of points of one cubic which correspond to the points of the other generate a rational quartic ruled surface, so that the planes joining the points of either cubic to the pairs of points of the other which correspond

to them touch a quadric cone (because of the five united points). We have a bitangent developable  $E_2 + F_2 + E_8^0$ .

**181.** Take now a prime section  $C_6^0$  of  $F$ ; an involution of pairs of generators gives chords of  $C_6^0$  forming a quintic ruled surface with a directrix conic  $\Gamma$ . Take any point  $A$  of  $\Gamma$ ; the chord of  $F$  through  $A$  meets it in two points through which pass  $\infty^1$  directrix quartics. Take one of these quartics  $E$  and an involution of pairs of generators of  $F$  such that the resulting cubic ruled surface formed by chords of  $E$  has its directrix line  $\lambda$  passing through  $A$ , the chord of  $C_6^0$  through  $A$  being common to this surface and the quintic surface. Then choose  $S$  to be the solid containing  $\lambda$  and  $\Gamma$ .

The chords of  $F$  which meet  $S$  meet  $F$  in  $E$ ,  $C_6^0$  and a curve  $C_{10}$  meeting every generator of  $F$  in two points.  $S$  meets  $M_5^6$  in  $\Gamma$ ,  $\lambda$  and a twisted cubic  $\vartheta_3^0$  which meets each of  $\Gamma$  and  $\lambda$  in two points. There are four tangents of  $C_{10}$  meeting  $\vartheta_3^0$  so that  $C_{10}$  is elliptic and has consequently two double points.

$C_{10}$  meets  $E$  in six points and  $C_6^0$  in ten points; four of the intersections with  $E$  are accounted for by the chords of  $F$  through the points common to  $\vartheta_3^0$  and  $\lambda$  and four of the intersections with  $C_6^0$  by the chords of  $F$  through the points common to  $\vartheta_3^0$  and  $\Gamma$ . Four further intersections with  $C_6^0$  are associated in pairs with the two double points of  $C_{10}$ ; the two remaining ones are associated with the two remaining intersections with  $E$  and with the two intersections of  $E$  and  $C_6^0$  which do not lie on the chord through  $A$ .

Hence on projection we have in  $\Sigma$  a ruled surface  $f$  whose double curve is  $C_2 + C_3^0 + C_5^0$ .  $C_5^0$  has two double points which lie on  $C_3^0$ ; it meets  $C_3^0$  in four further points through two of which  $C_2$  also passes, while  $C_2$  and  $C_3^0$  have one other intersection.

Since  $F$  is generated by a (1, 1) correspondence between  $C_6^0$  and  $E$  with four united points, to generate  $f$  we take in  $\Sigma$  a conic  $C_2$  and a twisted cubic  $C_3^0$  with three intersections, placing them in (2, 2) correspondence with two ordinary united points and one doubly united point. The joins of the pairs of points of the conic which correspond to the points of the cubic touch another conic; we thus obtain a quadric cone as part of the bitangent developable. The points of the conic give pairs of points of the cubic whose joins form a quartic ruled surface, and we thus obtain a developable of the third class. Thus the bitangent developable of the surface is  $E_3^0 + E_2 + E_5^0$ .

This surface is not strictly self-dual; the planes of the quadric cone are not formed by the generators which intersect in the points of  $C_2$  but by those which intersect in the points of  $C_3^0$ .

**182.** All the ruled surfaces in [3] so far obtained are such that their generators are represented by the points of a curve  $C$  on  $\Omega$  which belongs to the type I (A). Let us now suppose that we have a curve  $C$  of the

type II (A); since it lies in a [4] the generators of the corresponding ruled surface in [3] belong to a linear complex. The chords of  $C$  form an  $M_3^{10}$ , so that any plane of  $\Omega$  meets the [4] containing  $C$  in a line which is met by ten chords of  $C$ . It is shewn precisely as in § 173 that the chords of  $C$  lying on  $\Omega$  form a ruled surface  $R_2^{20}$  whose prime sections are of genus 3; hence the ruled surface in [3] has a double curve  $C_{10}^3$  and a bitangent developable  $E_{10}^3$ .

$C$  has four trisecants\*: these also lie on  $\Omega$ . Thus  $C_{10}^3$  has four triple points; through each of these points there pass three generators of the surface, and these lie in a plane which is a triple plane of  $E_{10}^3$ .

Since any surface whose generators belong to a linear complex is necessarily self-dual there is no need to seek for the other types of surfaces  $f$  when  $C$  is of the type II (A).

**183.** Given a rational sextic  $C$  in [5] we can always make some of the  $\infty^7$  quadrics containing it contain also one of its quadrisecant planes. Then the curve  $C$  represents a ruled surface in [3] for which either four generators pass through a point or else four generators lie in a plane. The quadrisecant plane of  $C$  counts for four among the trisecant planes of  $C$  which lie on  $\Omega$ ; we shall have this quadrisecant plane belonging to one system and four trisecant planes belonging to the opposite system.

Take four points, one on each of four generators of the normal surface  $F$ , and take  $S$  to meet the solid determined by these four points in a plane. If we project from  $S$  we have in  $\Sigma$  a surface  $f$  with four concurrent generators. The chords of  $F$  which meet  $S$  meet  $F$  in a curve  $C_{20}^9$  with triple points at each of the four selected points, so that the surface has a double curve  $C_{10}^3$  with a sextuple point. To generate  $f$  take a plane quartic with a triple point and a twisted cubic passing through the triple point and having one other intersection with the quartic; place them in (1, 1) correspondence with this last intersection as a united point.

To obtain the surface with a quadritangent plane we must project the normal surface with a directrix conic  $\Gamma$ . The chords of this surface also form an  $M_5^6$ , meeting an arbitrary solid  $S$  in a curve  $\mathfrak{D}_6^3$ , while the tangents form an  $M_4^8$ †. The quadritangent plane of the surface  $f$  is the intersection

\* If we take a rational normal sextic in [6] there are four of its trisecant planes meeting an arbitrary line. If, for instance, the curve is given by

$$x_0 : x_1 : x_2 : x_3 : x_4 : x_5 : x_6 = \theta^6 : \theta^5 : \theta^4 : \theta^3 : \theta^2 : \theta : 1$$

the trisecant planes form the quartic primal

$$\begin{vmatrix} x_0 & x_1 & x_2 & x_3 \\ x_1 & x_2 & x_3 & x_4 \\ x_2 & x_3 & x_4 & x_5 \\ x_3 & x_4 & x_5 & x_6 \end{vmatrix} = 0.$$

† If we set up a correspondence between the points  $B$  and  $C$  of  $\Gamma$ , the points  $B$  and  $C$  corresponding when the [5] containing  $S$  and the tangent of a given directrix

of  $\Sigma$  with the [6]  $ST$ ; to generate  $f$  we take a conic and a rational skew quartic in (1, 1) correspondence.

Since no two quadrisecant planes of a rational sextic in [5] can intersect we cannot have two quadrisecant planes of the same system on  $\Omega$ . But it is possible to have two quadrisecant planes on  $\Omega$  of different systems; then there will be no trisecant planes of  $C$  on  $\Omega$ ; the ruled surface in [3] has four concurrent generators and four coplanar generators. This is also obtained by projection from the normal surface with a directrix conic  $\Gamma$ .

*Surfaces with a multiple generator but without a directrix line*

**184.** Suppose now that we have a curve  $C$  on  $\Omega$  of the type I (B);  $C$  lies in [5] and has a double point  $P$ . Correspondingly we shall have in [3] a rational sextic ruled surface with a double generator  $G$ . The chords of  $C$  form an  $M_3^9$  on which  $C$  is a quadruple curve and  $P$  a sextuple point; the quartic cone formed by joining  $P$  to the other points of  $C$  is a double surface on  $M_3^9$ . Any plane of  $\Omega$  is met by nine chords of  $C$ ; hence, apart from the double generator, the surface in [3] has a double curve of the ninth order and a bitangent developable of the ninth class.

The chords of  $C$  lying on  $\Omega$  form a ruled surface  $R_2^{18}$  of order 18, the intersection of  $\Omega$  and  $M_3^9$ . Through any point of  $C$  there pass four generators of  $R_2^{18}$ , and there are twenty-four points of  $C$  at which two of these coincide. These, however, include the point  $P$  counted four times\*. Also there are eight tangents of  $C$  lying on  $\Omega$ . Hence, for the correspondence between  $C$  and a prime section  $C'$  of  $R_2^{18}$ , we have

$$\alpha = 2, \quad \alpha' = 4, \quad p = 0, \quad \eta = 20, \quad \eta' = 8,$$

and Zeuthen's formula gives  $p' = 2$ . Hence the ruled surface has a double curve  $G + G_9^2$  and a bitangent developable  $G + E_9^2$ .

The plane of the two tangents of  $C$  at  $P$  belongs to  $M_3^9$ †, so that it meets  $\Omega$  in two lines which are generators of  $R_2^{18}$ . The planes  $\omega$  through these lines represent intersections of  $C_9^2$  and  $G$ , while the planes  $\rho$  through them represent planes of  $E_9^2$  passing through  $G$ .

There are two chords  $PQ, PR$  of  $C$  which pass through  $P$  and lie on  $\Omega$ .

quartic  $E$ , at the point on the generator of  $F$  through  $B$ , meets the plane of  $\Gamma$  on the tangent at  $C$ , we have a (6, 2) correspondence. The eight coincidences give eight tangent solids of  $F$  which meet  $S$ . Cf. also the footnote to § 127.

\* Cf. § 265 below.

† This is seen at once when  $M_3^9$  is regarded as the projection of the  $M_3^{10}$  formed by the chords of a rational normal sextic. The tangent [3] of  $M_3^{10}$  at a point  $P$  contains the tangents of the normal curve at the two points where the chord through  $P$  meets it;  $M_3^{10}$  has the same tangent [3] at all points of a chord of the curve. The coordinates of a point of  $M_3^{10}$  are given parametrically by  $(\theta^6 + \kappa\phi^6, \theta^5 + \kappa\phi^5, \theta^4 + \kappa\phi^4, \theta^3 + \kappa\phi^3, \theta^2 + \kappa\phi^2, \theta + \kappa\phi, 1 + \kappa)$ ; and it is easily seen that the tangent solid at this point does not depend on  $\kappa$ .

The planes of  $\Omega$  through these lines are to be regarded as trisecant planes of  $C$ ; thus we have two double points of  $C_9^2$  on  $G$  and two double planes of  $E_9^2$  through  $G$ . There will be two other trisecant planes of  $C$  belonging to each system of planes on  $\Omega$ ; hence  $C_9^2$  has two triple points, and  $E_9^2$  has two triple planes.

If we project  $C_9^2$  on to a plane from one of its double points we obtain a curve  $C_7^2$  with a quadruple point (the intersection of the plane with  $G$ ) and two triple points, and therefore with one other double point. There is thus one line through either double point of  $C_9^2$  which meets it in two further points, and this must be a generator of the surface. We thus have the two generators which meet  $G$  and are represented on  $\Omega$  by the points  $Q$  and  $R$ .

185. Consider again the normal surface  $F$  in [7], and take two generators  $g$  and  $g'$ ; let  $S$  contain a line  $l$  in the solid  $gg'$ . Then the projected surface  $f$  has a double generator  $G$ , the intersection of  $\Sigma$  with the [5]  $Sgg'$ .

The directrix cubics  $\Delta$  of  $F$  meet  $g$  and  $g'$  in related ranges; hence the solids  $K$  meet the solid  $gg'$  in the lines of a regulus. There are two lines of this regulus meeting  $l$ , so that  $S$  meets two further solids  $K$ . Hence the surface  $f$  has two trisecant planes which do not pass through  $G$ .

$S$  meets  $M_5^6$  in  $l$  and a curve  $\vartheta_5^2$  which has  $l$  for a chord. The chords of  $F$  which meet  $S$  meet  $F$  in  $g, g'$ , and a curve  $C_{18}$  meeting every generator of  $F$  in four points. Since there are eight tangents of  $C_{18}$  which meet  $\vartheta_5^2$  it is of genus 7 and therefore has eight double points. The chords of  $F$  which pass through the intersections of  $l$  and  $\vartheta_5^2$  are chords of  $C_{18}$ ; we thus account for two intersections of  $C_{18}$  with each of  $g$  and  $g'$ . One of the other intersections of  $C_{18}$  and  $g$  is associated with one of its other intersections with  $g'$  and with one of its double points; similarly, for the remaining pair of intersections. The other six double points of  $C_{18}$  lie three in each of two planes which meet  $S$  in lines.

Hence on projection we have a surface  $f$  whose double curve is  $G + C_9^2$ .  $C_9^2$  has two double points and two ordinary points on  $G$ ; it has also two triple points.

To generate this surface we take two twisted cubics in (1, 1) correspondence, the correspondence being specialised to give the double generator.

186. On a rational normal sextic there are  $\infty^1$  involutions which contain, as a pair of corresponding points, the ends of a given chord; each of these gives rise to a quintic ruled surface of which the chord is a generator. On projecting from a point of the chord on to a [5] each of these surfaces

is projected into a quartic ruled surface, one of whose generators passes through the double point  $P$  of the projected curve.

There are  $\infty^8$  quadric fourfolds containing  $C$ , and  $\infty^5$  of these contain such a quartic surface\*. Take one of these latter to be  $\Omega$ . The planes of the conics on the quartic surface form a  $V_3^3$ ;  $\Omega$  meets this in the quartic surface and in two of its planes, these belonging to opposite systems on  $\Omega$ . The quartic surface meets a general plane of  $\Omega$  in two points.

The chords of  $C$  which lie on  $\Omega$  form this quartic ruled surface  $R_2^4$  and a ruled surface  $R_2^{14}$  which has  $C$  as a triple curve. There are two chords  $PQ, PR$  of  $C$  which pass through  $P$  and lie on  $\Omega$ ; these are double generators of  $R_2^{14}$ , and there is one other generator of  $R_2^{14}$  passing through  $P$ . There are twelve points of  $C$  at which two of the three generators of  $R_2^{14}$  coincide†; there are two tangents of  $C$  belonging to  $R_2^4$  and six belonging to  $R_2^{14}$ . Considering then the correspondence between  $C$  and a prime section  $C'$  of  $R_2^{14}$  we have

$$\alpha = 2, \quad \alpha' = 3, \quad p = 0, \quad \eta = 12, \quad \eta' = 6,$$

whence  $p' = 1$ .  $R_2^{14}$  meets a general plane of  $\Omega$  in seven points; thus we have in [3] a ruled surface whose double curve is given by  $C_2 + G + C_7^1$  and bitangent developable by  $E_2 + G + E_7^1$ .

$R_2^4$  and  $R_2^{14}$  have two common generators‡.

If we take any of the four trisecant planes (two of either system) of  $C$  which lie on  $\Omega$  there is a plane of  $\Omega$  of the same system meeting  $R_2^4$  in a conic. Then, considering the intersection of  $R_2^4$  with the [4] containing the two planes, we must conclude that the trisecant plane of  $C$  contains a generator of  $R_2^4$ .  $G$  passes through two double points of  $C_7^1$  and meets it in a further point, while also meeting  $C_2$ . Also  $C_2$  passes through two double points of  $C_7^1$ , while meeting it in two further points: there are exactly similar statements for the bitangent developable.

**187.** Take an involution of pairs of generators on the normal surface

\* It is sufficient to make one of the quadrics through  $C$  contain three generators of the quartic surface in order that it should contain the whole of it.

† These coincidences do not include  $P$ . If we regard  $P$  as two points  $P_1$  and  $P_2$  on different branches of  $C$  there are three generators  $P_1P_2, P_1Q, P_1R$  of  $R_2^{14}$  through  $P_1$  and three others  $P_2P_1, P_2Q, P_2R$  through  $P_2$ . Thus corresponding to the point of  $C$  on either branch at  $P$  we have three distinct points.

This may be contrasted with § 24 and with § 184 above and § 265 below.

‡ Not three, as is apparently given by Brill's formula; this includes a false coincidence at  $P$ . The tangents of  $C$  at  $P$  determine a plane containing a generator of  $R_2^4$  and a generator of  $R_2^{14}$ , which do not of course coincide. This discrepancy is further illustrated by considering the projections of two quintic surfaces, determined by involutions on the normal curve; projecting from a point of their common generator we have two  $g_2^1$ 's on  $C$  giving two quartic surfaces without a common generator.

$F$ , thus obtaining, by means of a directrix quartic  $E$ , a cubic ruled surface with a directrix line  $\lambda$ . Let  $g, g'$  be a pair of generators belonging to the involution, and take a line  $l$  lying in the solid  $gg'$  and passing through the point where  $\lambda$  meets it. Then project from a solid  $S$  containing  $l$  and  $\lambda$ .  $S$  meets  $M_5^6$  in  $l, \lambda$  and an elliptic quartic  $\mathfrak{D}_4^1$  which meets  $l$  and has  $\lambda$  as a chord.

The chords of  $F$  which meet  $S$  meet  $F$  in  $g, g', E$  and a curve  $C_{14}$  meeting each generator in three points and  $E$  in eight points. Since there are six tangents of  $C_{14}$  meeting  $\mathfrak{D}_4^1$  it is of genus 4, so that it must have four double points. The chords of  $F$  passing through the intersections of  $\lambda$  and  $\mathfrak{D}_4^1$  account for four intersections of  $C_{14}^4$  with  $E$ ; the remaining intersections fall into two pairs, with each of which is associated a double point of  $C_{14}^4$ . The other double points of  $C_{14}^4$  are associated with pairs of intersections of  $C_{14}^4$  with  $g$  and  $g'$ ; the remaining pair of these intersections is joined by the chord of  $F$  which passes through the common point of  $l$  and  $\mathfrak{D}_4^1$ .

Hence on projection the surface  $f$  has a double curve  $C_2 + G + C_7^1$ ;  $C_7^1$  has two double points and two simple points on  $C_2$ , two double points and one simple point on  $G$ ;  $G$  meets  $C_2$ .

To generate this surface we take a conic and a cubic in (1, 2) correspondence with a united point; the correspondence being so determined that one point of the conic gives rise to the pair of points of the cubic collinear with it. The joins of the pairs of points on the cubic form a regulus, two lines of which pass through the corresponding points of the conic; thus the planes joining the points of the conic to the pairs of points of the cubic which correspond to them are the tangent planes of a quadric cone. The surface has a bitangent developable  $E_2 + G + E_7^1$ .

188. Let us take two involutions on the rational normal sextic and project on to a [5] from a point of their common chord. Then we obtain a curve  $C$  with a double point  $P$  and two quartic ruled surfaces containing  $C$ ; there are  $\infty^2$  quadrics containing both these surfaces  $R_2^4$  and  $S_2^4$ ; let us take one of these to be  $\Omega$ . The plane of the tangents to  $C$  at  $P$  meets  $\Omega$  in two lines which are generators of  $R_2^4$  and  $S_2^4$  respectively.

The chords of  $C$  which lie on  $\Omega$  form the two ruled surfaces  $R_2^4, S_2^4$  and a third ruled surface  $R_2^{10}$  on which  $C$  is a double curve. There are four tangents of  $C$  belonging to  $R_2^{10}$  and there are four points of  $C$  at which the two generators of  $R_2^{10}$  coincide; this is sufficient to shew that the prime sections of  $R_2^{10}$  are rational curves.  $R_2^{10}$  meets the planes of  $\Omega$  in five points. Thus the points of  $C$  represent the generators of a ruled surface in [3] whose double curve is  $C_2 + D_2 + G + C_5^0$  and bitangent developable  $E_2 + F_2 + G + E_5^0$ .

$R_2^{10}$  has two generators in common with each of  $R_2^4$  and  $S_2^4$ , but these two surfaces themselves have not a common generator. Each trisecant plane of  $C$  which lies on  $\Omega$  contains both a generator of  $R_2^4$  and a generator of  $S_2^4$ . Hence  $C_2$  and  $D_2$  have two intersections through both of which  $C_5^0$  passes,  $C_5^0$  meeting each of them in two further points.  $G$  meets both  $C_2$  and  $D_2$ , and passes through two double points of  $C_5^0$ . There are exactly similar statements for the bitangent developable.

189. Take two involutions of pairs of generators on the normal surface  $F$ . One of these determines, by means of a directrix quartic  $E$ , a cubic ruled surface with a directrix line  $\lambda$ ; the other similarly determines, by means of a directrix quartic  $E'$ , a cubic ruled surface with a directrix line  $\lambda'$ . Let us project from the solid  $S$  containing  $\lambda$  and  $\lambda'$ .

Let  $g, g'$  be the pair of generators common to the two involutions.  $S$  contains the line  $l$  joining the points where the solid  $gg'$  is met by  $\lambda$  and  $\lambda'$ . It meets  $M_5^6$  in  $l, \lambda, \lambda'$  and a cubic  $\mathfrak{D}_3^0$  having  $\lambda$  and  $\lambda'$  as chords.

The chords of  $F$  which meet  $S$  meet  $F$  in  $g, g', E, E'$ , and a curve  $C_{10}$  which meets every generator in two points. Since there are four tangents of  $C_{10}$  meeting  $\mathfrak{D}_3^0$  it is an elliptic curve and has two double points; these are associated with its intersections with  $g$  and  $g'$ . On projection we obtain a ruled surface with a double curve exactly as in the last article.

To generate such a surface we take two conics and place them in (2, 2) correspondence with two united points, the correspondence being specialised to give the double generator. The joins of the pairs of points of either conic which correspond to the points of the other touch another conic, and the planes joining the points of one of the conics to the pairs of points of the other which correspond to them are the tangent planes of a quadric cone. The bitangent developable of the surface thus includes two developables of the second class; it is  $E_2 + F_2 + G + E_5^0$ .

190. The three types of surfaces so far obtained in this section are all self-dual, and, in fact, there are particular cases of surfaces of these types which belong to a linear complex and are represented on  $\Omega$  by curves  $C$  of the type II (B).

A rational sextic in [4] with a double point  $P$  has two trisecants\*; its chords form a locus  $M_3^9$  on which  $C$  is quadruple, the trisecants triple and  $P$  sextuple. Any quadric threefold containing  $C$  contains also its trisecants, and any line on this quadric is met by nine chords of  $C$ . The chords of  $C$  which lie on the quadric form a ruled surface  $R_2^{18}$  whose

\* The trisecant planes of a rational normal sextic form an  $M_5^4$  on which the chords of the curve are a double locus  $M_3^{10}$ . Hence a line meeting a chord of the curve meets two trisecant planes not containing the chord.



prime sections are of genus 2, precisely as in § 184. If we regard the quadric as a prime section of  $\Omega$  we have in [3] a ruled surface whose generators belong to a linear complex, the double curve is given by  $G + C_9^2$  and the bitangent developable by  $G + E_9^2$ .  $C_9^2$  has two triple points; at either of these there intersect three generators of the ruled surface, these lying in a plane which is a triple plane of  $E_9^2$ .  $G$  meets  $C_9^2$  and lies in planes of  $E_9^2$  just as before.

There are  $\infty^2$  quadrics containing  $C$ ; the base of this net of quadrics consists of  $C$  and the two trisecants.

If we take in [5] a rational sextic with a double point there are  $\infty^1$  involutions on the curve containing the pair of points at the double point; each of these gives a quartic ruled surface. Projecting from a point in the plane of a conic of one of these surfaces we obtain in [4] a rational sextic  $C$ , with a double point  $P$ , lying on a quartic ruled surface which has a double line; this double line is itself a chord of  $C$ . There are  $\infty^1$  quadrics containing this ruled surface, and if one of these is regarded as a prime section of  $\Omega$  we obtain in [3] a ruled surface whose double curve is  $C_2 + G + C_7^1$  and bitangent developable  $E_2 + G + E_7^1$ ; but now its generators belong to a linear complex.

Again, we can project from a point of [5] which is common to two planes meeting two different quartic surfaces in conics. We have then two quartic ruled surfaces containing the curve  $C$  in [4]; each of these has a double line which is a chord of  $C$ . There is a quadric containing  $C$  and both these ruled surfaces, and if this is regarded as a prime section of  $\Omega$  we have in [3] a ruled surface whose generators belong to a linear complex, the double curve being  $C_2 + D_2 + G + C_5^0$  and the bitangent developable  $E_2 + F_2 + G + E_5^0$ .

**191.** Take an involution of pairs of points on a rational normal sextic in [6] and project on to a [5] from a point of a chord which does not join a pair of the involution. The joins of the pairs of the involution form a rational normal quintic ruled surface through any two points of which there passes a cubic curve; hence on projection we have a curve  $C$  with a double point  $P$  and a quintic ruled surface with a double point  $P$ ; there is on this surface a plane cubic curve with a double point at  $P$ .

There are quadrics containing this ruled surface. Such a quadric  $\Omega$  necessarily contains the plane of the cubic curve; this plane and the quintic surface form the complete intersection of  $\Omega$  and the cubic cone which projects the surface from  $P$ . The surface meets all planes of  $\Omega$  of the same system as this one in two points, all planes of  $\Omega$  of the opposite system in three points.

There are two chords  $PQ$ ,  $PR$  of  $C$  which pass through  $P$  and lie on  $\Omega$ ; these are generators of the quintic surface  $R_2^5$ . The chords of  $C$  which lie

on  $\Omega$  generate this surface  $R_2^5$  and a surface  $R_2^{13}$  on which  $C$  is a triple curve; this latter surface meets all planes of  $\Omega$  of one system in seven points and all planes of the other system in six points.  $PQ$  and  $PR$  are generators of  $R_2^{13}$ , while  $R_2^{13}$  and  $R_2^5$  have three other common generators. There are also two generators of  $R_2^{13}$  passing through  $P$  and lying in the plane of the two tangents to  $C$  at  $P$ .

There are eight points of  $C$  other than  $P$  at which two of the three generators of  $R_2^{13}$  coincide, and there are six tangents of  $C$  which are generators of  $R_2^{13}$ . We thus find that the prime sections of  $R_2^{13}$  are rational curves.

A trisecant plane of  $C$  which lies on  $\Omega$  contains three generators of  $R_2^{13}$  if it is of the opposite system to the plane of the cubic on  $R_2^5$ , but if of the same system it must contain a generator of  $R_2^5$ .

Suppose that the plane cubic on  $R_2^5$  lies in a  $\rho$ -plane of  $\Omega$ . Then in [3] we have a ruled surface whose double curve is  $C_2 + G + C_7^0$ .  $G$  lies in the plane of  $C_2$  and  $C_7^0$  passes through the intersections of  $G$  and  $C_2$ ;  $C_7^0$  meets  $G$  in two other points and  $C_2$  in three other points, having two triple points. The bitangent developable is  $E_3^0 + G + E_6^0$ .  $E_6^0$  has two double planes which are planes of  $E_3^0$ ; it has two planes through  $G$  in common with  $E_3^0$  and three further planes in common with  $E_3^0$ ; it has also two other planes through  $G$ .

Similarly, if the plane cubic on  $R_2^5$  lies in a  $\varpi$ -plane of  $\Omega$  we obtain the dual surface in [3]; the double curve is  $C_3^0 + G + C_6^0$  and the bitangent developable  $E_2 + G + E_7^0$ .

192. Take on the normal surface  $F$  in [7] an involution of pairs of generators; then a directrix quartic  $E$  gives rise to a cubic ruled surface with a directrix line  $\lambda$ . Choose  $S$  to contain  $\lambda$  and to meet the solid containing a pair of generators  $g, g'$  in a line  $l$ ;  $g, g'$  not being a pair of the involution. Then the [6]  $ES$  contains  $g$  and  $g'$ , so that the surface  $f$  has a double generator lying in the plane of a double conic.

$S$  meets  $M_5^6$  in  $\lambda, l$  and a rational quartic  $\mathfrak{d}_4^0$  having  $\lambda$  for a trisecant and  $l$  for a chord.

The chords of  $F$  meeting  $S$  meet  $F$  in  $g, g', E$  and a curve  $C_{14}$  meeting every generator in three points. There are six tangents of  $C_{14}$  meeting  $\mathfrak{d}_4^0$ , so that it is a  $C_{14}^2$  with six double points; these double points lie three in each of two planes meeting  $S$  in lines.  $C_{14}^2$  has eight intersections with  $E$ ; six of these are accounted for by the chords of  $F$  which pass through the points where  $\lambda$  meets  $\mathfrak{d}_4^0$ , the other two are the points of  $E$  which correspond in the involution to its intersections with  $g$  and  $g'$ .

Thus on projection we obtain a surface  $f$  whose double curve is  $C_2 + G + C_7^0$ ;  $G$  lies in the plane of  $C_2$  and  $C_7^0$  passes through the two

points where it meets  $C_2$ ;  $C_7^0$  meets  $G$  in two other points and  $C_2$  in three other points, and has two triple points.

To generate this surface we take a conic and a twisted cubic in (1, 2) correspondence with a united point. The cubic meets the plane of the conic in two other points; the points of the conic which correspond to these are to be those two points in which it meets the line joining them. The planes which join the points of the conic to the pairs of corresponding points on the cubic form a developable of the third class  $E_3^0$ ; the bitangent developable of the surface is  $E_3^0 + G + E_6^0$ .

**193.** If we take a prime section  $C_6$  of  $F$ , an involution of pairs of generators gives a quintic ruled surface with a directrix conic  $\Gamma$ . A solid  $S$  through the plane of  $\Gamma$  meets  $M_5^6$  in  $\Gamma$  and an elliptic quartic  $\vartheta_4^1$  having three intersections with  $\Gamma$ . This quartic will meet the plane of  $\Gamma$  in a fourth point, and through this point there passes a transversal to two generators  $g, g'$  of  $F$  (not forming a pair of the involution). Let us take a line  $l$  through this point and lying in the solid  $gg'$ , and then choose  $S$  to contain  $\Gamma$  and  $l$ .  $S$  meets  $M_5^6$  further in a twisted cubic  $\vartheta_3^0$  meeting  $\Gamma$  in three points and  $l$  in two points.

The chords of  $F$  meeting  $S$  meet  $F$  in  $C_6, g, g'$  and a curve  $C_{12}$  meeting every generator in three points; there are six tangents of  $C_{12}$  meeting  $\vartheta_3^0$ , so that it is a  $C_{12}^2$ , and has two double points. It meets  $C_6$  in twelve points: six of these arise from the three chords of  $F$  through the intersections of  $\vartheta_3^0$  and  $\Gamma$ ; four more are associated as two pairs with the double points of  $C_{12}^2$ , while the remaining two are the points of  $C_6$  which correspond in the involution to its intersections with  $g$  and  $g'$ .

Thus on projection the surface  $f$  has a double curve  $C_3^0 + G + C_6^0$ ;  $G$  is a chord of  $C_3^0$  and  $C_6^0$  passes through their two intersections, meeting  $G$  in two further points and  $C_3^0$  in three further points. Also  $C_6^0$  has two double points on  $C_3^0$ .

To generate this surface we take two twisted cubics in (1, 2) correspondence with three united points. The cubics have a common chord; to the points in which this meets the second cubic correspond those in which it meets the first. The planes joining the points of the first cubic to the pairs of points which correspond to them on the second cubic are the tangent planes of a quadric cone. The bitangent developable of the surface is  $E_2 + G + E_7^0$ .

**194.** We now pass on to consider curves on  $\Omega$  of the type II (C);  $C$  is contained in a [4] and has two double points  $P$  and  $Q$ . These represent two double generators  $G$  and  $H$  of the surface in [3]. The chords of  $C$  form a locus  $M_3^8$ , there being eight of them meeting an arbitrary line in [4].

If we regard a quadric threefold containing  $C$  as a prime section of  $\Omega$ , every plane of  $\Omega$  meets the [4] containing  $C$  in a line which is met by eight chords of  $C$ . Hence the ruled surface in [3] has a double curve  $G + H + C_8$  and a bitangent developable  $G + H + E_8$ .

The chords of  $C$  lying on  $\Omega$  form a ruled surface  $R_2^{16}$  (the intersection of  $\Omega$  and  $M_3^8$ ) on which  $C$  is a quadruple curve. There are apparently twenty-four points of  $C$  at which two of the four generators of  $R_2^{16}$  coincide, but these include  $P$  and  $Q$ , each counted four times. There are eight tangents of  $C$  which lie on  $\Omega$ , so that we have a correspondence between  $C$  and a prime section  $C'$  of  $R_2^{16}$  for which

$$\alpha = 2, \quad \alpha' = 4, \quad p = 0, \quad \eta = 16, \quad \eta' = 8,$$

whence  $p' = 1$ , and we have a double curve  $G + H + C_8^1$  and a bitangent developable  $G + H + E_8^1$ .

There are two chords  $PX, PY$  of  $C$  which pass through  $P$  and lie on  $\Omega$ ; these are double generators of  $R_2^{16}$ . There are two other generators of  $R_2^{16}$  passing through  $P$ ; they lie in the plane containing the two tangents of  $C$  at  $P$ .  $C_8^1$  has two double points on  $G$  and meets it in two further points;  $E_8^1$  has two double planes and also two simple planes through  $G$ . Similarly for  $H$ .

195. Consider again the normal surface  $F$  in [7]; let us take a line  $l$  lying in the solid containing a pair of generators  $g, g'$  and a line  $m$  lying in the solid containing another pair of generators  $h, h'$ . Choose  $S$  to be the solid  $lm$ : the projected surface in  $\Sigma$  has two double generators  $G$  and  $H$ ; the intersections of  $\Sigma$  with the [5]'s  $Sgg'$  and  $Shh'$  respectively.

The chords of  $F$  meeting  $S$  meet  $F$  in the four generators  $g, g', h, h'$  and a curve  $C_{16}$  meeting each generator in four points.  $S$  meets  $M_3^6$  in  $l, m$  and an elliptic quartic  $\mathfrak{D}_4^1$  having  $l$  and  $m$  as chords. Since there are eight tangents of  $C_{16}$  meeting  $\mathfrak{D}_4^1$  it is a  $C_{16}^5$ , which has four double points. The chords of  $F$  passing through the intersections of  $l$  and  $\mathfrak{D}_4^1$  meet  $F$  in points common to  $C_{16}^5$  and  $g$  or  $g'$ ; the other four intersections of  $C_{16}^5$  with  $g$  and  $g'$  are associated in pairs with two of the double points of  $C_{16}^5$ .

Hence on projection we obtain a surface  $f$  whose double curve is  $G + H + C_8^1$ ;  $C_8^1$  has two double points on  $G$  and two double points on  $H$ , meeting each of  $G$  and  $H$  in two further points. The surface is generated by a (1, 1) correspondence between two twisted cubics, the correspondence being specialised to give the double generators.

196. If the quadric containing  $C$  is also made to contain the line  $PQ$  we have a particular ruled surface for which the double generators  $G$  and  $H$  intersect. Their point of intersection is a quadruple point of  $C_8^1$  which meets each of them in two further points; the plane containing them is a quadruple plane of  $E_8^1$ , which has two further planes passing through each of them.

In order to obtain this surface by projection we take the normal surface in [7] with a directrix conic  $\Gamma$ ; take a line  $l$  in the solid determined by a pair of generators  $g, g'$  and a line  $m$  in the solid determined by another pair of generators  $h, h'$ . Then project from the solid  $S$  containing  $l$  and  $m$ . We have on projection a surface with two coplanar double generators; their plane meets the surface further in a conic and is the intersection of  $\Sigma$  with the [6] containing  $S, g, g', h, h'$  and  $\Gamma$ .

To generate this surface we take a conic and a rational skew quartic in (1, 1) correspondence; the points in which the plane of the conic meets the quartic fall into two pairs, and the pair of points of the conic corresponding to either of these pairs is given by the intersections of the conic with the line joining the pair.

197. Let us project a rational normal sextic in [6] from a point on one of its chords. We obtain a rational sextic in [5] with a double point  $P$ . A quintic ruled surface which contains the chord of the normal curve gives on projection a quartic ruled surface with  $\infty^1$  directrix conics; each of these conics meets the projected curve twice, one of them passing through  $P$ .

Now let us project this curve on to a [4]; the point of projection is to be chosen in the plane of one of the conics of the quartic surface and also on the chord of the curve which lies in that plane. We obtain in [4] a rational sextic  $C$  with two double points  $P$  and  $Q$ ; there is a quartic ruled surface containing the curve, with a double line passing through  $Q$ .

There are  $\infty^3$  quadrics containing  $C$ , so that there are  $\infty^1$  containing the quartic ruled surface; let us regard one of these latter quadrics as a prime section of  $\Omega$ .

The chords of  $C$  which lie on  $\Omega$  form the quartic surface  $R_2^4$  and a ruled surface  $R_2^{12}$  on which  $C$  is a triple curve. Through  $P$  there pass two chords of  $C$ ,  $PX$  and  $PY$ , which lie on  $\Omega$ ; these are double generators of  $R_2^{12}$ . Similarly we have  $QZ$  and  $QT$ , but these are common generators of  $R_2^4$  and  $R_2^{12}$ . The plane of the tangents to  $C$  at  $P$  meets  $\Omega$  in a generator of  $R_2^4$  and a generator of  $R_2^{12}$ ; the plane of the tangents to  $C$  at  $Q$  meets  $\Omega$  in two generators of  $R_2^{12}$ .

The twelve points of  $C$  at which two generators of  $R_2^{12}$  coincide include  $Q$  counted four times, so that there are eight other points. There are six tangents of  $C$  belonging to  $R_2^{12}$ , so that we can deduce that the prime sections of  $R_2^{12}$  are rational curves. We have thus in [3] a ruled surface whose double curve is  $G + H + C_2 + C_6^0$  and bitangent developable  $G + H + E_2 + E_6^0$ .  $H$  lies in the plane of  $C_2$ ;  $C_6^0$  passes through the intersections of  $H$  with  $C_2$  and meets  $H$  in two further points;  $C_2$  and  $C_6^0$  have two further intersections.  $G$  meets  $C_2$ ; it passes through two double points of  $C_6^0$  and meets  $C_6^0$  in one further point.

198. Take an involution of pairs of generators of  $F$ ; we then have, through any directrix quartic  $E$ , a cubic ruled surface with a directrix line  $\lambda$ . Take a line  $l$  meeting  $\lambda$  and lying in the solid containing the pair of generators  $g, g'$ ; this being a pair of the involution. There is a point of the plane  $l\lambda$  which is not on either of the lines  $l$  or  $\lambda$  and through which passes a chord of  $F^*$ ; we can then take a line  $m$  through this point which lies in the solid containing a pair of generators  $h, h'$  (not belonging to the involution). Let us project from the solid  $S$  containing  $\lambda, l$  and  $m$ .

$S$  meets  $M_5^6$  in  $l, \lambda, m$  and a cubic  $\vartheta_3^0$  meeting  $l$  and having  $\lambda$  and  $m$  as chords. The chords of  $F$  which meet  $S$  meet  $F$  in  $g, g', h, h', E$  and a curve  $C_{12}$  which meets every generator in three points; since there are six tangents of  $C_{12}$  meeting  $\vartheta_3^0$  it is a  $C_{12}^2$  and therefore has two double points.  $C_{12}^2$  has six intersections with  $E$ ; four of these are accounted for by the chords of  $F$  passing through the intersections of  $\lambda$  and  $\vartheta_3^0$ . The other two are associated with the points where  $E$  is met by the generators which are paired with  $h$  and  $h'$  in the involution and with points in which  $C_{12}^2$  meets  $h$  and  $h'$ . The remaining intersections of  $C_{12}^2$  with  $h$  and  $h'$  are accounted for by the chords of  $F$  passing through the intersections of  $m$  and  $\vartheta_3^0$ . There is also a chord of  $F$  passing through the point common to  $l$  and  $\vartheta_3^0$ , this accounts for an intersection of  $C_{12}^2$  with each of  $g$  and  $g'$ ; the other intersections of  $C_{12}^2$  with these generators are associated with its double points. We thus obtain in  $\Sigma$  a surface with a double curve precisely as in the last article.

To generate this surface we take a conic and a twisted cubic in (1, 2) correspondence with a united point. The points of the conic which correspond to the other two points in which the cubic meets its plane are the intersections of the conic with the line joining these points. Also there is a point of the conic for which the pair of corresponding points on the cubic consists of the ends of the chord through this point. The planes of the pairs of generators which intersect in the points of the conic are the tangent planes of a quadric cone. Hence the bitangent developable is  $G + H + E_2 + E_6^0$ .

199. Given any chord of a rational normal sextic there are  $\infty^1$  quintic ruled surfaces which contain it. Through each point of the chord there pass  $\infty^1$  cubic curves on each of these ruled surfaces, and each cubic curve meets the normal sextic in two points. From any point on any chord of the curve we can obtain any other chord of the curve; given any two chords of the curve, through any point of either we have a cubic curve meeting the other in its two intersections with the curve. It will be possible to choose the pair of chords and the points on them so that the

solids containing the two cubic curves have a plane in common. Then project on to a [4] from the line joining the points in which the cubic curves meet the chords. We obtain a rational sextic  $C$  with two double points  $P$  and  $Q$ . There are two quartic ruled surfaces composed of chords of  $C$ , each of these surfaces having a double line; one double line passes through  $P$  and the other through  $Q$ , and the two double lines intersect.

There are  $\infty^1$  quadric threefolds containing each ruled surface; these two systems are both contained in the system of  $\infty^2$  quadrics which contain  $C$  and the intersection of the double lines of the two surfaces. Hence there is a quadric containing both the ruled surfaces; let us regard this quadric as a prime section of  $\Omega$ .

The chords of  $C$  which lie on  $\Omega$  form three ruled surfaces; a ruled surface  $R_2^8$  on which  $C$  is a double curve, a quartic ruled surface  $R_2^4$  with a double line through  $P$  and a quartic ruled surface  $S_2^4$  with a double line through  $Q$ . The two chords  $PX$ ,  $PY$  of  $C$ , which pass through  $P$  and lie on  $\Omega$ , are common generators of  $R_2^8$  and  $R_2^4$ , while the plane of the two tangents of  $C$  at  $P$  meets  $\Omega$  in two lines, one of which is a generator of  $R_2^8$  and the other of  $S_2^4$ . Similarly, the two chords  $QZ$ ,  $QT$  of  $C$  which pass through  $Q$  and lie on  $\Omega$  are common generators of  $R_2^8$  and  $S_2^4$ , while the plane of the two tangents of  $C$  at  $Q$  meets  $\Omega$  in two lines, one of which is a generator of  $R_2^8$  and the other of  $R_2^4$ . There is another generator common to  $R_2^8$  and  $R_2^4$  and another common to  $R_2^8$  and  $S_2^4$ ;  $R_2^4$  and  $S_2^4$  also have a generator in common. There are four points of  $C$  at which the two generators of  $R_2^8$  coincide, while there are four tangents of  $C$  which are generators of  $R_2^8$ ; whence the prime sections of  $R_2^8$  are rational curves.

We thus have in [3] a ruled surface whose double curve is  $G + H + C_2 + D_2 + C_4^0$  and bitangent developable  $G + H + E_2 + F_2 + E_4^0$ .

The plane of  $C_2$  contains  $G$ ;  $C_4^0$  passes through the intersections of  $C_2$  and  $G$ , meeting each of  $G$  and  $C_2$  once further. The plane of  $D_2$  contains  $H$ ;  $C_4^0$  passes through the intersections of  $D_2$  and  $H$ , meeting each of  $H$  and  $D_2$  once further.  $C_2$  and  $D_2$  have one intersection. There are similar statements for the bitangent developable.

**200.** Take two pairs of generators  $g, g'$  and  $h, h'$  of the normal surface  $F$  in [7]. An involution containing the pair  $gg'$  gives an axis  $\lambda$  of a directrix quartic  $E$ , while another involution containing the pair  $hh'$  gives an axis  $\mu$  of another directrix quartic  $E'$ . We take the common pair of the two involutions to be the generators through the two intersections of  $E$  and  $E'$  and take  $\lambda$  and  $\mu$  to intersect in a point of the common chord of  $E$  and  $E'$ .

The [5]  $\lambda\mu gg'$  meets the [5]  $\lambda\mu hh'$  in a [3]  $S$ ;  $S$  contains a line  $l$  in the [3]  $gg'$  and a line  $m$  in the [3]  $hh'$ . We project from  $S$  on to  $\Sigma$ .  $S$  meets

$M_5^6$  in the four lines  $\lambda, \mu, l, m$  and a conic  $\vartheta_2$  which meets each of these lines once.  $l$  meets  $\lambda$  and  $m$  meets  $\mu$ .

The chords of  $F$  which meet  $S$  meet  $F$  in  $g, g', h, h', E, E'$  and a curve  $C_3$  meeting every generator in two points. There are four tangents of  $C_3$  meeting  $\vartheta_2$  so that it is an elliptic curve without double points. It meets  $E$  in four points; two of these are accounted for by the chord of  $F$  passing through the intersection of  $\vartheta_2$  and  $\lambda$ ; the others are associated with the points in which  $E$  is met by the generators corresponding to  $h$  and  $h'$  in the first involution and with intersections of  $C_3$  with  $h$  and  $h'$ . Similarly for the intersections of  $C_3$  with  $E'$ . On projection we have in  $\Sigma$  a surface whose double curve is  $G + H + C_2 + D_2 + C_4^0$  precisely as in the last article.

To generate this surface we place two conics  $C_2$  and  $D_2$  in (2, 2) correspondence with a doubly united point. To the other point in which  $C_2$  meets the plane of  $D_2$  correspond a pair of points of  $D_2$  collinear with it, and to the second point in which  $D_2$  meets the plane of  $C_2$  correspond a pair of points of  $C_2$  collinear with it. The planes joining the points of either conic to the pairs of points of the other conic which correspond to them are the tangent planes of a quadric cone. We thus have a bitangent developable  $G + H + E_2 + F_2 + E_4^0$ .

**201.** We now consider the curve  $C$  on  $\Omega$  of the type II (D); we have a rational sextic in [4] with a triple point  $P$ —the projection of a normal sextic from a line in one of its trisecant planes. The chords of  $C$  form a locus  $M_3^7$  on which  $C$  is a quadruple curve and  $P$  a sextuple point.

The chords of  $C$  which lie on  $\Omega$  form a ruled surface  $R_2^{14}$ , the intersection of  $\Omega$  with  $M_3^7$ ; there are no lines through  $P$  lying on  $\Omega$  and meeting  $C$  again but there are six generators of  $R_2^{14}$  passing through  $P$ , viz. the lines in which  $\Omega$  is met by the three planes containing the pairs of tangents of  $C$  at  $P$ .

There are apparently twenty-four points of  $C$  at which two of the four generators of  $R_2^{14}$  coincide; but these include  $P$  counted twelve times\*. There are eight tangents of  $C$  which are generators of  $R_2^{14}$ , and we deduce that the prime sections of  $R_2^{14}$  are rational curves. We have in [3] a ruled surface whose double curve is  $3G + C_7^0$  and bitangent developable  $3G + E_7^0$ ; the triple generator  $G$  being represented on  $\Omega$  by the point  $P$ .  $G$  meets  $C_7^0$  in six points, while six planes of  $E_7^0$  pass through  $G$ .

**202.** Take now the normal surface  $F$  in [7], and let  $S$  be chosen to lie in the [5] containing three generators  $g_1, g_2, g_3$ . The solids containing the pairs of generators  $g_2g_3, g_3g_1, g_1g_2$  meet  $S$  in lines  $l_1, l_2, l_3$  respectively. The projected surface has a triple generator  $G$ , the intersection of  $\Sigma$  with the [5] containing  $S, g_1, g_2, g_3$ .

\* There are three different branches of  $C$  passing through  $P$ .



The curves  $\Delta$  meet  $g_1, g_2, g_3$  in related ranges, so that the solids  $K$  containing them meet the [5] containing the three generators in the planes of a  $V_3^3$ . Thus  $S$  meets the  $M_4^4$  formed by the solids  $K$  in a cubic curve  $\vartheta_3^0$ ; it meets  $M_5^6$  in  $\vartheta_3^0, l_1, l_2, l_3$ , and  $l_1, l_2, l_3$  are chords of  $\vartheta_3^0$ .

The chords of  $F$  which meet  $S$  meet  $F$  in the generators  $g_1, g_2, g_3$  each counted twice and in a curve  $C_{14}$  meeting each generator of  $F$  four times. There are eight tangents of  $C_{14}$  meeting  $\vartheta_3^0$ , so that it is a  $C_{14}^3$  and has no double points. A chord of  $F$  through an intersection of  $\vartheta_3^0$  and  $l_1$  meets  $F$  in two points on  $C_{14}^3$ , one of which is on  $g_2$  and the other on  $g_3$ . Hence on projection we have a surface  $f$  with a double curve  $3G + C_7^0$ ;  $G$  meets  $C_7^0$  in six points.

To generate this surface we take two rational plane cubics in (1, 1) correspondence, there being three pairs of corresponding points on the line of intersection of their planes.

*Surfaces with a directrix line which is not a generator*

203. Consider on  $\Omega$  a curve  $C$  of the type III (A); it lies in the tangent prime at a point  $O$ , meeting every plane  $\omega$  through  $O$  in five points and every plane  $\rho$  through  $O$  in one point. Then we have in [3] a ruled surface with a directrix line  $R$ ; through every point of  $R$  there pass five generators, while any plane through  $R$  contains one generator.

Any plane  $\rho$  of  $\Omega$  meets the tangent prime at  $O$  in a line, and the plane  $\omega$  containing this line joins it to  $O$ ; all the ten chords of  $C$  meeting the line are contained in this plane  $\omega$ . Thus the double curve of the ruled surface in [3] is its directrix  $R$  counted ten times.

A plane  $\omega$  of  $\Omega$  meets the tangent prime at  $O$  in a line which is met by ten chords of  $C$ . The chords of  $C$  lying on  $\Omega$  form a ruled surface, and the genus of the prime sections of the ruled surface is found to be three as in former instances. Hence the surface in [3] has a bitangent developable  $E_{10}^3$ .

The four trisecants of  $C$  are all joined to  $O$  by  $\omega$ -planes; the  $\rho$ -planes through these trisecants represent four planes which are triple planes of  $E_{10}^3$  and tritangent planes of the ruled surface.

Dually, for a curve  $C$  of the type III (B), we have a double curve  $C_{10}^3$  with four triple points and a bitangent developable  $10R$ .

204. To obtain the first of these surfaces by projection, we take the normal surface  $F$  in [7] and project from a solid  $S$  which lies in a [5] containing a rational quintic curve of  $F$  (there are  $\infty^5$  such curves on  $F$ ). Then the [5] meets  $\Sigma$  in a line  $R$  which is a directrix of  $f$ ; through any point of  $R$  there pass five generators, while any plane through  $R$  contains one generator. The only chords of  $F$  which meet  $S$  are those of the quintic

curve, we thus have a  $\mathfrak{D}_6^3$  in  $S$ . These chords meet  $F$  in the quintic curve counted four times, four of them passing through each point of the curve.

To generate  $f$  we take a twisted cubic and one of its chords, placing them in (5, 1) correspondence with two united points.

To obtain the second surface we must project the normal surface in [7] which has a directrix line  $\lambda$ . Projecting on to  $\Sigma$  from a solid  $S$  we obtain a surface  $f$  with a directrix line  $R$ —the intersection of  $\Sigma$  with the [5]  $S\lambda$ ; through each point of  $R$  there passes one generator, while every plane through  $R$  contains five generators. To generate  $f$  we take a line and a rational quintic in (1, 1) correspondence.

The chords of this surface  $F$  also form an  $M_5^6$  meeting  $S$  in a  $\mathfrak{D}_6^3$ . Also the tangent solids of  $F$  meet an arbitrary [5] in the tangents of a rational normal quintic, so that they form an  $M_4^8$ . We have on  $F$  a curve  $C_{20}^9$ , eight of whose tangents meet  $\mathfrak{D}_6^3$ ; it has twelve double points which lie three in each of four planes which meet  $S$  in lines. Thus the double curve of the projected surface is a  $C_{10}^3$  with four triple points.

**205.** Take now a curve  $C$  on  $\Omega$  of the type III (C). The  $\rho$ -planes through  $O$  cut out a  $g_2^1$  on  $C$ ; of the eight tangents of  $C$  which lie on  $\Omega$  two lie in  $\rho$ -planes and the other six in  $\omega$ -planes. We find that the ruled surface in [3] has a double curve  $6R + C_4^0$  and a bitangent developable  $R + E_9^1$ .  $R$  is a trisecant of  $C_4^0$ ; its three intersections with  $C_4^0$  are represented on  $\Omega$  by the  $\omega$ -planes which contain the chords of  $C$  passing through  $O$ . Similarly,  $E_9^1$  has three planes passing through  $R$ . Also  $E_9^1$  has four triple planes; these are represented on  $\Omega$  by the  $\rho$ -planes through the four trisecants of  $C$ , which are all joined to  $O$  by  $\omega$ -planes.

To obtain this surface by projection we choose the solid  $S$  to meet a [4] containing a directrix quartic  $E$  of the normal surface  $F$  in a plane. Then the [5] containing  $S$  and  $E$  meets  $\Sigma$  in a directrix  $R$  of  $f$ ; through any point of  $R$  there pass four generators, while any plane through  $R$  contains two generators.

The chords of  $E$  meet  $S$  in an elliptic cubic  $\mathfrak{D}_3^1$ , there are three of the chords passing through each point of  $E$ . The chords of  $F$  meet  $S$  in  $\mathfrak{D}_3^1$  and a twisted cubic  $\mathfrak{D}_3^0$  meeting  $\mathfrak{D}_3^1$  in three points. These chords meet  $F$  in  $E$  counted three times and a rational curve  $C_8^0$  meeting each generator of  $F$  once only; there are six tangents of  $E$  meeting  $\mathfrak{D}_3^1$  and two tangents of  $C_8^0$  meeting  $\mathfrak{D}_3^0$ .  $C_8^0$  has six intersections with  $E$ ; these are accounted for by the three chords of  $F$  through the intersections of  $\mathfrak{D}_3^1$  and  $\mathfrak{D}_3^0$ . Hence, on projection, the double curve of  $f$  is  $6R + C_4^0$ ,  $R$  being a trisecant of  $C_4^0$ . The surface has four tritangent planes since  $S$  meets four of the solids  $K$ .

To generate  $f$  we take a line and a twisted cubic with one intersection, placing them in (1, 4) correspondence with a united point.

206. When  $C$  is of the type III (D) we have in [3] a ruled surface whose double curve is  $R + C_9^1$  and bitangent developable  $6R + E_4^0$ ;  $R$  is a trisecant of  $C_9^1$ , which has four triple points, and there are three planes of  $E_4^0$  passing through  $R$ .

To obtain this surface by projection we take the normal surface  $F$  in [7] which has a directrix conic  $\Gamma$ , and project from a solid  $S$  which meets the plane of  $\Gamma$  in a point  $O$ . Then the [5]  $S\Gamma$  meets  $\Sigma$  in a line  $R$  which is a directrix of  $f$ ; through any point of  $R$  there pass two generators, while any plane through  $R$  contains four generators.

The chords of  $F$  form an  $M_5^6$  on which the plane of  $\Gamma$  is a triple plane\*, and  $S$  meets  $M_5^6$  in an elliptic sextic  $\vartheta_6^1$  with a triple point at  $O$ . The chords which meet  $S$  meet  $F$  in  $\Gamma$  and a curve  $C_{18}$  meeting every generator of  $F$  in three points; there are two tangents of  $\Gamma$  passing through  $O$  and six tangents of  $C_{18}$  meeting  $\vartheta_6^1$ . Thus  $C_{18}$  is of genus 4, and has twelve double points; these lie three in each of four planes, each plane meeting  $S$  in a line.  $C_{18}$  has six intersections with  $\Gamma$ , these lie on three lines through  $O$ .

Hence on projection we have in  $\Sigma$  a ruled surface  $f$  whose double curve is  $R + C_9^1$ ;  $C_9^1$  has four triple points and meets  $R$  three times.

This surface  $f$  is generated by a line and a rational quartic in (1, 2) correspondence. The joins of the pairs of points of the quartic which correspond to the points of the line form a cubic ruled surface; thus we have a developable of the fourth class formed by the bitangent planes of the surface which do not pass through  $R$ . There are three planes of this developable passing through  $R$  since there are three generators of the cubic ruled surface meeting  $R$ .

207. Any prime section  $C_6^0$  of  $F$  meets  $\Gamma$  in two points; let us take a  $g_2^1$  on  $C_6^0$  containing this pair of points; then we have a quintic ruled surface with a directrix conic  $\gamma$  meeting the plane of  $\Gamma$  in a point  $O$ . Let us choose  $S$  to contain  $\gamma$ . Then  $S$  meets  $M_5^6$  in  $\gamma$  and a rational quartic  $\vartheta_4^0$  having a double point at  $O$ ,  $\vartheta_4^0$  meeting  $\gamma$  in two other points.

\* The equations of  $F$  can be put in the form

$$\frac{x_0}{x_1} = \frac{x_1}{x_2} = \frac{x_2}{x_3} = \frac{x_3}{x_4} = \frac{x_5}{x_6} = \frac{x_6}{x_7},$$

and the chords form the  $M_5^6$

$$\left\| \begin{array}{cccc} x_0 & x_1 & x_2 & x_5 \\ x_1 & x_2 & x_3 & x_6 \\ x_2 & x_3 & x_4 & x_7 \end{array} \right\| = 0.$$

A plane meeting the plane of  $\Gamma$  meets  $M_5^6$  in three further points, since

$$\left\| \begin{array}{cccc} \alpha x_0' + \beta x_0'' & \alpha x_1' + \beta x_1'' & \alpha x_2' + \beta x_2'' & \alpha x_5' + \beta x_5'' + \gamma \xi_5 \\ \alpha x_1' + \beta x_1'' & \alpha x_2' + \beta x_2'' & \alpha x_3' + \beta x_3'' & \alpha x_6' + \beta x_6'' + \gamma \xi_6 \\ \alpha x_2' + \beta x_2'' & \alpha x_3' + \beta x_3'' & \alpha x_4' + \beta x_4'' & \alpha x_7' + \beta x_7'' + \gamma \xi_7 \end{array} \right\| = 0$$

has three solutions in  $\alpha : \beta : \gamma$  other than  $\alpha = \beta = 0$ .

The chords of  $F$  which meet  $S$  meet  $F$  in  $\Gamma$ ,  $C_6^0$  and a curve  $C_{12}$  meeting each generator in two points. There are two tangents of  $\Gamma$  passing through  $O$  and two tangents of  $C_6^0$  meeting  $\gamma$ ; so that there are four tangents of  $C_{12}$  meeting  $\mathfrak{D}_4^0$ . Hence  $C_{12}$  is an elliptic curve and has four double points.  $C_{12}$  meets  $C_6^0$  in twelve points; four of these are accounted for by the chords of  $F$  through the two intersections (other than  $O$ ) of  $\gamma$  and  $\mathfrak{D}_4^0$ ; the rest are associated in pairs with the double points of  $C_{12}$ .

On projection we have in  $\Sigma$  a surface  $f$  whose double curve is  $R + C_3^0 + C_6^0$ .  $C_6^0$  has four double points, through all of which  $C_3^0$  passes; these two curves have two other intersections.  $R$  meets  $C_3^0$  once and  $C_6^0$  twice.

The surface is generated by a line  $R$  and a twisted cubic  $C_3^0$  with one intersection, these being in (2, 2) correspondence with a doubly united point. The pairs of points of the cubic corresponding to the points on the line give chords of the cubic which form a quartic ruled surface, one generator of this surface meeting the line in the point which gives rise to it. Hence the bitangent planes of the surface which do not pass through  $R$  form a rational developable  $E_4^0$ , three of whose planes pass through  $R$ .

We have also the dual surface whose double curve is  $6R + C_4^0$  and bitangent developable  $R + E_3^0 + E_6^0$ . To generate this we take a line  $R$  and a twisted cubic in (1, 4) correspondence with a united point; but now the  $g_4^1$  on the cubic is such that each of its sets is made up of two pairs of an involution. The planes joining  $R$  to the pairs of points of this involution give the developable  $E_3^0$ , one of whose planes passes through  $R$ .

**208.** Consider now a normal sextic ruled surface  $F$  in [7] and two of its prime sections. Suppose that we have an involution of pairs of generators of  $F$ , thus obtaining a quintic ruled surface through one prime section, while another involution of pairs of generators gives similarly a quintic ruled surface through the other prime section. Let us suppose further that the directrix conics of these two quintic surfaces have two intersections; in what circumstances can the solid  $S$  containing the planes of these two conics be chosen as a solid from which we can project  $F$  on to  $\Sigma$ ?

The restriction on  $S$  is that it must not meet  $F$ ; hence we immediately conclude that the two involutions must be different and that the solid containing the pair of generators common to them necessarily passes through one of the two intersections of the conics. Through the other intersection  $O$  there will pass two chords of  $F$ , so that the four generators of  $F$  so determined all meet the same plane and lie in a [6]. Hence  $F$  has a directrix conic  $\Gamma$ , the plane of  $\Gamma$  being that of the two chords through  $O$ , and  $S$  meets the plane of  $\Gamma$  in  $O$ .

Taking then the normal surface with a directrix conic  $\Gamma$  let us choose  $S$  in this way; it meets the plane of  $\Gamma$  in  $O$  and contains two conics  $\gamma$  and  $\delta$  which pass through  $O$  and have another intersection.  $S$  meets  $M_5^6$  further in a third conic  $\mathfrak{D}_2$ , which also passes through  $O$  and meets  $\gamma$  and  $\delta$  each in one other point. The chords of  $F$  meeting  $S$  meet  $F$  in  $\Gamma$  and three prime sections; on projection we have a surface  $f$  whose double curve consists of a directrix  $R$  and three twisted cubics;  $R$  meets all the twisted cubics which have four points in common, any two of them having a further intersection. The bitangent developable is again  $6R + E_4^0$ .

We have also the dual surface whose double curve is  $6R + C_4^0$ ; the bitangent developable consisting of the pencil of planes through  $R$  and three cubic developables. This is also generated by a line  $R$  and a twisted cubic in (1, 4) correspondence with a united point; but now the  $g_4^1$  on the cubic is such that each of its sets is made up of pairs of three different involutions. If  $a_1 a_2 a_3 a_4$  typifies a set of the  $g_4^1$  then  $a_1 a_2$ ,  $a_3 a_4$  are pairs of one involution;  $a_1 a_3$ ,  $a_2 a_4$  are pairs of another and  $a_1 a_4$ ,  $a_2 a_3$  are pairs of a third. The planes joining the points of  $R$  to the pairs of any involution which correspond to them form a cubic developable with a plane passing through  $R$ .

**209.** We now consider a curve  $C$  of the type III (E); two of its chords pass through  $O$  and it has a double point  $P$ ; it meets every  $\varpi$ -plane through  $O$  in four points and every  $\rho$ -plane through  $O$  in two points. The  $\rho$ -planes cut out a  $g_2^1$  on  $C$ , the two points on the different branches of  $C$  at  $P$  forming a pair of this  $g_2^1$ . Hence the chords of  $C$  lying in the  $\rho$ -planes form a quartic ruled surface. An arbitrary plane  $\rho$  meets the [4] containing  $C$  in a line and the  $\varpi$ -plane containing this line joins it to  $O$ ; this contains six chords of  $C$  meeting the line; the other three are generators of the quartic ruled surface. We thus have in [3] a double curve  $6R + G + C_3^0$ . There is a generator of the quartic ruled surface through  $P$ , and the two chords of  $C$  passing through  $O$  are also generators of this surface; hence  $C_3^0$  meets  $G$  once and  $R$  twice.

The chords of  $C$  which lie in the  $\varpi$ -planes form a ruled surface  $R_2^{14}$  on which  $C$  is a triple curve, there are six tangents of  $C$  which are generators of this surface. There are twelve points of  $C$  at which two of the three generators of  $R_2^{14}$  coincide (these do not include  $P$ ) so that the prime sections of  $R_2^{14}$  are elliptic curves. An arbitrary plane  $\varpi$  meets the [4] containing  $C$  in a line; one chord of  $C$  meeting this line is contained in the  $\rho$ -plane joining it to  $O$ , the other eight are all generators of  $R_2^{14}$ . Hence the surface in [3] has a bitangent developable  $R + G + E_8^1$ .

The two chords of  $C$  which lie on  $\Omega$  and pass through  $P$  are double generators of  $R_2^{14}$ ; there is another generator of  $R_2^{14}$  passing through  $P$ . Hence  $G$  lies in two double planes and one ordinary plane of  $E_8^1$ .

Also  $R$  is an axis of  $E_8^1$ ; the two planes of  $E_8^1$  through  $R$  are represented on  $\Omega$  by the  $\rho$ -planes containing the two chords of  $C$  which pass through  $O$ . Again, the two trisecants of  $C$  are joined to  $O$  by  $\pi$ -planes, so that  $E_8^1$  has two triple planes represented on  $\Omega$  by the  $\rho$ -planes through these trisecants.

**210.** To obtain this last surface by projection we choose  $S$  to meet the [4] containing a directrix quartic  $E$  of the general normal surface  $F$  in a plane; the chords of  $E$  meet this plane in an elliptic cubic  $\mathfrak{D}_3^1$ . But further,  $S$  is so chosen that it lies in a [5] with a pair of generators  $g, g'$ ; the chord of  $E$ , joining the points where it is met by  $g$  and  $g'$ , meets  $\mathfrak{D}_3^1$ .  $S$  meets the solid containing  $g$  and  $g'$  in a line  $l$  meeting  $\mathfrak{D}_3^1$ . It meets  $M_5^6$  in  $\mathfrak{D}_3^1$ ,  $l$  and a conic  $\mathfrak{D}_2$ , meeting  $l$  once and  $\mathfrak{D}_3^1$  twice. The chords of  $F$  which meet  $S$  meet  $F$  in the quartic  $E$  counted three times,  $g, g'$  and a prime section  $C_6^0$ . The intersections of  $C_6^0$  with  $g$  and  $g'$  lie on the chord of  $F$  through the intersection of  $l$  and  $\mathfrak{D}_2$ ; the four intersections of  $C_6^0$  and  $E$  lie in pairs on the chords of  $F$  through the intersections of  $\mathfrak{D}_3^1$  and  $\mathfrak{D}_2$ . Hence on projection we have a surface with a double curve  $6R + G + C_3^0$  precisely as in the last article.

To generate  $f$  we take a line  $R$  and a twisted cubic in (1, 4) correspondence with a united point; one of the points of the line gives rise to four points of the cubic which include the two intersections of the cubic with the chord  $G$  through it.

**211.** If we have on  $\Omega$  a curve  $C$  of the type III (F) we have in [3] a ruled surface whose double curve is  $R + G + C_8^1$  and bitangent developable  $6R + G + E_3^0$ .  $R$  and  $G$  intersect;  $C_8^1$  has two triple points, it has also two double points which lie on  $G, G$  meeting it in a further point;  $R$  is a chord of  $C_8^1$ . There is a plane of  $E_3^0$  passing through  $G$  and there are two planes of  $E_3^0$  passing through  $R$ .

In order to obtain this surface by projection we take the normal surface  $F$  with a directrix conic  $\Gamma$ , and project from a solid  $S$  which meets the plane of  $\Gamma$  in a point  $O$ . But here  $S$  is further chosen to lie in a [5] with two generators  $g, g'$  of  $F$ . Since this [5] is not to be the same as the [5] through  $S$  and  $\Gamma$  the two points in which  $\Gamma$  is met by  $g$  and  $g'$  must lie on a line through  $O$ . The solid  $gg'$  meets  $S$  in a line  $l$  passing through  $O$ .

$S$  meets  $M_5^6$  in  $l$  and an elliptic quintic  $\mathfrak{D}_5^1$  with a double point at  $O$ ,  $\mathfrak{D}_5^1$  meeting  $l$  again. The chords of  $F$  which meet  $S$  meet  $F$  in  $\Gamma, g, g'$  and  $C_{16}$ , where  $C_{16}$  meets every generator in three points. There are six tangents of  $C_{16}$  meeting  $\mathfrak{D}_5^1$ , so that it is of genus 4; it has eight double points. Six of these double points lie three in each of two planes

meeting  $S$  in lines; if  $A$  is either of the others then we have two chords  $AB$  and  $AC$  of  $C_{16}$  meeting  $\vartheta_5^1$ ; the chord  $BC$  meets  $l$ ,  $B$  being on  $g$  and  $C$  on  $g'$ . Thus we account for two intersections of  $C_{16}$  with each of  $g$  and  $g'$ ; the remaining pair is given by the chord of  $F$  through the intersection of  $l$  and  $\vartheta_5^1$ .  $C_{16}$  meets  $\Gamma$  in four points which lie on two lines through  $O$ . Hence on projection we have in  $\Sigma$  a surface  $f$  with a double curve  $R + G + C_8^1$  as required.

To generate  $f$  we take a line and a rational quartic in (1, 2) correspondence; one point of the line giving rise to a pair of points of the quartic collinear with it. The joins of the pairs of points on the quartic form a cubic ruled surface, one generator of which passes through the point of the line which gives rise to it; hence the bitangent planes of the surface which do not pass through the line or the double generator form a developable of the third class  $E_3^0$ , two of whose planes pass through the line. We have a bitangent developable  $6R + G + E_3^0$ .

**212.** Take a point  $O$  in the plane of  $\Gamma$ ; then a line  $l$  passing through  $O$  and lying in the solid which contains a pair of generators  $g, g'$  meeting  $\Gamma$  in a pair of points collinear with  $O$ ; then a conic  $\vartheta_2$  passing through  $O$ , this being the directrix conic of the quintic ruled surface determined by an involution on a prime section  $C_6^0$  of  $F$ . Project on to  $\Sigma$  from the solid  $S$  containing  $l$  and  $\vartheta_2$ .

$S$  meets  $M_5^0$  in  $l, \vartheta_2$  and a twisted cubic  $\vartheta_3^0$ ;  $\vartheta_3^0$  passes through  $O$ , meeting  $l$  once again and  $\vartheta_2$  twice again. The chords of  $F$  meeting  $S$  meet  $F$  in  $\Gamma, g, g', C_6^0$  and an elliptic curve  $C_{10}^1$ , four of whose tangents meet  $\vartheta_3^0$ ;  $C_{10}^1$  meeting every generator in two points. It meets  $\Gamma$  twice and  $C_6^0$  ten times, having two double points.

A pair of intersections of  $C_{10}^1$  with  $g$  and  $g'$  are given by the chord of  $F$  through the intersection (other than  $O$ ) of  $l$  and  $\vartheta_3^0$ ; the other pair is associated with the intersections of  $C_6^0$  with  $g$  and  $g'$  and with two intersections of  $C_6^0$  with  $C_{10}^1$ . Four other intersections of  $C_6^0$  and  $C_{10}^1$  are given by the chords of  $F$  passing through the two intersections (other than  $O$ ) of  $\vartheta_2$  and  $\vartheta_3^0$ ; the remaining four are associated in pairs with the two double points of  $C_{10}^1$ .

Hence on projection we have a surface  $f$  with a double curve  $R + G + C_3^0 + C_5^0$ .  $R$  meets each of  $G, C_3^0$  and  $C_5^0$ ;  $G$  is a chord of  $C_3^0$  and  $C_5^0$  passes through their two intersections, meeting  $G$  in one other point;  $C_5^0$  has two double points through each of which  $C_3^0$  passes,  $C_3^0$  and  $C_5^0$  having two further intersections.

The surface is generated by a line  $R$  and a twisted cubic  $C_3^0$  in (2, 2) correspondence with a doubly united point; one point of the line giving rise to the intersections of  $C_3^0$  with the chord which passes through it.

The lines joining the pairs of points of the cubic form a quartic ruled surface, there being two generators of this surface passing through points of the line which give rise to them. Hence the bitangent planes of the surface which do not pass through  $R$  or  $G$  form a developable  $E_3^0$ , two of whose planes pass through  $R$ . We have thus a bitangent developable  $6R + G + E_3^0$ ; two planes of  $E_3^0$  passing through  $R$  and one through  $G$ , while  $R$  and  $G$  are co-planar.

We have also the dual surface whose double curve is  $6R + G + C_3^0$  and bitangent developable  $R + G + E_3^0 + E_5^0$ . To generate this surface we take a line  $R$  and a twisted cubic in (1, 4) correspondence with a united point, one of the points of the line giving rise to four points of the cubic which include the two intersections of the cubic with the chord  $G$  through it. But here the  $g_4^1$  on the cubic is such that each of its sets consists of two sets of an involution (which does not contain the extremities of this chord as a pair of corresponding points). The joins of the pairs of the involution give a regulus, and there is a (1, 2) correspondence between the points of  $R$  and the lines of this regulus, there being one united element. We thus have a developable  $E_3^0$ , two of whose planes pass through  $G$  and one through  $R$ .

**213.** Now take  $S$  to meet the plane of  $\Gamma$  in a point  $O$  and to contain an axis  $\lambda$  of a directrix quartic  $E$ . The lines through  $O$  in the plane of  $\Gamma$  determine an involution of pairs of generators of  $F$ , while the generators of the cubic ruled surface having  $\lambda$  for its directrix determine another. If  $g, g'$  is the pair of generators common to these two involutions the solid  $gg'$  meets  $S$  in a line  $l$  passing through  $O$  and meeting  $\lambda$ .

$S$  meets  $M_5^6$  in  $\lambda, l$  and a rational quartic  $\vartheta_4^0$  which has a double point at  $O$  and meets  $\lambda$  twice. The chords of  $F$  meeting  $S$  meet  $F$  in  $E, \Gamma, g, g'$ , and an elliptic curve  $C_{12}^1$ , four of whose tangents meet  $\vartheta_4^0$ ;  $C_{12}^1$  meets every generator in two points. It has four double points, and meets  $\Gamma$  in four points (collinear with  $O$  in pairs) and  $E$  in eight points. Four of these intersections with  $E$  are given by the chords of  $F$  through the two intersections of  $\lambda$  and  $\vartheta_4^0$ ; the remaining four are associated in pairs with two of the double points of  $C_{12}^1$ . The other two double points are associated with the intersections of  $C_{12}^1$  with  $g$  and  $g'$ .

Thus on projection from  $S$  we have in  $\Sigma$  a surface  $f$  whose double curve is  $R + G + C_2 + C_6^0$ .  $R$  meets  $G$  and is a chord of  $C_6^0$ , while  $G$  and  $C_2$  intersect.  $C_6^0$  has four double points, two of which are on  $G$ ; the other two are on  $C_2$ , which meets  $C_6^0$  in two other points.

To generate  $f$  we take a line  $R$  and a conic  $C_2$  in (2, 2) correspondence; to a branch point of the correspondence on  $R$  there corresponds a branch point of the correspondence on  $C_2$  and the line  $G$  joining these points is a



double generator. The joins of the pairs of points of  $C_2$  which correspond to the points of  $R$  touch another conic; there are two tangents of this latter passing through the point in which  $R$  meets the plane of  $C_2$ . Hence those bitangent planes of the surface which do not pass through  $R$  or  $G$  give a developable  $E_3^0$ , two of whose planes pass through  $R$  and one through  $G$ . The bitangent developable is  $6R + G + E_3^0$ .

There is also the dual surface whose double curve is  $6R + G + C_3^0$  and bitangent developable  $R + G + E_2 + E_6^0$ . To generate this surface we take a line  $R$  and a twisted cubic in (1, 4) correspondence with a united point; one of the points of  $R$  gives rise to a set of four points on the cubic which includes its intersections with the chord through this point of  $R$ . And here the  $g_4^1$  on the cubic is such that each of its sets consists of two pairs of points of an involution, which includes the intersections of the cubic with the chord just mentioned as a pair of corresponding points. The joins of the pairs of the involution give a regulus and there is a (1, 2) correspondence with two united elements between the points of  $R$  and the lines of this regulus. We thus have a quadric cone with one of its tangent planes passing through the double generator.

214. Take again the normal surface  $F$  with a directrix conic in [7], and an involution of pairs of generators giving a quintic ruled surface through a prime section  $C_6^0$ ; the directrix conic  $\vartheta_2$  of this surface meeting the plane of  $\Gamma$  in a point  $O$ . It is possible to take a solid  $S$  through  $\vartheta_2$  which also contains an axis  $\lambda$  of a directrix quartic  $E$ , where  $\lambda$  meets  $\vartheta_2$ , and a line  $l$  through  $O$ , meeting  $\lambda$ , which lies in a solid containing a pair of generators  $g, g'$  of  $F$ .

$S$  meets  $M_5^6$  in  $l, \lambda, \vartheta_2$  and another conic  $\phi_2$  passing through  $O$ ;  $\phi_2$  meeting  $\vartheta_2$  again and also meeting  $\lambda$ . The chords of  $F$  meeting  $S$  meet  $F$  in  $\Gamma, g, g', E, C_6^0$  and another prime section  $D_6^0$ . It is easily seen how the mutual intersections of these curves arise; so that on projection we have a surface whose double curve is  $R + G + C_2 + C_3^0 + D_3^0$ .  $R$  meets  $G, C_3^0$  and  $D_3^0$ , while  $G$  meets  $C_2$  and passes through two intersections of  $C_3^0$  and  $D_3^0$ . There are two points common to all of  $C_2, C_3^0, D_3^0$ ;  $C_2$  meeting each of  $C_3^0$  and  $D_3^0$  in one further point; also there is one more intersection of  $C_3^0$  and  $D_3^0$ . We have again a bitangent developable  $6R + G + E_3^0$ .

The dual surface has a double curve  $6R + G + C_3^0$  and a bitangent developable  $R + G + E_2 + E_3^0 + F_3^0$ . It is generated by a line  $R$  and a twisted cubic in (1, 4) correspondence with a united point; one of the points of  $R$  giving rise to four points of the cubic which include the two intersections of the cubic with the chord  $G$  through it. Here, however, the  $g_4^1$  on the cubic is such that each of its sets consists of two pairs of points of three

different involutions\*. One of the involutions contains the ends of the chord  $G$  as a pair of corresponding points; this gives a developable  $E_2$  with a plane passing through  $G$ . Each of the other two involutions gives a developable of the third class with two planes through  $G$  and one through  $R$ , and we get the same two planes through  $G$  for each of these two developables.

**215.** Take now a curve  $C$  of the type III (G) with two double points  $P$  and  $Q$ ; it meets every  $\omega$ -plane through  $O$  in four points and every  $\rho$ -plane through  $O$  in two points, a chord of  $C$  passing through  $O$ . The  $\rho$ -planes cut out a  $g_2^1$  on  $C$  which includes the pairs of points on the two branches of  $C$  at  $P$  and on the two branches of  $C$  at  $Q$ . Hence the chords of  $C$  lying in the  $\rho$ -planes form a cubic ruled surface  $R_2^{3\dagger}$ . The chords of  $C$  lying in the  $\omega$ -planes through  $O$  form a ruled surface  $R_2^{13}$  on which  $C$  is a triple curve, there being six tangents of  $C$  which are generators of  $R_2^{13}$ . There are twelve points of  $C$  at which two of the three generators of  $R_2^{13}$  coincide (these do not include  $P$  or  $Q$ ); so that the prime sections of  $R_2^{13}$  are elliptic curves.

A  $\rho$ -plane of  $\Omega$  meets the [4] containing  $C$  in a line, and the  $\omega$ -plane through this line joins it to  $O$  and contains six chords of  $C$ . The two other chords meeting the line are generators of  $R_2^3$ , and we have in [3] a ruled surface whose double curve is  $6R + G + H + C_2$ . There are generators of  $R_2^3$  passing through  $P$  and  $Q$ , and the chord of  $C$  through  $O$  is also a generator of  $R_2^3$ . Hence  $C_2$  meets  $G$ ,  $H$ ,  $R$  each in one point.  $G$  and  $H$  both meet  $R$ .

A  $\omega$ -plane of  $\Omega$  meets the [4] containing  $C$  in a line, and the  $\rho$ -plane through this line joins it to  $O$  and contains a chord of  $C$ . The other seven chords meeting the line are generators of  $R_2^{13}$ . We have then for the bi-tangent developable of the surface  $R + G + H + E_7^1$ . The two chords of  $C$  which pass through  $P$  and lie on  $\Omega$  are double generators of  $R_2^{13}$ , which has also another generator passing through  $P$ . Hence there are two double planes and a simple plane of  $E_7^1$  passing through  $G$ . Likewise there are two double planes and a simple plane of  $E_7^1$  passing through  $H$ ; there is also a plane of  $E_7^1$  passing through  $R$ , represented on  $\Omega$  by the  $\rho$ -plane which contains the chord of  $C$  passing through  $O$ .

**216.** We now obtain this surface by projection. We take the general normal surface  $F$  in [7] with  $\infty^1$  directrix cubics  $\Delta$  and choose the solid  $S$  to meet the [4] containing a directrix quartic  $E$  in a plane. We also suppose that there are two pairs of generators  $g, g'$  and  $h, h'$  such that each

\* Cf. § 208.

† The projection of a quintic ruled surface in [6] from a line meeting two of its generators.

pair lies in a [5] with  $S$ . The solid  $gg'$  meets  $S$  in a line  $l$  and the solid  $hh'$  meets  $S$  in a line  $m$ . The chords of  $E$  meet  $S$  in the points of a plane cubic  $\mathfrak{D}_3^1$ , and as we assume that the [5] containing  $S$  and  $E$  does not contain a generator of  $F$  the two lines  $l$  and  $m$  must meet  $\mathfrak{D}_3^1$ .  $S$  meets  $M_5^6$  in  $\mathfrak{D}_3^1$ ,  $l$ ,  $m$  and a line  $\lambda$ , which is a transversal of  $l$  and  $m$ , meeting  $\mathfrak{D}_3^1$ . The chords of  $F$  meeting  $S$  meet  $F$  in  $E$  counted three times,  $g$ ,  $g'$ ,  $h$ ,  $h'$  and another directrix quartic  $E'$ . Hence we have on projection a surface  $f$  with a double curve  $6R + G + H + C_2$ ;  $C_2$  meeting  $G$ ,  $H$  and  $R$ , while  $G$  and  $H$  both meet  $R$ .

To generate  $f$  we take a line  $R$  and a twisted cubic in (1, 4) correspondence with a united point. There are two points of  $R$  giving rise to sets of four points on the cubic which include the two intersections of the cubic with the chords,  $G$  and  $H$ , which pass through the respective points.

217. We have in [3] the dual surface whose double curve is  $R + G + H + C_7^1$  and bitangent developable  $6R + G + H + E_2$ .  $G$  and  $H$  meet  $R$ , while there is a plane of  $E_2$  passing through each of  $G$ ,  $H$  and  $R$ .  $C_7^1$  has two double points and one ordinary point on each of  $G$  and  $H$  and it also meets  $R$ .

To obtain this surface by projection we take the normal surface  $F$  with a directrix conic  $\Gamma$ . Take two pairs of generators  $g$ ,  $g'$  and  $h$ ,  $h'$ , and let the chords of  $\Gamma$  determined by these pairs of generators meet in  $O$ . Take a line  $l$  through  $O$  lying in the solid  $gg'$  and a line  $m$  through  $O$  lying in the solid  $hh'$ ; then project from a solid  $S$  containing  $l$  and  $m$ .

$S$  meets  $M_5^6$  in  $l$ ,  $m$  and an elliptic quartic  $\mathfrak{D}_4^1$  passing through  $O$  and meeting each of  $l$  and  $m$  in a further point. The chords of  $F$  meeting  $S$  meet  $F$  in  $\Gamma$ ,  $g$ ,  $g'$ ,  $h$ ,  $h'$  and  $C_{14}$ , this latter curve meeting every generator in three points. There are six tangents of  $C_{14}$  meeting  $\mathfrak{D}_4^1$ , so that it is of genus 4; hence it has four double points.  $C_{14}$  meets  $\Gamma$  in two points collinear with  $O$ ; the chord of  $F$  through the intersection of  $l$  and  $\mathfrak{D}_4^1$  other than  $O$  passes through intersections of  $C_{14}$  with  $g$  and  $g'$ ; the remaining intersections of  $C_{14}$  with  $g$  and  $g'$  are associated in pairs with two of its double points. Similarly for its intersections with  $h$  and  $h'$ .

Hence on projection we have a surface with a double curve  $R + G + H + C_7^1$  precisely as required.

To generate this surface we take a line and a rational quartic in (1, 2) correspondence; two points of the line give rise to pairs of points of the quartic whose joins respectively pass through them. The joins of the pairs of points of the quartic form a cubic ruled surface, two of whose generators meet the line in the points which give rise to them; there is one other generator of the cubic ruled surface meeting the line. Hence the planes joining the points of the line to the pairs of points of the quartic which

correspond to them touch a quadric cone one of whose tangent planes passes through the line. There is also a tangent plane through each of the double generators. We have a bitangent developable  $6R + G + H + E_2$ .

**218.** Consider again the surface  $F$  with a directrix conic, and take two involutions of pairs of generators; the first of these gives a pencil of lines in the plane of  $\Gamma$  passing through a point  $O$ , while the second, by means of any directrix quartic  $E$ , gives a cubic ruled surface with a directrix line  $\lambda$ . There is a pair of generators  $g, g'$  common to the two involutions; the solid containing them meets  $\lambda$  and passes through  $O$ . Take a solid  $S$  containing  $O$  and  $\lambda$ ; it meets the solid  $gg'$  in a line  $l$  passing through  $O$  and meeting  $\lambda$ ; further, we choose  $S$  to meet the solid containing another pair of generators in a line  $m$  passing through  $O$ .

$S$  meets  $M_5^6$  in  $l, m, \lambda$  and a twisted cubic  $\vartheta_3^0$  passing through  $O$ ;  $\vartheta_3^0$  meets  $m$  again and meets  $\lambda$  twice. The chords of  $F$  meeting  $S$  meet  $F$  in  $\Gamma, g, g', h, h', E$  and an elliptic curve  $C_{10}^1$  four of whose tangents meet  $\vartheta_3^0$ .  $C_{10}^1$  meets every generator in two points and has two double points; it meets  $\Gamma$  in two points collinear with  $O$ , and  $E$  in six points. Four of these intersections with  $E$  are accounted for by the two chords of  $F$  through the intersections of  $\lambda$  and  $\vartheta_3^0$ ; either of the others is associated with an intersection of  $E$  with  $h$  or  $h'$  and with an intersection of  $C_{10}^1$  with  $h'$  or  $h$  respectively. The other intersections of  $C_{10}^1$  with  $h$  and  $h'$  are on the chord of  $F$  through the intersection (other than  $O$ ) of  $m$  and  $\vartheta_3^0$ ; the intersections of  $C_{10}^1$  with  $g$  and  $g'$  are associated in pairs with the double points.

Hence on projection from  $S$  we obtain a surface  $f$  with a double curve  $R + G + H + C_2 + C_5^0$ .  $R$  meets  $G, H$  and  $C_5^0$ .  $H$  is a chord of  $C_2$ , and  $C_5^0$  passes through their intersections meeting  $H$  in one other point and  $C_2$  in two other points.  $G$  meets  $C_2$  and passes through two double points of  $C_5^0$ .

The surface is generated by a line  $R$  and a conic  $C_2$  in (2, 2) correspondence. To the point in which  $R$  meets the plane of  $C_2$  corresponds a pair of points on  $C_2$  whose join  $H$  passes through this point, while there are mutually corresponding double points on  $R$  and  $C_2$ . The joins of the pairs of points of  $C_2$  which correspond to the points of  $R$  touch another conic; the planes joining the points of  $R$  to these pairs of points touch a quadric cone with a tangent plane passing through each of  $R, G$  and  $H$ . The bitangent developable is  $6R + G + H + E_2$ .

There is also in [3] the surface dual to this one whose double curve is given by  $6R + G + H + C_2$  and bitangent developable by  $R + G + H + E_2 + E_5^0$ . This surface is generated by a line  $R$  and a twisted cubic in (1, 4) correspondence as in § 216; there are two points of  $R$  giving rise to sets of four points on the cubic which include the two intersections of the cubic with

the chords  $G$  and  $H$  passing through these points. But here the  $g_4^1$  on the cubic is such that each of its sets consists of two pairs of an involution, which includes as a pair of corresponding points the intersections of the cubic with either  $G$  or  $H$ , say with  $G$ . The joins of the pairs of this involution form a regulus, and the planes joining the points of  $R$  to the corresponding pairs of the involution touch a quadric cone which has one tangent plane through  $G$  and two tangent planes through  $H$ .

**219.** Take an involution of pairs of generators on the normal surface  $F$  with a directrix conic  $\Gamma$ , and let this give a quintic ruled surface, through a prime section  $C_6^0$  of  $F$ , whose directrix conic  $\vartheta_2$  meets the plane of  $\Gamma$  in a point  $O$ . Then we can choose a solid  $S$  to contain  $\vartheta_2$ , to meet the solid containing a pair of generators  $g, g'$  in a line  $l$  through  $O$  and to meet the solid containing another pair of generators  $h, h'$  in a line  $m$  through  $O$ . Of course neither of the pairs  $g, g', h, h'$  need belong to the involution.  $S$  meets  $M_5^6$  in  $l, m, \vartheta_2$  and a conic  $\phi_2$  which meets  $l$  and  $m$  each once and  $\vartheta_2$  twice.

The chords of  $F$  meeting  $S$  meet  $F$  in  $\Gamma, g, g', h, h', C_6^0$  and an elliptic curve  $C_8^1$  four of whose tangents meet  $\phi_2$ . This curve meets every generator of  $F$  twice and has no double points.  $C_8^1$  does not meet  $\Gamma$ , but it meets  $C_6^0$  in eight points; four of these are given by the two chords of  $F$  through the intersections of  $\vartheta_2$  and  $\phi_2$ . The other intersections of  $C_8^1$  and  $C_6^0$  are associated with the intersections of  $C_6^0$  with  $g, g', h, h'$  and with intersections of  $C_8^1$  with  $g', g, h', h$  (one with each). The other intersections of  $C_8^1$  with these four generators are given by the chords of  $F$  through the intersections of  $\phi_2$  with  $l$  and  $m$ .

We thus have in  $\Sigma$  a surface  $f$  whose double curve is  $R + G + H + C_3^0 + C_4^0$ .  $R$  meets  $G, H$  and  $C_3^0$ .  $G$  and  $H$  are chords of  $C_3^0$ ;  $C_4^0$  passes through the four intersections, meeting each of  $G$  and  $H$  in one other point and  $C_3^0$  in two other points.

To generate this surface we take a line  $R$  and a twisted cubic  $C_3^0$  in (2, 2) correspondence with a doubly united point; there are two points of  $R$  which give rise to pairs of points of  $C_3^0$  on the chords  $G$  and  $H$  which pass through them. The pairs of points of  $C_3^0$  which correspond to the points of  $R$  are joined by lines forming a quartic ruled surface; the planes joining the points of  $R$  to these lines touch a quadric cone, there being tangent planes of this cone through  $G, H$  and  $R$ . We have a bitangent developable  $6R + G + H + E_2$ .

We have also the dual surface whose double curve is  $6R + G + H + C_2$  and bitangent developable  $R + G + H + E_3^0 + E_4^0$ . To generate this surface we take a line  $R$  and a twisted cubic in (1, 4) correspondence giving two double generators  $G$  and  $H$  as in § 216; but here the  $g_4^1$  on the

cubic is such that each of its sets consists of two pairs of an involution, which does not contain the pair of intersections of the cubic with either  $G$  or  $H$  as a pair of corresponding points. The pairs of points of this involution give lines forming a regulus; there is thus set up a (1, 2) correspondence between the points of  $R$  and the lines of this regulus, and the planes joining the points of  $R$  to the lines of the regulus give (because of the united point) the developable  $E_3^0$ , which has a plane passing through  $R$  and two planes through each of  $G$  and  $H$ .

220. Take the surface  $F$  with a directrix conic  $\Gamma$ , and any two of its directrix quartics  $E$  and  $E'$ . Take an axis  $\lambda$  of  $E$  and an axis  $\mu$  of  $E'$ , these intersecting on the common chord of  $E$  and  $E'$ . Then choose a solid  $S$  containing  $\lambda$  and  $\mu$  and meeting the plane of  $\Gamma$  in a point  $O$ .  $S$  also contains a line  $l$ , passing through  $O$  and meeting  $\lambda$ , which lies in the solid containing a pair of generators  $g, g'$ , and a line  $m$ , passing through  $O$  and meeting  $\mu$ , which lies in the solid containing a pair of generators  $h, h'$ .  $S$  meets  $M_5^6$  in  $l, m, \lambda, \mu$  and a conic  $\vartheta_2$  passing through  $O$  and meeting  $\lambda$  and  $\mu$ . The chords of  $F$  meeting  $S$  meet  $F$  in  $E, E', \Gamma, g, g', h, h'$  and a prime section  $C_6^0$ . On projection from  $S$  we have in  $\Sigma$  a surface whose double curve is  $R + C_2 + D_2 + G + H + C_3^0$ .  $R$  meets  $G, H$  and  $C_3^0$ .  $C_2$  meets  $G$  and has  $H$  for a chord,  $C_3^0$  passing through the intersections of  $H$  with  $C_2$  and meeting  $C_2$  again.  $D_2$  meets  $H$  and has  $G$  for a chord,  $C_3^0$  passing through the intersections of  $G$  with  $D_2$  and meeting  $D_2$  again.  $C_2$  and  $D_2$  have one intersection.

There is also the dual surface whose double curve is  $6R + G + H + C_2$  and bitangent developable  $R + E_2 + F_2 + G + H + E_3^0$ . This is generated by a line  $R$  and a twisted cubic in (1, 4) correspondence as in § 216, but here the  $g_4^1$  on the cubic is such that each of its sets consists of two pairs of points of three different involutions. One of these involutions contains the pair of points in which  $G$  meets the cubic; the regulus arising from this involution gives, when taken with  $R$ , the developable  $E_2$  which has a plane through  $G$  and two planes through  $H$ . Similarly another of the involutions contains the pair of points in which  $H$  meets the cubic; we have then the developable  $F_2$  with a plane through  $H$  and two planes through  $G$ . The third involution does not contain the extremities either of  $G$  or of  $H$ ; we thus have the developable  $E_3^0$  with a plane through  $R$  and two planes through each of  $G$  and  $H$ . The planes of  $E_3^0$  through  $G$  are the same as the planes of  $F_2$  through  $G$ , and the planes of  $E_3^0$  through  $H$  are the same as the planes of  $E_2$  through  $H$ .

221. We now proceed to consider the curve  $C$  of the type III (I) on  $\Omega$ ; it meets every plane of  $\Omega$  through  $O$  in three points and has four of its chords passing through  $O$ . The chords of  $C$  which lie in either system of

planes form a ruled surface  $R_2^{10}$  on which  $C$  is a double curve, and there are four tangents of  $C$  which are generators of this ruled surface. Also there are four points of  $C$  at which the two generators of the ruled surface coincide; and the prime sections of  $R_2^{10}$  are rational curves.

Any plane of  $\Omega$  meets the [4] containing  $C$  in a line, and the plane of the opposite system through this line joins it to  $O$  and contains three chords of  $C$ ; there are seven other chords of  $C$  meeting the line. Hence we have in [3] a ruled surface whose double curve is  $3R + C_7^0$  and bitangent developable  $3R + E_7^0$ .  $C_7^0$  meets  $R$  in four points; these are represented on  $\Omega$  by the  $\omega$ -planes through the four chords of  $C$  which pass through  $O$ ; similarly  $E_7^0$  has four planes passing through  $R$ .

Of the four trisecants of  $C$  two lie in  $\omega$ -planes and two in  $\rho$ -planes;  $C_7^0$  has two triple points and  $E_7^0$  two triple planes. The section of  $R_2^{10}$  by a solid is a rational curve of order 10 lying on a quadric; it meets all generators of one system in three points and all of the other system in seven points, having six double points and two triple points.

**222.** In order to obtain this surface by projection we take in [7] the normal surface  $F$  with  $\infty^1$  directrix cubics and project from a solid  $S$  which meets the solid  $K$  containing one of these cubics  $\Delta$  in a line  $l$ . Then  $S$  meets the  $M_4^4$  formed by the solids  $K$  in two further points\* so that the surface  $f$  in  $\Sigma$  has two tritangent planes not passing through its directrix. Its directrix is the line  $R$  in which  $\Sigma$  is met by the [5] containing  $S$  and  $K$ ; through any point of  $R$  there pass three generators of  $F$ , while any plane through  $R$  contains three generators of  $F$ .

$S$  meets  $M_5^6$  in  $l$  and a rational quintic  $\mathfrak{D}_5^0$  of which  $l$  is the quadrisecant. There are two chords of  $\Delta$  passing through any point of it and meeting  $l$ , while there are four tangents of  $\Delta$  meeting  $l$ . Hence the chords of  $F$  meeting  $S$  meet  $F$  in  $\Delta$  counted twice and an elliptic curve  $C_{14}^1$ , four of whose tangents meet  $\mathfrak{D}_5^0$ .  $C_{14}^1$  meets every generator of  $F$  in two points and has six double points. It meets  $\Delta$  in eight points; these lie two on each of the four chords of  $F$  through the intersections of  $l$  and  $\mathfrak{D}_5^0$ . The six double points lie three in each of two planes which meet  $S$  in lines.

Hence on projection we have a surface with a double curve  $3R + C_7^0$ ;  $C_7^0$  has two triple points and meets  $R$  four times.

This surface is generated by a line  $R$  and a twisted cubic in (1, 3) correspondence.

**223.** We now pass to the curve  $C$  on  $\Omega$  of the type III (J); the chords of  $C$  which lie in either system of planes include one trisecant and four tangents and form a ruled surface  $R_2^9$  whose prime sections are rational

\* A [5] through  $S$  meets  $M_4^4$  in a quartic ruled surface, and any solid through a generator of this surface meets it in two other points.

curves. The ruled surface  $R_2^9$  has  $C$  as a double curve, the trisecant as a triple generator and also a double generator passing through the double point of  $C$ .

We have in [3] a surface whose double curve is  $3R + G + C_6^0$  and bitangent developable  $3R + G + E_6^0$ .  $C_6^0$  has a triple point and meets  $R$  three times; it has a double point on  $G$  and meets  $G$  in one other point. Similarly for  $E_6^0$ .  $R$  and  $G$  intersect.

To obtain the surface by projection we choose the solid  $S$  to meet the solid  $K$  containing a directrix cubic  $\Delta$  of  $F$  in a line  $l$  and also to meet the solid containing a pair of generators  $g, g'$  in a line  $m$ . Since  $S$  is not to meet  $F$ ,  $l$  and  $m$  must intersect, the chord of  $\Delta$  through their intersection meeting  $\Delta$  in its intersections with  $g$  and  $g'$ . The curves  $\Delta$  meet  $g$  and  $g'$  in related ranges whose joins form a regulus; this meets  $m$  in two points, including its intersection with  $l$ . Hence there is only one solid  $K$  meeting  $S$  in a point not on  $l$  or  $m$ ; so that the projected surface has one tritangent plane passing neither through its directrix nor its double generator.

$S$  meets  $M_5^6$  in  $l, m$  and a rational quartic  $\vartheta_4^0$  meeting  $m$  and having  $l$  as a trisecant. The chords of  $F$  meeting  $S$  meet  $F$  in  $\Delta$  counted twice,  $g, g'$  and an elliptic curve  $C_{12}^1$  four of whose tangents meet  $\vartheta_4^0$ .  $C_{12}^1$  meets every generator of  $F$  in two points and has four double points. It meets  $\Delta$  in six points which lie two on each of the chords of  $F$  which pass through the three intersections of  $l$  and  $\vartheta_4^0$ . The chord through the intersection of  $m$  and  $\vartheta_4^0$  gives intersections of  $C_{12}^1$  with  $g$  and  $g'$ ; the line joining the other pair of intersections does not meet  $\vartheta_4^0$  but gives two chords of  $F$  which do so, these intersecting in a double point of  $C_{12}^1$ . The remaining three double points of  $C_{12}^1$  lie in a plane passing through a trisecant of  $\vartheta_4^0$ .

Hence on projection we have a surface with a double curve  $3R + G + C_6^0$ , precisely as required.

To generate this surface take a line  $R$  and a twisted cubic in (1, 3) correspondence; one point of  $R$  giving rise to a triad of points on the cubic which include the two intersections with the chord  $G$  through the point.

**224.** When  $C$  is of the type III (K) the chords which lie in either system of planes include four tangents and form a ruled surface  $R_2^8$ ; this has  $C$  for a double curve and has two double generators, one through each double point of  $C$ ; there is one other generator through each of these double points. The prime sections of  $R_2^8$  are rational curves. We have in [3] a ruled surface whose double curve is  $3R + G + H + C_5^0$  and bitangent developable  $3R + G + H + E_5^0$ .  $C_5^0$  has a double point and a simple point on  $G$ , while  $E_5^0$  has a double plane and a simple plane through  $G$ ; and similarly for  $H$ .  $R$  meets  $G$  and  $H$ . Also  $R$  is a chord of  $C_5^0$ ; the two points of intersection are represented on  $\Omega$  by the  $\omega$ -planes which contain



the two chords of  $C$  passing through  $O$ . Similarly there are two planes of  $E_5^0$  passing through  $R$ .

To obtain this surface by projection take a line  $l$  in a solid containing a directrix cubic  $\Delta$  of the normal surface  $F$ . Through a given point  $A$  of  $l$  there passes a chord of  $\Delta$ ; through the intersections of  $\Delta$  with this chord we have a pair of generators  $g, g'$  of  $F$ , and we can take a line  $m$  through  $A$  lying in the solid containing these two generators. Similarly we have a line  $n$  meeting  $l$  in a point  $B$  and lying in the solid containing a pair of generators  $h, h'$  of  $F$ . Then take  $S$  to be the solid containing  $l, m$  and  $n$ .  $S$  meets  $M_5^6$  in  $l, m, n$  and a cubic  $\mathfrak{D}_3^0$  which meets  $m$  and  $n$  and has  $l$  for a chord.

The chords of  $F$  meeting  $S$  meet  $F$  in  $\Delta$  counted twice,  $g, g', h, h'$  and an elliptic curve  $C_{10}^1$ , four of whose tangents meet  $\mathfrak{D}_3^0$ .  $C_{10}^1$  meets every generator of  $F$  in two points and has two double points.  $C_{10}^1$  meets  $\Delta$  in four points; these are on the chords of  $F$  through the two intersections of  $l$  and  $\mathfrak{D}_3^0$ . The chord of  $F$  through the intersection of  $m$  and  $\mathfrak{D}_3^0$  accounts for an intersection of  $C_{10}^1$  with each of  $g$  and  $g'$ ; the other pair of intersections is associated with a double point of  $C_{10}^1$ . Similarly for the intersections with  $h$  and  $h'$ .

Hence on projection we have a surface with a double curve as required.

To generate this surface we take a line  $R$  and a twisted cubic in (1, 3) correspondence; there are two points of  $R$  for which the corresponding triad of points on the cubic contains the two intersections with the chord through the point.

225. When  $C$  is of the type III (L) the chords which lie in either system of planes form a ruled surface  $R_2^7$  on which  $C$  is a double curve; the prime sections of  $R_2^7$  are rational curves. The triple point  $P$  of  $C$  represents a triple generator  $G$  of the ruled surface in [3]; the ruled surface  $R_2^7$  has three generators passing through  $P$ .

The ruled surface in [3] has a double curve  $3R + 3G + C_4^0$  and a bitangent developable  $3R + 3G + E_4^0$ .  $C_4^0$  meets  $R$  and has  $G$  for a trisecant, while  $R$  and  $G$  intersect.  $E_4^0$  has one plane passing through  $R$  and three planes passing through  $G$ .

The [5] containing three generators  $g, g', g''$  of the normal surface  $F$  meets the solid containing one of its directrix cubics  $\Delta$  in a plane; take  $S$  to lie in the [5] and meet the plane in a line  $l$ . The solids  $g'g'', g''g, gg'$  meet  $S$  in lines  $m, m', m''$  which all meet  $l$ .  $S$  meets  $M_5^6$  in  $l, m, m', m''$  and a conic  $\mathfrak{D}_2$  meeting these four lines each in one point. There are two chords of  $F$  through any point of  $g, g', g''$  or  $\Delta$  which meet  $S$ ; hence the chords of  $F$  meeting  $S$  meet  $F$  in  $\Delta, g, g', g''$  each counted twice and an elliptic curve  $C_8^1$  four of whose tangents meet  $\mathfrak{D}_2$ .  $C_8^1$  meets every generator of  $F$  in two points and has no double points.

On projection we have a surface with a double curve  $3R + 3G + C_4^0$  as required.

The cubics of  $F$  meet  $g, g', g''$  in related ranges; hence the solids  $K$  meet the [5] containing these generators in planes of a  $V_3^3$ .  $S$  meets the  $M_4^4$  formed by these solids  $K$  in  $l$  and the conic  $\mathfrak{D}_2$ ; the projected surface  $f$  contains  $\infty^1$  plane cubics whose planes pass through  $G$ ; these cubics have double points at those intersections of the planes with  $C_4^0$  which are not on  $G$ .

The surface is generated by a line  $R$  and a rational plane cubic in (1, 3) correspondence; to the point in which  $R$  meets the plane of the cubic correspond three points of the cubic collinear with it.

### *Surfaces with a directrix line which is also a generator*

**226.** Take a curve  $C$  on  $\Omega$  of the type IV (A); it gives in [3] a ruled surface with a directrix line  $R$  which is also a generator; through any point of  $R$  there pass four other generators, while any plane through  $R$  contains one other generator. Thus the double curve is the line  $R$  counted ten times, while the bitangent developable consists of the pencil of planes through  $R$  and a developable of the ninth class. The four trisecants of  $C$  are all contained in  $\omega$ -planes, so that this last developable has four triple planes represented on  $\Omega$  by the  $\rho$ -planes through the trisecants. The developable will then necessarily be elliptic. We have a double curve  $10R$  and a bitangent developable  $R + E_9^1$ , where  $E_9^1$  has four triple planes. There are three planes of  $E_9^1$  passing through  $R$ ; there is a  $\omega$ -plane through the tangent of  $C$  at  $O$ , and these three planes of  $E_9^1$  are represented on  $\Omega$  by the  $\rho$ -planes which contain the three chords of  $C$  joining  $O$  to the other three intersections of  $C$  with the  $\omega$ -plane through its tangent at  $O$ .

To obtain this surface by projection we choose  $S$  to lie in the [5] determined by a directrix quartic  $E$  and a generator  $g$  of the normal surface. This [5] meets  $\Sigma$  in a line  $R$  which is a directrix and also a generator of  $f$ ; through any point of  $R$  there pass four other generators, while any plane through  $R$  contains one other generator. The surface has four tritangent planes since  $S$  meets four of the solids  $K$ . It is generated by a line and a twisted cubic with two intersections placed in (1, 4) correspondence with one united point.

**227.** Dually, when  $C$  is of the type IV (B), we have in [3] a ruled surface whose double curve is  $R + C_9^1$  and bitangent developable  $10R$ ;  $C_9^1$  has  $R$  for a trisecant and has four triple points.

To obtain this surface by projection we take the normal surface  $F$  in [7] which has a directrix line  $\lambda$ , and then choose  $S$  to meet the plane containing  $\lambda$  and some generator  $g$  in a point  $O$ . The [5] joining  $S$  to  $\lambda$  also contains  $g$ , and meets  $\Sigma$  in a line  $R$  which is a directrix and also a

generator of  $f$ ; through any point of  $R$  there passes one other generator, while any plane through  $R$  contains four other generators.

The  $M_5$  containing the chords of  $F$  is made up of the  $\infty^3$  solids joining  $\lambda$  to the pairs of generators of  $F$ , and meets an arbitrary [5] in the chords of a rational normal quintic  $c_5$ . Hence\* the chords of  $F$  form an  $M_5^6$  which meets an arbitrary solid  $S$  in a sextic curve  $\mathfrak{D}_6^3$  of genus 3. Since the quintic curve  $c_5$  is a triple curve on the locus of its chords, any plane joining  $\lambda$  to a generator is a triple plane on  $M_5^6$ . We thus have  $\infty^1$  planes through  $\lambda$  forming an  $M_3^5$  which is a triple locus on  $M_5^6$ .

The tangent solids of  $F$  all pass through  $\lambda$  and meet the [5] in the tangents of  $c_5$ ; they thus form a locus  $M_4^8$ , there being eight of them which meet an arbitrary solid  $S$ . If, however, a line in [5] meets  $c_5$  there are only six further tangents of  $c_5$  meeting the line; so that if  $S$  meets a plane of  $M_3^5$  in a point  $O$  there are only six tangent solids of  $F$  meeting  $S$  in points other than  $O$ .

Thus the solid  $S$  from which we are to project meets  $M_5^6$  in an elliptic† sextic  $\mathfrak{D}_6^1$  with a triple point at  $O$ . The chords of  $F$  meeting  $S$  meet  $F$  in  $\lambda$ ,  $g$  and a curve  $C_{18}$ ; since there are six tangents of  $C_{18}$  meeting  $\mathfrak{D}_6^1$  it is of genus 4. It meets every generator of  $F$  in three points and  $\lambda$  in three points, and has twelve double points. These lie three in each of four planes containing trisecants of  $\mathfrak{D}_6^1$ . The intersections of  $C_{18}$  with  $g$  are in perspective from  $O$  with its intersections with  $\lambda$ .

Hence on projection we have a surface  $f$  with a double curve  $R + C_9^1$ ;  $C_9^1$  has  $R$  for a trisecant and has four triple points.

The surface is generated by a line  $R$  and a rational quintic with one intersection, these being placed in (1, 1) correspondence without a united point.

**228.** Suppose now, still considering the normal surface with a directrix line, that we take the directrix conic  $\mathfrak{D}_2$  of a quintic ruled surface determined by an involution on a prime section  $C_6^0$  of  $F$ . There is a generator of the quintic ruled surface passing through the intersection of  $\lambda$  and  $C_6^0$ ; this meets  $\mathfrak{D}_2$  in a point  $O$  and  $C_6^0$  again on a generator  $g$  of  $F$ . Thus  $O$  lies in

\* Cf. § 95. If  $F$  is generated by a (1, 1) correspondence between  $\lambda$  and one of its directrix quintic curves (there are  $\infty^5$  such curves on  $F$ ) its equations can be written

$$\frac{x_0}{x_1} = \frac{x_1}{x_2} = \frac{x_2}{x_3} = \frac{x_3}{x_4} = \frac{x_4}{x_5} = \frac{x_5}{x_6} = \frac{x_6}{x_7},$$

and the equations of  $M_5^6$  are

$$\left\| \begin{array}{cccc} x_0 & x_1 & x_2 & x_3 \\ x_1 & x_2 & x_3 & x_4 \\ x_2 & x_3 & x_4 & x_5 \end{array} \right\| = 0.$$

† If we take a plane through a point of a rational quintic curve in [4], it meets the locus  $V_3^6$  of § 95 in a sextic curve with a triple point at this point. This sextic curve has double points at the six remaining intersections of the plane with  $F_2^7$ , so that it is an *elliptic* curve.

the plane containing  $\lambda$  and  $g$ ; so that if  $S$  is chosen to contain  $\vartheta_2$  it automatically meets the plane containing  $\lambda$  and  $g$  in a point  $O$  of  $\vartheta_2$ .

$S$  meets  $M_5^6$  in  $\vartheta_2$  and a rational quartic  $\vartheta_4^0$  having a double point at  $O$  and meeting  $\vartheta_2$  in two other points. The chords of  $F$  which meet  $S$  meet  $F$  in  $\lambda$ ,  $g$ ,  $C_6^0$  and an elliptic curve  $C_{12}^1$ , four of whose tangents meet  $\vartheta_4^0$ ;  $C_{12}^1$  meets every generator of  $F$  in two points and  $\lambda$  in two points, and has four double points. Its intersections with  $\lambda$  are joined by lines through  $O$  to its intersections with  $g$ . It meets  $C_6^0$  in twelve points; four of these are given by the chords of  $F$  through the two intersections (other than  $O$ ) of  $\vartheta_2$  and  $\vartheta_4^0$ ; the remaining ones fall into four pairs associated with the four double points of  $C_{12}^1$ .

On projection we have a surface  $f$  with a double curve  $R + C_3^0 + C_6^0$ .  $R$  meets  $C_3^0$  and is a chord of  $C_6^0$ ;  $C_6^0$  has four double points through all of which  $C_3^0$  passes, and it meets  $C_3^0$  in two other points.

The surface is generated by a line  $R$  and a twisted cubic  $C_3^0$  in (2, 1) correspondence with a united point.

We have also in [3] the dual surface whose double curve is  $10R$  and bitangent developable  $R + E_3^0 + E_6^0$ . This is generated by a line  $R$  and a twisted cubic placed in (1, 4) correspondence with one united point,  $R$  being a chord of the cubic; but here the  $g_4^1$  on the cubic is such that each of its sets consists of two pairs of an involution. The joins of pairs of this involution form a regulus, and we have a (1, 2) correspondence between the points of  $R$  and the lines of this regulus; hence, because of the united point, the planes joining the points of  $R$  to the corresponding lines of the regulus form a developable  $E_3^0$ , one of whose planes passes through  $R$ .

**229.** We have seen that the directrix conics of all the quintic ruled surfaces given by involutions on prime sections of  $F$  necessarily meet one of the planes of  $M_3^5$ . Let us enquire whether we can project from a solid  $S$  containing two of these conics  $\vartheta_2$  and  $\phi_2$ .  $\vartheta_2$  meets the plane containing  $\lambda$  and a generator  $g$  in a point; hence, since the directrix  $R$  of the surface in  $\Sigma$  is to be a simple and not a double generator of  $f$ ,  $\phi_2$  must meet the same plane in a point. Moreover, since  $S$  is not to meet  $F$ ,  $\vartheta_2$  and  $\phi_2$  must meet this plane in the same point  $O$ .  $\vartheta_2$  and  $\phi_2$  have a second intersection through which passes a common generator of the two quintic ruled surfaces to which they belong.

$S$  meets  $M_5^6$  in  $\vartheta_2$ ,  $\phi_2$  and a third conic  $\psi_2$  passing through  $O$  and meeting each of  $\vartheta_2$  and  $\phi_2$  in one further point. The chords of  $F$  meeting  $S$  meet  $F$  in  $\lambda$ ,  $g$  and three prime sections  $B_6^0$ ,  $C_6^0$ ,  $D_6^0$ . Projecting from  $S$  we obtain in  $\Sigma$  a surface with a double curve  $R + B_3^0 + C_3^0 + D_3^0$ .  $R$  meets each cubic in one point; there are four points common to all the cubics, while any two of them have one further intersection.

There is also in [3] the dual surface whose double curve is  $10R$  and bitangent developable  $R + E_3^0 + F_3^0 + G_3^0$ . To generate this we take a line  $R$  and a twisted cubic meeting it in two points and place them in (1, 4) correspondence with one united point; but here the  $g_4^1$  on the cubic is such that each of its sets is made up of two pairs of three different involutions. Any one of the involutions gives  $\infty^1$  chords of the cubic forming a regulus, and there is thus a (1, 2) correspondence between the points of  $R$  and the lines of this regulus. We thus have, because of the united point, three developables of the third class, each of which has a plane passing through  $R$ .

**230.** Take now a curve  $C$  on  $\Omega$  of the type IV (C). The  $\rho$ -planes through  $O$  cut out a  $g_2^1$  on  $C$ , so that the ruled surface which is formed by those chords of  $C$  lying in the  $\rho$ -planes and not passing through  $O$  is rational. There are six tangents of  $C$ , other than that at  $O$ , which lie on  $\Omega^*$ ; of these two are in  $\rho$ -planes and four in  $\varpi$ -planes. Those chords of  $C$  which lie in  $\varpi$ -planes and do not pass through  $O$  form a ruled surface on which  $C$  is a double curve, and there are four points of  $C$  at which the two generators of this ruled surface coincide. The prime sections of this ruled surface are rational curves †.

A general plane  $\rho$  of  $\Omega$  meets the [4] containing  $C$  in a line; the  $\varpi$ -plane through this line joins it to  $O$  and contains six chords of  $C$ ; there are four other chords of  $C$  meeting the line. Hence the double curve of the ruled surface in [3] is  $6R + C_4^0$ .  $R$  is a trisecant of  $C_4^0$ ; two of the intersections are represented on  $\Omega$  by the  $\varpi$ -planes through the two trisecants of  $C$  which pass through  $O$ , the remaining one is represented on  $\Omega$  by the  $\varpi$ -plane through that chord of  $C$  which lies in the  $\rho$ -plane containing the tangent of  $C$  at  $O$ .

A general plane  $\varpi$  of  $\Omega$  meets the [4] containing  $C$  in a line; the  $\rho$ -plane through this line joins it to  $O$  and contains three chords of  $C$ ; there are seven other chords of  $C$  meeting the line. Hence the bitangent developable of the surface in [3] is  $3R + E_7^0$ . There are four planes of  $E_7^0$  passing through  $R$ ; two of these are represented on  $\Omega$  by the  $\rho$ -planes through the two trisecants of  $C$  which pass through  $O$ , the other two are represented by the two  $\rho$ -planes through those chords of  $C$  which join  $O$  to the other intersections with  $C$  of the  $\varpi$ -plane through the tangent of  $C$  at  $O$ . The two trisecants of  $C$  which do not pass through  $O$  are joined to  $O$  by  $\varpi$ -planes; the  $\rho$ -planes through these trisecants represent two triple planes of  $E_7^0$ .

\* For there are six tangents of a rational *quintic* in [3] which lie on a quadric containing the curve.

† Such a prime section is a curve of order 10 lying on a quadric in [3]; it meets all generators of one system in seven points and all of the other in three points, having two triple points and six double points.

231. To obtain this surface by projection we take the normal surface in [7] with  $\infty^1$  directrix cubics and choose  $S$  to meet the [4] determined by a directrix cubic  $\Delta$  and a generator  $g$  in a plane. Then this [4] and  $S$  determine a [5] through  $S$  which meets  $\Sigma$  in a line  $R$ ,  $R$  being a directrix and also a generator of the projected surface  $f$ . Through any point of  $R$  there pass three other generators, while any plane through  $R$  contains two other generators.

$S$  meets the solid containing  $\Delta$  in a line  $l$ . The [4] containing  $g$  and  $\Delta$  meets  $S$  in a plane through  $l$ , and  $\Delta$  is projected from  $g$  on to this plane into a conic  $\vartheta_2$  meeting  $l$  twice; through any point of  $\vartheta_2$  there passes a line meeting  $\Delta$  and  $g$ . Then  $S$  meets  $M_5^6$  in  $l$ ,  $\vartheta_2$  and a twisted cubic  $\vartheta_3^0$  which meets  $\vartheta_2$  once and  $l$  twice. Through any point of  $g$  there pass three lines meeting  $\vartheta_2$  and  $\Delta$ ; through any point of  $\Delta$  there pass two of its chords meeting  $l$  and one line meeting  $g$  and  $\vartheta_2$ . Thus the chords of  $F$  meeting  $S$  meet  $F$  in  $\Delta$  and  $g$  each counted three times and in a rational curve  $C_8^0$  meeting every generator once.  $C_8^0$  meets  $\Delta$  in five points; four of these are accounted for by the chords of  $F$  through the intersections of  $l$  and  $\vartheta_3^0$ ; the other lies, with the intersection of  $C_8^0$  with  $g$ , on the chord through the intersection of  $\vartheta_2$  and  $\vartheta_3^0$ .

Hence on projection we have in  $\Sigma$  a surface with a double curve  $6R + C_4^0$ ;  $C_4^0$  having  $R$  for a trisecant.

To generate this surface we take a line  $R$  and a twisted cubic with one intersection, placing them in (1, 3) correspondence without a united point. Each point of  $R$  gives a triad of points on the cubic, the plane of the triad meeting  $R$  in another point; we have thus a (1, 1) correspondence between the points of  $R$ , there being one triad having its plane passing through any point of  $R^*$ ; there are two coincidences, or two triads whose planes pass through the corresponding points on  $R$ . These give the two triple planes of  $E_7^0$ . There are four planes of  $E_7^0$  passing through  $R$ ; two of these arise from the triad of points containing the intersection of  $R$  with the cubic. The pencil of planes through  $R$  gives a  $g_2^1$  on the cubic, and this has two sets belonging to the  $g_3^1$  given by the triads†; these give the other two planes of  $E_7^0$  through  $R$ .

The tritangent planes of  $f$  are at once obtained from the projection;  $S$  meets the  $M_4^4$  formed by the solids  $K$  in  $l$  and two other points‡, the solids  $K$  through these other points project from  $S$  into the two tritangent planes of  $f$ .

The primes through the [5] containing  $S$  and  $\Delta$ —since they all contain the generator  $g$ —give pairs of generators of  $F$  which give a  $g_2^1$  on  $\Delta$ ; the joins of the pairs of this  $g_2^1$  are lines of a regulus, and there are two of

\* There is one of these planes passing through any point of  $\Sigma$ ; the equation of the plane is expressible linearly in terms of a parameter.

† Cf. the footnote to § 134.

‡ Cf. § 222.

these joins meeting  $l$ . Thus on projection there are two points of  $R$  such that, at either of these points, the plane of the pair of intersecting generators passes through  $R$ . This is of course clear from the representation on  $\Omega$ ; the points are represented by the  $\pi$ -planes containing the two trisecants of  $C$  which pass through  $O$ . If, however, it happened that one of these pairs of points on  $\Delta$  gave a pair of generators whose solid met  $S$  in a line, the projected surface would have a double generator and  $C$  would be of the type IV (E); similarly we might have two double generators,  $C$  being of the type IV (G).

**232.** The dual of the surface which we have been considering arises when  $C$  is of the type IV (D); the double curve is  $3R + C_7^0$ ,  $C_7^0$  having two triple points and having  $R$  as a quadrisecant; the bitangent developable is  $6R + E_4^0$ , there being three planes of  $E_4^0$  passing through  $R$ .

To obtain this surface by projection take the normal surface  $F$  in [7] with a directrix conic  $\Gamma$ , and take  $S$  to lie in a [5] containing  $\Gamma$  and a generator  $g$ ,  $S$  meeting the plane of  $\Gamma$  in a point  $O$ . Then the [5] meets  $\Sigma$  in a line  $R$  which is a directrix and also a generator of  $f$ ; through any point of  $R$  there pass two other generators, while any plane through  $R$  contains three other generators.

The [3] containing  $g$  and  $\Gamma$  meets  $S$  in a line  $l$  passing through  $O$ ; through each point of  $l$  there passes a line meeting  $g$  and  $\Gamma$ .  $S$  meets  $M_5^6$  in  $l$  and a rational quintic  $\mathfrak{D}_5^0$  which has a double point at  $O$  and meets  $l$  in two other points.

Through any point of  $g$  there pass two lines meeting both  $l$  and  $\Gamma$ , while through any point of  $\Gamma$  we have one line meeting  $l$  and  $g$  and also the line joining the point to  $O$ . Thus the chords of  $F$  which meet  $S$  meet  $F$  in  $\Gamma$  and  $g$ , each counted twice, together with an elliptic curve  $C_{14}^1$ , four\* of whose tangents meet  $\mathfrak{D}_5^0$ . This curve meets every generator of  $F$  in two points and  $\Gamma$  in six points; of these intersections with  $\Gamma$  four are collinear with  $O$  in pairs, the other two being associated with the intersections of  $g$  with  $C_{14}^1$ .  $C_{14}^1$  has six double points; these lie three in each of two planes meeting  $S$  in lines.

Hence the projected surface has a double curve  $3R + C_7^0$ ;  $C_7^0$  has two triple points and has  $R$  as a quadrisecant.

The surface is generated by a line  $R$  and a rational twisted quartic with one intersection, these being placed in (1, 2) correspondence without a united point. The pairs of points on the quartic corresponding to the points of  $R$  give a cubic ruled surface; thus the bitangent planes of the ruled surface other than those through  $R$  form a rational developable of the fourth class, which itself has three planes passing through  $R$ .

\* The tangent solid of  $F$  along  $g$  meets  $l$ .

**233.** Suppose now that  $C$  is of the type IV (E); the double point  $P$  represents a double generator  $G$  of the ruled surface in [3]. The chords of  $C$  lying in either system of planes and not passing through  $O$  form a rational ruled surface\*.

An arbitrary plane  $\rho$  of  $\Omega$  meets the [4] containing  $C$  in a line and the plane  $\omega$  through this line joins it to  $O$ ; this plane contains six chords of  $C$ , and there are three others meeting the line. Hence the surface has a double curve  $6R + G + C_3^0$ .  $R$  and  $G$  intersect, while  $R$  is a chord of  $C_3^0$ .  $C_3^0$  also meets  $G$ .

An arbitrary plane  $\omega$  of  $\Omega$  meets the [4] containing  $C$  in a line and the plane  $\rho$  through this line joins it to  $O$ ; this plane contains three chords of  $C$ , and there are six others meeting the line. Hence the surface has a bitangent developable  $3R + G + E_6^0$ ; there are three planes of  $E_6^0$  passing through  $R$ , while  $E_6^0$  has a double plane and an ordinary plane passing through  $G$ . The trisecant of  $C$  which does not pass through  $O$  is joined to  $O$  by a  $\omega$ -plane; the  $\rho$ -plane containing this trisecant represents a triple plane of  $E_6^0$ .

**234.** To obtain this surface by projection take a directrix cubic  $\Delta$  on the normal surface and a line  $l$  in the solid containing  $\Delta$ . Then take a pair of generators  $h, h'$  of  $F$  which meet  $\Delta$  at its intersections with one of its chords which meet  $l$ , and take a line  $m$  meeting  $l$  and lying in the solid containing  $h$  and  $h'$ . Then choose  $S$  to pass through  $l$  and  $m$  and lie in a [5] containing  $\Delta$  and a generator  $g$ . We have as before a conic  $\vartheta_2$  whose plane passes through  $l$ , and  $S$  meets  $M_5^6$  in  $l, m, \vartheta_2$  and a conic  $\phi_2$  meeting each of  $l, m, \vartheta_2$  in one point. The chords of  $F$  meeting  $S$  meet  $F$  in  $\Delta$  and  $g$  both counted three times,  $h, h'$  and a prime section  $C_6^0$ . On projection we have a double curve  $6R + G + C_3^0$  as required.

The projected surface has a tritangent plane,  $S$  meeting  $M_4^4$  in one point not on  $l$  or  $m$ .

This surface is generated by a line  $R$  and a twisted cubic  $C_3^0$  in (2, 3) correspondence;  $R$  is a chord of  $C_3^0$ , there being one doubly united point and one ordinary united point. The correspondence is such that to one of the branch-points on  $R$  there corresponds one of the branch-points on  $C_3^0$ ; the line joining these points is the double generator  $G$ .

\* The chords of  $C$  which lie in the  $\rho$ -planes and do not pass through  $O$  form a rational quartic ruled surface; they join the pairs of a  $g_2^1$  on  $C$  which includes the pair of points on the two branches at  $P$ .

The chords of  $C$  which lie in the  $\omega$ -planes and do not pass through  $O$  form a ruled surface of order 9; a prime section of this is a curve of order 9 lying on a quadric, meeting all generators of one system in three points and all of the other system in six points. It has one triple point and seven double points.



**235.** When  $C$  is of the type IV (F) we have a double curve  $3R + G + C_6^0$  and a bitangent developable  $6R + G + E_3^0$ .

Consider now the normal surface with a directrix conic  $\Gamma$ ; take a point  $O$  in the plane of  $\Gamma$  and a line  $m$  through  $O$  lying in the solid containing a pair of generators  $h, h'$ . Choose  $S$  to pass through  $m$  and also to lie in a [5] with  $\Gamma$  and a generator  $g$ . We have as before a line  $l$  in  $S$  passing through  $O$ ;  $S$  meets  $M_5^6$  in  $l, m$  and a rational quartic  $\mathfrak{D}_4^0$  passing through  $O$ , meeting  $m$  once again and  $l$  twice again.

The chords of  $F$  which meet  $S$  meet  $F$  in  $\Gamma$  and  $g$  both counted twice,  $h, h'$  and an elliptic curve  $C_{12}^1$  meeting every generator in two points and having four double points. Three of these lie in a plane containing a trisecant of  $\mathfrak{D}_4^0$ , the other being associated with intersections of  $C_{12}^1$  with  $h$  and  $h'$ ; the remaining intersections of  $C_{12}^1$  with  $h$  and  $h'$  lie on the chord of  $F$  passing through the intersection of  $m$  and  $\mathfrak{D}_4^0$  other than  $O$ . Two of the four intersections of  $C_{12}^1$  with  $\Gamma$  are collinear with  $O$ ; the other two are associated with the intersections of  $C_{12}^1$  with  $g$ , lying with them on the two chords of  $F$  which pass through the two intersections (other than  $O$ ) of  $l$  and  $\mathfrak{D}_4^0$ .

Thus the projected surface has a double curve  $3R + G + C_6^0$ ;  $R$  is a trisecant of  $C_6^0$  while  $C_6^0$  meets  $G$  and has also a double point on  $G$ .  $R$  and  $G$  intersect and  $C_6^0$  has a triple point.

This surface is generated by a line  $R$  and a rational skew quartic in (1, 2) correspondence; there is one intersection but no united point. The correspondence is specialised to give the double generator; one of the points of  $R$  gives rise to a pair of points of the quartic collinear with it. The joins of the pairs of points of the quartic form a cubic ruled surface and the planes joining the points of  $R$  to the corresponding pairs form a developable  $E_3^0$  with two planes passing through  $R$  and one through  $G$ . The bitangent developable is  $6R + G + E_3^0$ .

**236.** When  $C$  is of the type IV (G) the double curve of the ruled surface is  $6R + G + H + C_2$  and the bitangent developable  $3R + G + H + E_5^0$ .  $G$  and  $H$  meet  $R$ , while  $C_2$  meets each of  $R, G$  and  $H$ . There are two planes of  $E_5^0$  passing through  $R$ , while through  $G$  there pass a double plane and a simple plane of  $E_5^0$ , as also through  $H$ .

To obtain the surface by projection we choose the solid  $S$  to lie in a [5] containing a directrix cubic  $\Delta$  and a generator  $g$  of  $F$ , there being two pairs of generators  $g', g''$  and  $h', h''$  whose solids meet  $S$  in lines  $m$  and  $n$ .  $S$  meets the solid containing  $\Delta$  in a line  $l$  which meets  $m$  and  $n$ ; it meets the [4] containing  $g$  and  $\Delta$  in a plane through  $l$ , and  $\Delta$  is projected from  $g$  into a conic  $\mathfrak{D}_2$  in this plane. Then  $S$  meets  $M_5^6$  in  $l, m, n, \mathfrak{D}_2$  and a line  $\lambda$  which meets  $m, n$  and  $\mathfrak{D}_2$ . The chords of  $F$  meeting  $S$  meet  $F$  in  $\Gamma$  and  $g$

both counted three times,  $g', g'', h', h''$  and a directrix quartic  $E$ ; the double curve of the projected surface is precisely as given above.

The surface is generated by a line  $R$  and a twisted cubic which meets it, these being in (1, 3) correspondence without a united point; the correspondence is specialised to give the two double generators. Or it can be generated by  $R$  and  $C_2$  in (2, 3) correspondence with a united point, there being two pairs of corresponding double elements.

**237.** When  $C$  is of the type IV (H) the double curve of the ruled surface is  $3R + G + H + C_5^0$  and the bitangent developable  $6R + G + H + E_2$ .  $R$  meets both  $G$  and  $H$ ;  $C_5^0$  has two double points, one of which lies on  $G$  and the other on  $H$ , each of these lines meeting  $C_5^0$  in a further point.  $R$  is a chord of  $C_5^0$ . There is one plane of  $E_2$  passing through each of  $R, G, H$ .

To obtain this surface by projection take in [7] the normal surface  $F$  with a directrix conic  $\Gamma$ . Take a point  $O$  in the plane of  $\Gamma$  and two lines  $m$  and  $n$  passing through  $O$ ,  $m$  lying in the solid containing a pair of generators  $g', g''$  and  $n$  lying in the solid containing a pair of generators  $h', h''$ . Then choose  $S$  to contain  $m$  and  $n$  and to lie in a [5] with  $\Gamma$  and a generator  $g$ . The solid  $g\Gamma$  meets  $S$  in a line  $l$  through  $O$ ; and  $S$  meets  $M_5^6$  in  $l, m, n$  and a twisted cubic  $\mathfrak{D}_3^0$  meeting  $m$  and  $n$  and having  $l$  as a chord.

The chords of  $F$  meeting  $S$  meet  $F$  in  $\Gamma$  and  $g$  each counted twice,  $g', g'', h', h''$  and an elliptic curve  $C_{10}^1$  meeting every generator of  $F$  twice and having two double points. These double points lie one in each of the solids  $g'g''$  and  $h'h''$  and are thus associated with intersections of  $C_{10}^1$  with these generators; the remaining intersections of  $C_{10}^1$  with these generators are given by the chords of  $F$  through the points in which  $\mathfrak{D}_3^0$  meets  $m$  and  $n$ .  $C_{10}^1$  meets  $\Gamma$  in two points; these are associated with its intersections with  $g$  and lie on the chords of  $F$  through the intersections of  $l$  and  $\mathfrak{D}_3^0$ . On projection we have a double curve as above described.

The surface is generated by a line  $R$  and a rational skew quartic in (1, 2) correspondence;  $R$  meets the quartic, but there is no united point. The correspondence is specialised to give the double generators; there are two points on  $R$  which give pairs of points of the quartic collinear with them. The joins of the pairs of points of the quartic form a cubic ruled surface, and the planes joining the points of  $R$  to the pairs of points of the quartic which correspond to them touch a quadric cone. There is a tangent plane of this cone passing through  $R$  and one through each double generator.

**238.** We now pass to curves  $C$  on  $\Omega$  of the type V. When  $C$  is of the type V (A) we have a ruled surface in [3] with a directrix line  $R$  which is also a double generator; through any point of  $R$  there pass three other

generators, while any plane through  $R$  contains one other generator. It is clear how to obtain such a surface by projection; we take  $S$  to meet the solid  $K$  containing a directrix cubic  $\Delta$  of the normal surface  $F$  in a line  $l$ , and so that  $S$  lies in the [5] containing  $\Delta$  and two generators  $g$  and  $h$ . Then this [5] meets  $\Sigma$  in the line  $R$ .

The surface is generated by a twisted cubic and one of its chords in (3, 1) correspondence without any united point.

An arbitrary plane  $\rho$  of  $\Omega$  meets the [4] containing  $C$  in a line; this line is joined to  $O$  by a  $\omega$ -plane which contains all the chords of  $C$  meeting the line; the double curve of the surface is thus the line  $R$  counted ten times.

An arbitrary plane  $\omega$  of  $\Omega$  meets the [4] containing  $C$  in a line, and the plane  $\rho$  through this line joins it to  $O$ . If  $C$  is projected on to a plane from this line we obtain a rational sextic with a triple point and therefore with seven other double points. Hence there are seven chords of  $C$  which meet the line and do not lie in the plane joining the line to  $O$ . Hence the ruled surface has a bitangent developable  $3R + E_7$ . The chords of  $C$  which lie in  $\omega$ -planes and do not pass through  $O$  form a ruled surface whose prime sections are rational curves\*, so that  $E_7$  is a rational developable  $E_7^0$ . There are four planes of  $E_7^0$  passing through  $R$ ; the  $\omega$ -plane through either tangent of  $C$  at  $O$  meets  $C$  in two points other than  $O$ , and the  $\rho$ -planes containing the chords joining  $O$  to these points represent two planes of  $E_7^0$  which pass through  $R$ .

$C$  has two trisecants; these are joined to  $O$  by  $\omega$ -planes, and the  $\rho$ -planes through them represent two triple planes of  $E_7^0$ .

**239.** The dual surface, represented on  $\Omega$  by a curve  $C$  of the type V (B), has a double curve  $3R + C_7^0$  and a bitangent developable  $10R$ .  $R$  is a quadrisecant of  $C_7^0$ , which has two triple points.

This surface is obtained by projection from the normal surface with a directrix line  $\lambda$ ;  $S$  must be chosen to meet the plane containing  $\lambda$  and a generator  $g$  in a point  $X$  and to meet the plane containing  $\lambda$  and another generator  $h$  in a point  $Y$ . Then the [5] containing  $S$  and  $\lambda$  also contains  $g$  and  $h$ , and meets  $\Sigma$  in a line  $R$  which is a directrix and also a double generator of  $f$ . Through any point of  $R$  there passes one other generator, while any plane through  $R$  contains three other generators.

The line  $XY$  is on the  $M_5^6$  formed by the chords of  $F$ ; through every point of  $XY$  there passes a transversal to  $g$  and  $h$ .  $S$  meets  $M_5^6$  in  $XY$  and a rational quintic  $\mathfrak{D}_5^0$  with double points at  $X$  and  $Y$ .

\* There are four points of  $C$  at which the two generators of this ruled surface coincide and also four tangents of  $C$ , other than that at  $O$ , lying in  $\omega$ -planes (this being the number of tangents of a rational quartic which lie on a quadric containing it). A prime section of the ruled surface is a curve of order ten lying on a quadric surface, meeting all generators of one system in three points and all of the other system in seven points; it has six double points and two triple points.

Through any point of  $\lambda$  there pass lines through  $X$  and  $Y$  meeting  $g$  and  $h$  respectively; through any point of  $g$  there passes a transversal to  $h$  and  $XY$  and also a line through  $X$  meeting  $\lambda$ ; through any point of  $h$  there passes a transversal to  $g$  and  $XY$  and also a line through  $Y$  meeting  $\lambda$ ; thus the chords of  $F$  which meet  $S$  meet  $F$  in the lines  $\lambda, g, h$  each counted twice and an elliptic curve  $C_{14}^1$  (four of whose tangents meet  $\mathfrak{S}_5^0$ ) meeting every generator of  $F$  in two points and  $\lambda$  in four points. This curve has six double points, and these lie three in each of two planes containing trisecants of  $\mathfrak{S}_5^0$ . Hence the double curve of  $f$  is  $3R + C_7^0$ ;  $R$  is a quadrisecant of  $C_7^0$ , which has two triple points.

The surface is generated by a rational quintic and one of its chords in 1, 1) correspondence without a united point.

**240.** Suppose now that  $C$  is of the type V (C); we have a line  $R$  which is a directrix and also a double generator of the ruled surface; through any point of  $R$  there pass two other generators, while any plane through  $R$  contains two other generators. The chords of  $C$  which lie in either system of planes and do not pass through  $O$  join pairs of points of a  $g_2^1$  and form a rational quintic ruled surface.

An arbitrary plane of  $\Omega$  meets the [4] containing  $C$  in a line, and the plane of the opposite system through this line joins it to  $O$ ; there are four chords of  $C$  which meet the line and do not lie in this plane. Hence the double curve of the ruled surface is  $6R + C_4^0$  and the bitangent developable  $6R + E_4^0$ .

$R$  is a trisecant of  $C_4^0$ ; the  $\omega$ -plane containing the line through  $O$  which meets  $C$  in two further points represents one of the three intersections. The  $\rho$ -plane through either tangent of  $C$  at  $O$  meets  $C$  in another point, and the  $\omega$ -plane containing the line joining this point to  $O$  represents one of the other intersections.

Similarly there are three planes of  $E_4^0$  passing through  $R$ .

**241.** To obtain this surface by projection we take the normal surface which has a directrix conic  $\Gamma$ , and choose  $S$  to lie in a [5] containing  $\Gamma$  and two generators  $g$  and  $h$ . Then  $S$  meets the plane of  $\Gamma$  in a point  $O$ , and the [5] meets  $\Sigma$  in the line  $R$ . The [4] containing  $\Gamma, g$  and  $h$  meets  $S$  in a plane passing through  $O$  and the solids  $gh, g\Gamma, h\Gamma$  meet  $S$  in three lines  $l, m, n$  which lie in this plane,  $m$  and  $n$  passing through  $O$ .  $S$  meets  $M_5^0$  in  $l, m, n$  and a twisted cubic which passes through  $O$  and meets  $m$  and  $n$  again.

Through any point of  $\Gamma$  there pass a transversal to  $g$  and  $m$ , a transversal to  $h$  and  $n$  and a line to  $O$ ; through any point of  $g$  there pass two transversals to  $\Gamma$  and  $m$  and a transversal to  $h$  and  $l$ ; through any point of  $h$  there pass two transversals to  $\Gamma$  and  $n$  and one transversal to  $g$  and  $l$ . Hence the chords of  $F$  which meet  $S$  meet  $F$  in  $\Gamma, g$  and  $h$  each counted

three times and a rational curve  $C_8^0$  meeting each generator once and  $\Gamma$  four times. Two of these intersections with  $\Gamma$  are collinear with  $O$ ; the other two lie on the chords of  $F$  through the other intersections of  $\mathfrak{D}_3^0$  with  $m$  and  $n$  and are associated with the intersections of  $C_8^0$  with  $g$  and  $h$ .

Hence on projection we have a double curve  $6R + C_4^0$ ,  $R$  being a tri-secant of  $C_4^0$ .

The surface is generated by a rational skew quartic and one of its chords placed in (2, 1) correspondence without any united point. The joins of the pairs of points of the quartic form a cubic ruled surface; so that the planes joining the points of the chord to the pairs of points of the quartic which correspond to them form a developable  $E_4^0$  with three planes passing through  $R$ .

**242.** The pencil of primes through the [5] containing  $S$ ,  $\Gamma$ ,  $g$  and  $h$  cuts out an involution of pairs of generators on  $F$ . There is one pair of this involution meeting  $\Gamma$  in a pair of points collinear with  $O$ ; these are in fact two intersections of  $C_8^0$  with  $\Gamma$ .

Now it may happen that the solid containing this pair of generators meets  $S$  in a line  $p$  through  $O$ ; in this case the projected surface has a double generator and corresponds to a curve  $C$  of the type V (D).  $S$  meets  $M_5^0$  in  $l$ ,  $m$ ,  $n$ ,  $p$  and a conic  $\mathfrak{D}_2$  meeting  $m$ ,  $n$  and  $p$ . On  $F$  we have  $\Gamma$ ,  $g$ ,  $h$  each counted three times and the new pair of generators  $k$ ,  $k'$  together with a prime section  $C_8^0$ .

The surface is generated by a rational skew quartic and one of its chords  $R$  placed in (2, 1) correspondence without any united point; but here the correspondence must be specialised to give the double generator  $G$ , one pair of points of the quartic having their join  $G$  meeting  $R$  in the point which gives rise to them. The joins of the pairs of points of the quartic form a cubic ruled surface, and we thus obtain a developable  $E_3^0$  with two planes passing through  $R$  and one through  $G$ . The bitangent developable of the surface is  $6R + G + E_3^0$ .

The properties of the surface are also easily deduced from those of the curve on  $\Omega$ . The double curve is  $6R + G + C_3^0$ .

**243.** We now suppose that  $C$  is of the type VI (A); the chords of  $C$  in the  $\omega$ -planes which do not pass through  $O$  form a rational quintic ruled surface. An arbitrary plane  $\rho$  meets the [4] containing  $C$  in a line and the plane  $\omega$  containing this line joins it to  $O$ ; there are no chords of  $C$  meeting the line other than those which lie in this  $\omega$ -plane. Thus the double curve of the ruled surface is  $10R$ . An arbitrary plane  $\omega$  of  $\Omega$  meets the [4] containing  $C$  in a line; there are four chords of  $C$  meeting this line which do not lie in the  $\rho$ -plane joining it to  $O$ . Hence the bitangent developable of the surface is  $6R + E_4^0$ .

The  $\omega$ -plane through any one of the three tangents of  $C$  at  $O$  meets  $C$  again in another point; the  $\rho$ -plane containing the line joining this point to  $O$  represents a plane of  $E_4^0$  which passes through  $R$ . Thus there are three planes of  $E_4^0$  passing through  $R$ .

To obtain this surface by projection we take the normal surface with a directrix conic  $\Gamma$  and choose  $S$  to lie in the [5] determined by  $\Gamma$  and three generators  $g, h, k$ . Then this [5] meets  $\Sigma$  in a line  $R$  which is a directrix and also a triple generator of  $F$ ; through any point of  $R$  there pass two other generators, while any plane through  $R$  contains one other generator.  $S$  meets  $M_5^6$  in three lines  $l, m, n$  passing through the point  $O$  in which it meets the plane of  $\Gamma$  and in three other lines  $\lambda, \mu, \nu$ , where  $\lambda$  meets  $m$  and  $n$ ,  $\mu$  meets  $n$  and  $l$ , and  $\nu$  meets  $l$  and  $m$ . The double curve of the projected surface is  $10R$ ; the chords of  $F$  meeting  $S$  meet  $F$  in  $\Gamma, g, h$  and  $k$ , each counted four times.

The surface is generated by a rational skew quartic and a line  $R$  in (2, 1) correspondence;  $R$  is a trisecant of the quartic but there are no united points. The joins of the pairs of points of the quartic form a cubic ruled surface, and the planes joining the points of  $R$  to the pairs of points of the quartic which correspond to them give a developable  $E_4^0$ , three of whose planes pass through  $R$ .

**244.** When  $C$  is of the type VI (B) we have the surface dual to this; the double curve is  $6R + C_4^0$ , where  $R$  is a trisecant of  $C_4^0$ , and the bitangent developable is  $10R$ .

To obtain this surface by projection take the normal surface with a directrix line  $\lambda$ ; choose  $S$  to meet the plane containing  $\lambda$  and a generator  $g$  in a point  $X$ , the plane containing  $\lambda$  and a generator  $h$  in a point  $Y$  and the plane containing  $\lambda$  and a generator  $k$  in a point  $Z$ . Then there is a [5] containing  $S, \lambda, g, h, k$  which meets  $\Sigma$  in a line  $R$  which is a directrix and also a triple generator of the surface. Through any point of  $R$  there passes one other generator, while any plane through  $R$  contains two other generators.

$S$  meets  $M_5^6$  in the lines  $YZ, ZX, XY$  and a rational cubic  $\mathfrak{D}_3^0$  passing through  $X, Y$  and  $Z$ . The chords of  $F$  which meet  $S$  meet  $F$  in  $\lambda, g, h, k$ , each counted three times and in a rational curve  $C_8^0$  meeting each generator once and  $\lambda$  three times, there being two tangents of  $C_8^0$  meeting  $\mathfrak{D}_3^0$ . On projection we have a double curve  $6R + C_4^0$ , where  $R$  is a trisecant of  $C_4^0$ .

The surface is generated by a rational skew quintic and one of its trisecants in (1, 1) correspondence without any united point.

### *Surfaces whose generators belong to a linear congruence*

**245.** Suppose now that  $C$  is of the type VII (A); through the solid containing the quadric surface on which  $C$  lies there pass two tangent

primes of  $\Omega$ , touching it in  $O$  and  $O'$ ; these represent two lines  $R$  and  $R'$  in [3] which are directrices of the ruled surface. Through any point of one of these (say  $R$ ) there pass five generators lying in a plane through  $R'$ . Hence the double curve of the ruled surface is  $10R$  and the bitangent developable is  $10R'$ .

To obtain this surface by projection take the normal surface  $F$  with a directrix line  $\lambda$  and project from a solid  $S$  lying in a [5] which contains one of its directrix quintics. Then this [5] meets  $\Sigma$  in a line  $R$ , while the [5] through  $S$  and  $\lambda$  meets  $\Sigma$  in a line  $R'$ ;  $R$  and  $R'$  are directrices of  $f$ ; through any point of  $R$  there pass five generators lying in a plane through  $R'$ .

The surface is generated by two lines  $R$  and  $R'$  in (1, 5) correspondence.

The solid  $S$  meets  $M_6^6$  in the curve  $\mathfrak{D}_6^3$  in which it is met by the chords of the directrix quintic; the chords of  $F$  which meet  $S$  meet  $F$  in the directrix quintic counted four times.

**246.** When  $C$  is of the type VII (B) the ruled surface has two directrices  $R$  and  $R'$  and three double generators  $G, H, K$ . Through any point of  $R$  there pass four generators lying in a plane through  $R'$ , while through any point of  $R'$  there pass two generators lying in a plane through  $R$ . The double curve is  $6R + G + H + K + R'$  and bitangent developable is  $R + G + H + K + 6R'$ .

To obtain this surface by projection we take the normal surface with a directrix conic  $\Gamma$ ; take  $S$  to meet the plane of  $\Gamma$  in a point  $O$  and to contain a plane  $\pi$  which lies in a [4] containing a directrix quartic  $E$ . Then the three double generators arise automatically, there being three pairs of generators whose solids meet  $S$  in lines  $l, m, n$  passing through  $O$  and meeting  $\pi^*$ .  $S$  meets  $M_5^6$  in  $l, m, n$  and a plane cubic  $\mathfrak{D}_3^1$  lying in  $\pi$  and passing through the intersections of  $\pi$  with  $l, m, n$ .

The surface is generated by a (2, 4) correspondence between  $R$  and  $R'$  specialised to give the three double generators.

The chords of  $F$  meeting  $S$  meet  $F$  in  $E$  counted three times,  $g, g', h, h', k, k'$  and  $\Gamma$ .

**247.** In type VII (C) the ruled surface has two directrices  $R$  and  $R'$  and four double generators  $G, H, J, K$ . Through any point of each directrix there pass three generators lying in a plane through the other. The double curve and bitangent developable are both  $3R + G + H + J + K + 3R'$ .

\* Through  $\pi$  there pass solids cutting out a  $g_4^1$  on  $E$ , while through  $O$  there pass lines cutting out a  $g_2^1$  on  $\Gamma$ ; there are three pairs of generators such that a pair meets  $\Gamma$  in a set of the  $g_2^1$  and meets  $E$  in two points of a set of the  $g_4^1$ . These are the three pairs whose solids meet  $S$  in lines.

Take the general surface  $F$  in [7] and project from a solid  $S$  containing a line  $l$  in the solid  $K$  containing a directrix cubic  $\Delta$  and a line  $l'$  in the solid  $K'$  containing a directrix cubic  $\Delta'$ . Then there are four pairs of generators whose solids meet  $S$  in lines  $m, n, p, q$  each of which meets  $l$  and  $l'$ .  $S$  meets  $M_5^6$  in  $l, l', m, n, p, q$ .

The surface is generated by two lines  $R$  and  $R'$  in (3, 3) correspondence, the correspondence being specialised to give the four double generators.

**248.** When  $C$  is of the type VIII (A) the surface has a single directrix  $R$  which is also a quadruple generator; through any point of  $R$  there passes one other generator. To obtain this by projection we take the normal surface in [7] with a directrix line  $\lambda$  and project from a solid  $S$  meeting four planes which join  $\lambda$  to four different generators of the surface, giving the surface in  $\Sigma$  as required.

The surface is generated by a rational skew quintic and its quadric secant in (1, 1) correspondence without any united point.

The double curve and bitangent developable are both  $10R$ .

**249.** When  $C$  is of the type VIII (B) the surface has a directrix  $R$  which is also a double generator; through any point of  $R$  there pass two other generators which are coplanar with  $R$ .

Take in [7] the normal surface with a directrix conic  $\Gamma$ , and the [4] which contains  $\Gamma$  and two generators. Two primes through this [4] give each two further generators of  $F$ , and we thus have two chords of  $\Gamma$  which intersect in a point  $O$ . The primes meet in a [5] containing the plane of  $\Gamma$  and also the first two generators; choose  $S$  to lie in this [5] and to pass through  $O$ . Since there are two of the pencil of primes through the [5] which meet  $F$  in pairs of generators whose solid passes through  $O$  it follows that all the primes of the pencil must do so, so that we have an infinity of pairs of generators whose solids meet  $S$  in  $O$  and which meet  $\Gamma$  in pairs of points collinear with  $O$ . Hence on projection we have the surface required. The double curve and bitangent developable are both  $7R + G + H + K$ .

The surface is generated by a (2, 1) correspondence between a rational skew quartic and one of its chords  $R$ ; the pairs of points of the quartic lying in planes through  $R$ . The joins of these pairs of points form a cubic ruled surface of which  $R$  is the double line. The range of points on  $R$  is thus related to the pencil of planes through  $R$ . There is incidentally a (1, 2) correspondence set up between the points of  $R$ ; through any point of  $R$  there pass two chords of the quartic (other than  $R$ ), while any plane through  $R$  gives another chord of the quartic meeting it. There will be three coincidences in this correspondence, thus shewing the existence of the double generators  $G, H, K$ .



250. When  $C$  is of the type VIII (C) we have a surface with a directrix  $R$ ; through any point of  $R$  there pass three generators which lie in a plane through  $R$ . There are four double generators; the double curve and bi-tangent developable are  $6R + G + H + J + K$ .

Take a directrix cubic  $\Delta$  of the normal surface  $F$ , and two primes through it. Each meets  $F$  further in three generators, giving thus two triads of points on  $\Delta$  whose planes meet in a line  $l$ . Take  $S$  to pass through  $l$  and to lie in the [5] common to the two primes. Then the pencil of primes through the [5] is such that any one of the primes meets  $F$  in  $\Delta$  and three generators, these generators meeting  $\Delta$  in a triad of points whose plane passes through  $l$ . On projection we have the surface required.

The surface is generated by a line and a twisted cubic in (1, 3) correspondence; the triads of points on the cubic being those cut out by the planes through the line. We thus have a (1, 3) correspondence between the points of the line; there are four coincidences, thus shewing the existence of the four double generators.

251. On pp. 306–308 we give tables of the different types of rational ruled surfaces of the sixth order in [3] (not including developable surfaces). There are eighty-one different types altogether. We have already obtained sixty-seven of these; the remaining ones will be obtained in Chapter VI.

## SECTION II

### ELLIPTIC SEXTIC RULED SURFACES

252. We first obtain the different types of elliptic sextic ruled surfaces in [3] by means of elliptic sextic curves on  $\Omega$ ; afterwards we obtain them as the projections of normal surfaces in [5].

In the first investigation we shall need to use the properties of certain ruled surfaces formed by chords of an elliptic sextic curve. The elliptic sextic curve is normal in [5]\*, all elliptic sextic curves being obtainable as the projections of curves in [5]. On the normal curve there are  $\infty^1$  linear series of pairs of points, the chords joining the pairs of points of such a linear series form a rational quartic ruled surface with, in general,  $\infty^1$  directrix conics, each conic meeting the curve in two points †. To each of these ruled surfaces there belong four tangents of the curve.

\* § 8.

† Segre, "Sur les transformations des courbes elliptiques," *Math. Ann.* 27 (1886), 296. Of the  $\infty^1$  series  $g_2^1$  there are nine for which the quartic ruled surface has a directrix line.

*Elliptic sextic curves which lie on quadrics*

**253.** We can at once divide the elliptic sextic curves  $C$  which lie on a quadric  $\Omega$  in [5] into the following seven classes:

I. The normal curve in [5].

II. The curve  $C$  is contained in a [4].

III.  $C$  is contained in a tangent prime  $T$  of  $\Omega$  but does not pass through the point of contact  $O$  of  $\Omega$  and  $T$ .

IV.  $C$  is contained in  $T$  and passes through  $O$ .

V.  $C$  is contained in  $T$  and has a double point at  $O$ .

VI.  $C$  lies on the quadric in which  $\Omega$  is met by a [3].

VII.  $C$  lies on a quadric cone in which  $\Omega$  is met by a [3] which touches it.

When  $C$  is contained in a space [4] it may or may not have a double point; if it has not a double point it has two trisecant chords\*.

We have two kinds of elliptic sextic curves on a quadric in [3]:

(A) The intersection with a quartic surface passing through two generators of the same system and touching the quadric twice. This curve meets all generators of one system in four points and all of the other system in two points; it has two double points.

(B) The intersection with a cubic surface touching the quadric in three points; this curve meets each generator of the quadric in three points and has three double points.

Also we have on a quadric cone:

(A) The intersection with a cubic surface passing through the vertex and touching the cone twice; this curve has a double point at the vertex and also two other double points; it meets every generator of the cone in two points other than the vertex.

(B) The intersection with a cubic surface touching the cone three times; this curve has three double points and meets every generator in three points.

**254.** We must not overlook the fact that there can exist on a quadric an elliptic sextic curve with a triple point. It is true that such a curve does not lie on a quadric in general, but if the tangents at the triple point are co-planar it will do so, the quadric being then determined by having to touch at the triple point the plane of the three tangents and to pass through six arbitrary points of the curve. The curve meets every generator of the quadric in three points, and we can also have such a curve lying on a quadric cone.

The quintic ruled surface formed by the trisecants of the curve † breaks up into the quadric counted twice and the plane of the tangents at the triple point.

\* Berzolari, *Palermo Rend.* 9 (1895), 195; Castelnuovo, *ibid.* 3 (1889), 28.

† § 89.

To obtain such a curve in [3] we must project the normal curve in [5] from a line in a trisecant plane; the three tangents of the curve at the points where it is met by the trisecant plane lying in a [4]. The existence of such planes is at once clear on using elliptic arguments for points of the curve; there are  $\infty^2$  such, four passing through any chord of the curve. Thus the curve in [3] exists for all values of the modulus.

We also have a curve in [4] such that the tangents at the three points where it is met by a trisecant lie in a [3].

**255.** We are now in a position to give the following more detailed classification of the elliptic sextic curves  $C$  on a quadric  $\Omega$  in [5]. The abbreviations  $\varpi_2\rho_4$ ,  $O\varpi_3\rho_2$ , etc., are explained in the footnote to § 171.

- I.  $C$  is a normal curve on  $\Omega$ .
- II. (A)  $C$  lies in a [4] which does not touch  $\Omega$ , and has no double point.
- II. (B)  $C$  lies in a [4] which does not touch  $\Omega$ , and has a double point.
- III.  $C$  lies in a tangent prime  $T$  but does not pass through  $O$ , the point of contact of  $\Omega$  and  $T$ .
  - (A)  $C$  is  $\varpi_2\rho_4$  with two chords through  $O$ .
  - (B)  $C$  is  $\varpi_4\rho_2$  with two chords through  $O$ .
  - (C)  $C$  is  $\varpi_2\rho_4$  with a chord through  $O$  and with a double point.
  - (D)  $C$  is  $\varpi_4\rho_2$  with a chord through  $O$  and with a double point.
  - (E)  $C$  is  $\varpi_3\rho_3$  with three chords through  $O$ .
  - (F)  $C$  is  $\varpi_3\rho_3$  with two chords through  $O$  and with a double point.
- IV.  $C$  lies in a tangent prime  $T$  and passes through  $O$ .
  - (A)  $C$  is  $O\varpi_2\rho_3$  with a trisecant through  $O$ .
  - (B)  $C$  is  $O\varpi_3\rho_2$  with a trisecant through  $O$ .
  - (C)  $C$  is  $O\varpi_2\rho_3$  with a double point.
  - (D)  $C$  is  $O\varpi_3\rho_2$  with a double point.

These four types can be written down at once because when  $C$ , which lies on the quadric point-cone with vertex  $O$  in which  $T$  meets  $\Omega$ , is projected from  $O$  on to a solid in  $T$  it becomes an elliptic quintic curve lying on a quadric\*. Similarly the curves under III must give elliptic sextics when thus projected †.

V.  $C$  lies in a tangent prime  $T$  and has a double point at  $O$ . It will meet every plane of  $\Omega$  passing through  $O$  in two points other than  $O$ .

VI.  $C$  lies in a [3] through which two tangent primes of  $\Omega$  pass; it thus lies on an ordinary quadric surface.

(A)  $C$  meets all generators of one system in four points and all of the other system in two points, having two double points.

(B)  $C$  meets every generator in three points and has three double points.

(C)  $C$  meets every generator in three points and has a triple point.

\* Cf. § 151.

† Cf. § 253.

VII.  $C$  lies in a [3] which touches  $\Omega$ , and therefore on a quadric cone with vertex  $V$ .

(A)  $C$  has a double point at  $V$  and also two other double points, meeting every generator in two points other than  $V$ .

(B)  $C$  has three double points and meets every generator in three points.

(C)  $C$  has a triple point and meets every generator in three points.

*Surfaces whose generators do not belong to a linear complex*

256. If we have a normal elliptic sextic curve  $C$  in [5] the projection of  $C$  from a plane on to a plane is an elliptic sextic curve with nine double points, while the projection from a plane meeting  $C$  is an elliptic quintic curve with five double points. Hence the chords of  $C$  form a locus  $M_3^9$  of three dimensions and the ninth order, on which  $C$  is a quadruple curve.  $C$  has no quadrisecant planes, because an elliptic quartic in [3] has no double points, so that there is no double surface on  $M_3^9$ .

If there is a quadric  $\Omega$  passing through  $C$  every plane of  $\Omega$  is met by nine chords of  $C$ , so that in the most general type of elliptic sextic ruled surface in [3] the double curve  $C_9$  is of order nine and the bitangent developable  $E_9$  of class  $9^*$ .

There are four trisecant planes of  $C$  which lie on  $\Omega$  †; there are, in fact, two of these belonging to each system of planes on  $\Omega$ . Hence  $C_9$  has two triple points and  $E_9$  two triple planes.

The chords of  $C$  which lie on  $\Omega$  form a ruled surface  $R_2^{18}$  of order 18, the intersection of  $\Omega$  and  $M_3^9$ ; there are four such chords passing through any point of  $C$ . A prime section  $C'$  of  $R_2^{18}$  is in (1, 1) correspondence with  $C_9$  and  $E_9$  ‡.

In the  $(\alpha, \alpha')$  correspondence between  $C$  and  $C'$  determined by the condition that two points correspond when the line joining them is a chord of  $C$  we have  $\alpha = 2$  and  $\alpha' = 4$ . The number  $\eta'$  of branch points of this correspondence on  $C'$  is the number of tangents of  $C$  which lie on  $\Omega$ , so that  $\eta' = 12$  ||. Through each point of  $C$  there pass four of its chords which lie on  $\Omega$ ; the number  $\eta$  of branch points of the correspondence on  $C$  is the number of points of  $C$  for which two of these four generators of  $R_2^{18}$  coincide. But if, through a point  $P$  of  $C$ , we have the four chords  $PA, PB, PC, PD$  lying on  $\Omega$ , the points  $A, B, C, D$  are a set of points on an elliptic curve of a (4, 4) correspondence of valency 2. Hence  $\eta = 24$  ¶.

Zeuthen's formula then shews that  $p' = 4^{**}$ .

Hence the most general elliptic sextic ruled surface in [3] has a double curve  $C_9^4$  and a bitangent developable  $E_9^4$ .

\* § 33.  
|| § 32.

† § 35.  
¶ § 14.

‡ § 33.  
\*\* § 16.

257. Consider now a quartic ruled surface, with  $\infty^1$  directrix conics, containing the curve; each generator of this surface is a chord of the curve, while each directrix conic meets it in two points.

There are  $\infty^{20}$  quadrics in [5], and of these there are  $\infty^8$  containing the curve  $C$  of order six and genus 1; a quadric meeting  $C$  in twelve general points necessarily containing it entirely\*. If, in addition, the quadric is made to contain three generators of the quartic surface it will contain the whole of the surface, so that we have  $\infty^5$  such quadrics†. This quadric contains the planes of two directrix conics of the surface‡; the  $\infty^1$  planes form a  $V_3^3$  which is met by the quadric in two of the planes and in the quartic surface. The planes are of opposite systems on the quadric.

The ruled surface formed by the chords of  $C$  which lie on  $\Omega$  now breaks up into this quartic surface and a ruled surface of the fourteenth order on which  $C$  is a triple curve. Every plane of  $\Omega$  meets the quartic surface in two points and therefore the other surface in seven points. We thus obtain in [3] an elliptic sextic ruled surface whose double curve consists of a conic and a curve of the seventh order and whose bitangent developable consists of a quadric cone and a developable of the seventh class.

Through any point  $P$  of  $C$  there pass four of its chords which lie entirely on  $\Omega$ ; one of these  $PQ$  is a generator of the quartic ruled surface, while the other three  $PX, PY, PZ$  are generators of the ruled surface of order 14. Since the set of points  $P + Q$  and also the set  $2P + Q + X + Y + Z$  vary in two linear series for different positions of  $P$  on  $C$  so does the set  $P + X + Y + Z$ . The generators of the ruled surface of order 14 thus set up on  $C$  a (3, 3) correspondence of valency 1, so that there are sixteen points of  $C$  at which two of the three generators of this ruled surface coincide.

Again, of the twelve tangents of  $C$  which lie on  $\Omega$  there are four belonging to the quartic; there must then be eight belonging to the ruled surface of order 14.

Considering then the correspondence between  $C$  and the curve  $C'$  in which the ruled surface of order 14 is met by a [4] we have

$$\alpha = 2, \quad \alpha' = 3, \quad p = 1, \quad \eta = 16, \quad \eta' = 8,$$

whence  $p' = 3$ , so that  $C'$  is of genus 3.

Thus we have in ordinary space a ruled surface whose double curve is  $C_2 + C_7^3$  and bitangent developable  $E_2 + E_7^3$ .

The generators of the quartic surface set up on  $C$  a (1, 1) correspondence of valency 1, while those of the ruled surface of order 14 set up a (3, 3)

\* The intersections of an elliptic curve with a primal are such that any one of them is determined by the rest.

† That there are six linearly independent quadrics containing the quartic ruled surface follows at once when its equations are written in the form

$$X/Y = Y/Z = X'/Y' = Y'/Z'.$$

‡ Cf. § 91.

correspondence of valency 1; these correspondences are both symmetrical and have therefore two common pairs of points, which means simply that there are two chords of  $C$  generators of both ruled surfaces\*. This shews that  $C_2$  and  $C_7^3$  intersect in two points.

**258.** There are  $\infty^1$  quartic ruled surfaces arising from the  $\infty^1 g_2^1$ 's on  $C$ ; we have thus  $\infty^2$  conics, lying in planes, on these surfaces. We may call these planes secant planes of  $M_3^9$ ; they meet  $M_3^9$  in conics, whereas an ordinary plane meets  $M_3^9$  in nine points. There are  $\infty^2$  secant planes; they form a locus of four dimensions. We can use arguments for these planes similar to those used in § 91.

Take any one of the secant planes and the conic  $\Gamma$  therein meeting  $C$  in  $X$  and  $Y$ ; take any point  $P$  in the plane and let  $PX, PY$  meet  $\Gamma$  again in  $X'$  and  $Y'$ . Then there are chords of  $C$  passing through  $X'$  and  $Y'$ , and these joined to  $X$  and  $Y$  respectively give the two trisecant planes of  $C$  which pass through  $P$ .

Hence we conclude that if there is a secant plane passing through a point  $P$  it must intersect both the trisecant planes of  $C$  through  $P$  in lines, provided that  $P$  is not on the conic in the secant plane.

**259.** Take a trisecant plane of  $C$  lying on  $\Omega$ , and suppose for the moment that none of the three chords of  $C$  which it contains is a generator of the quartic ruled surface. Then there are three generators of the quartic ruled surface passing one through each of the points in which the trisecant plane meets  $C$ . The secant plane which belongs to the same system of planes on  $\Omega$  as that to which the trisecant plane belongs meets the trisecant plane in a point and the quartic ruled surface in a conic; we thus have a [4] meeting the quartic ruled surface in a conic and three generators, which is impossible. We therefore conclude that the trisecant plane contains a generator of the quartic ruled surface. Incidentally, it must meet the other secant plane on  $\Omega$  in a line.

Suppose now that  $\Omega$  contains the planes  $\omega_1, \omega_2, \rho_1, \rho_2$  trisecant to  $C$  and also the secant planes  $\omega_0$  and  $\rho_0$ . Then  $\omega_0$  meets  $\rho_1$  and  $\rho_2$  in lines, while  $\rho_0$  meets  $\omega_1$  and  $\omega_2$  in lines.

Now  $\rho_0$  represents the plane of the double conic  $C_2$ . Hence  $C_7^3$  has two double points on  $C_2$  as well as the two simple points already mentioned; they are represented on  $\Omega$  by the planes  $\omega_1$  and  $\omega_2$ . The plane of  $C_2$  meets the sextic ruled surface in  $C_2$  counted twice and in two generators (these generators being represented on  $\Omega$  by the two points in which  $C$  is met by  $\rho_0$ ); these intersect in a point of  $C_7^3$ , and all the seven intersections of  $C_7^3$  with the plane of  $C_2$  are thus accounted for. The generators of the surface are trisecant to  $C_7^3$ .

The two points in which  $\rho_0$  meets  $C$  lie one in  $\omega_1$  and the other in  $\omega_2$ .

The projection of  $C_7^3$  on to a plane from one of its double points is a quintic with three double points. Thus there are two lines through the double point of  $C_7^3$  each of which meets  $C_7^3$  in two further points, and these are generators of the surface; the third generator through this point lies in the plane of  $C_2$ .

There are similar statements for the developable  $E_7^3$  and the planes  $\rho_1, \rho_2$ .

**260.** Of the  $\infty^5$  quadrics containing  $C$  and one secant plane (and therefore two) there are  $\infty^1$  which contain another (and therefore still another) secant plane. Then the ruled surface formed by the chords of  $C$  which lie on  $\Omega$  breaks up into two quartic surfaces and a ruled surface of order ten of which two generators pass through any point of  $C$ . By the same argument as in § 257, there are eight points of  $C$  for which these two generators coincide, and each ruled surface contains four tangents of  $C$ .

Any plane of  $\Omega$  is met by two generators of each quartic surface and by five generators of the other surface; we have thus in ordinary space a sextic ruled surface whose double curve is  $C_2 + D_2 + C_5$  and bitangent developable  $E_2 + F_2 + E_5$ .

Taking a section of the ruled surface of order ten by a [4] we have a curve  $C'$  of order ten, and for the correspondence between  $C$  and  $C'$

$$\alpha = 2, \quad \alpha' = 2, \quad p = 1, \quad \eta = 8, \quad \eta' = 4,$$

whence  $p' = 2$ , and  $C'$  is of genus 2.

Thus for the double curve of the ruled surface we have  $C_2 + D_2 + C_5^2$  and for the bitangent developable  $E_2 + F_2 + E_5^2$ .

Each quartic surface sets up on  $C$  a (1, 1) correspondence of valency 1, while the ruled surface of order ten sets up a (2, 2) correspondence of valency zero. This shews that, all these correspondences being symmetrical, there are no generators common to the two quartic surfaces, while each quartic surface has two generators in common with the ruled surface of order ten. Hence  $C_5^2$  has two simple intersections with each of  $C_2$  and  $D_2$ , while  $E_5^2$  has two simple planes in common with each of  $E_2$  and  $F_2$ .

**261.** Suppose that the quadric  $\Omega$  has  $\omega_1, \omega_2, \rho_1, \rho_2$  for trisecant planes of  $C$  and also contains the secant planes  $\omega_0, \rho_0, \omega_0', \rho_0'$ . Then the two intersections of  $\omega_0$  with  $C$  will lie one on  $\rho_1$  and the other on  $\rho_2$ , with similar statements for the other secant planes.

We conclude that  $C_2$  and  $D_2$  have two intersections each of which lies on  $C_5^2$ . The plane of either  $C_2$  or  $D_2$  meets the ruled surface in the double conic and two generators whose point of intersection lies on  $C_5^2$ . All the intersections of  $C_5^2$  with these planes are thus accounted for. The generators are chords of  $C_5^2$ .

The three generators through one of the two triple points consist of one in the plane of  $C_2$ , one in the plane of  $D_2$  and the trisecant of  $C_5^2$ .

The trisecants of  $C_5^2$  form a quadric surface; the intersection of this with the sextic surface consists of the two particular trisecants just mentioned and the curve  $C_5^2$  counted twice.

*Surfaces whose generators belong to a linear complex which is not special*

**262.** Suppose that we have on  $\Omega$  an elliptic sextic curve  $C$  contained in a prime which does not touch  $\Omega$ . The curve has two trisecants, and these lie on  $\Omega$ . We have in [3] correspondingly an elliptic sextic ruled surface with a double curve of order nine which has two triple points at each of which the three generators are co-planar; by the same argument as in § 256 the double curve is a  $C_9^4$  of genus 4.

Let us obtain this genus in another way.

The chords of  $C$  form a locus  $M_3^9$  on which the two trisecants are triple lines and  $C$  is a quadruple curve. Since any chord of  $C$  is met by two\* chords, other than those which pass through its intersections with  $C$ , there is a double surface  $F_2$  on  $M_3^9$ .

The order of  $F_2$  is the number of double points of a plane section of  $M_3^9$ . The genus of such a plane section can be calculated by an application of Zeuthen's formula, and the number of its double points is thereby determined. Take any plane section  $c$  of  $M_3^9$ , and consider the correspondence between  $C$  and  $c$ , points of the two curves corresponding when the line joining them is a chord of  $C$ . The correspondence is a (2, 5) correspondence; for through each point of  $c$  there passes a chord of  $C$  and through each point of  $C$  there pass five of its chords which meet  $c$ , these lying in the solid determined by the point of  $C$  and the plane of  $c$ . The number of branch-points of the correspondence on  $c$  is the number of tangents of  $C$  which meet  $c$ ; this number is 12—the order of the surface formed by the tangents of  $C$ . If  $P$  is a branch-point of the correspondence on  $C$  then two of the five chords of  $C$  which pass through  $P$  and meet  $c$  must coincide, so that the solid  $Pc$  must touch  $C$  at a point other than  $P$ . The tangent of  $C$  at this other point must be one of the 12 tangents of  $C$  which meet  $c$ , so that the correspondence has 48 branch-points on  $C$ . Thus

$$a = 2, \quad a' = 5, \quad p = 1, \quad \eta = 48, \quad \eta' = 12,$$

and Zeuthen's formula gives  $p' = 10$ . A plane curve of order nine and genus 10 has 18 double points; hence the double surface on  $M_3^9$  is a surface  $F_2^{18}$  of order 18. It can be shewn that  $C$  is a double curve on  $F_2^{18}$ ; the section of  $M_3^9$  by, for example, a quadriseccant plane of  $C$  consists of six chords of  $C$  together with a cubic having a double point and passing through the four points of  $C$  in the plane. Thus  $F_2^{18}$  meets  $\Omega$  in  $C$  counted twice, the trisecants counted three times and a residual curve of order 18.

Hence the chords of  $C$  which lie on  $\Omega$  form a ruled surface of order 18 (the intersection of  $\Omega$  and  $M_3^9$ ) on which  $C$  is a quadruple curve and the two trisecants are triple generators, there being also a double curve of order 18. The section of this ruled surface by a solid is a curve of order 18 lying on a

\* The projection of  $C$  from a chord on to a plane is an elliptic quartic curve with two double points.



quadric and meeting each generator in nine points; it has six quadruple points, two triple points, and eighteen double points. Thus its genus is

$$136 - 72 - 36 - 6 - 18 = 4,$$

so that  $C_9^4$  and  $E_9^4$  are also of genus 4.

**263.** If we project a normal elliptic sextic curve on to a [4] from a point lying in a secant plane this secant plane meets [4] in a line which intersects the two trisecants of the projected curve  $C$ ; through any point of the line pass two chords of  $C$ , and all these chords form a quartic ruled surface.

There are  $\infty^{14}$  quadrics in [4]; hence there are  $\infty^3$  passing through  $C$ , which will incidentally contain the two trisecants, and  $\infty^1$  of these contain the quartic ruled surface.

If we consider one of these quadrics as a section of  $\Omega$  by the [4] containing  $C$ , the points of  $C$  represent the generators of an elliptic sextic ruled surface in [3] whose double curve is  $C_2 + C_7^3$  exactly as in § 257;  $C_7^3$  has two double points and two simple points on  $C_2$ . Similarly for the bitangent developable  $E_2 + E_7^3$ .

**264.** Given a  $g_2^1$  on the normal elliptic sextic curve the joins of pairs of points form a quartic ruled surface with  $\infty^1$  directrix conics each meeting the curve in two points; but these pairs of points on the conics themselves form a  $g_2^1$ , they are cut out by [4]'s passing through two generators of the quartic surface. Given two points on the curve, that  $g_2^1$  is determined in which they form a pair; but there are four series  $g_2^1$  giving rise to quartic ruled surfaces such that this pair of points lies on a conic of the surface; this last statement is at once clear on using elliptic arguments.

The planes of the directrix conics of a quartic ruled surface form a locus  $V_3^3$ , and in [5] two loci  $V_3^3$  without a common surface meet in a curve of order nine. It is therefore clear that there are points not lying on the elliptic curve, or on any of its chords, through which there pass two secant planes. If we project from such a point on to a [4] the two secant planes meet [4] in lines, each of which meets both the trisecants of the projected curve  $C$ ; through any point of either line there pass two chords of  $C$ , and all such chords form a quartic ruled surface on which the line is a double line.

Of the  $\infty^2$  quadrics passing through  $C$  there is one containing both the quartic ruled surfaces; regarding this as a prime section of  $\Omega$  the points of  $C$  represent the generators of an elliptic sextic ruled surface in ordinary space whose double curve is  $C_2 + D_2 + C_5^2$  and bitangent developable  $E_2 + F_2 + E_5^2$  exactly as in § 260. Here the generators of the surface belong to a linear complex, and the three generators at a triple point lie in a tritangent plane.

**265.** Suppose now that the curve  $C$  lying on  $\Omega$  is contained in a [4] and has a double point  $P$ . Then  $P$  represents a double generator  $G$  of

the ruled surface. The chords of  $C$  form a locus  $M_3^8$ , and any plane on  $\Omega$  meets the [4] containing  $C$  in a line meeting this locus in eight points. Thus the ruled surface in [3] has a double curve  $G + C_8$  and a bitangent developable  $G + E_8$ .

The chords of  $C$  lying on  $\Omega$  form a ruled surface of order 16; there are four of them passing through every point of  $C$ , except that through  $P$  there pass only two  $PQ, PR$ , while through  $Q$  and  $R$  there pass only three. The twelve tangents of  $C$  which lie on  $\Omega$  belong to this ruled surface. We have on  $C$  a (4, 4) correspondence of valency 2; there are twenty-four points of  $C$  for which two of the four corresponding points coincide. Now these points include  $P$ . For we regard the double point of  $C$  as lying on two distinct branches of the curve; suppose that it is  $P_1$  regarded as a point of one branch and  $P_2$  regarded as a point of the other. Then the tangent prime of  $\Omega$  at  $P_1$  meets  $C$  twice at  $P_1$ , twice at  $P_2$ , and once at each of  $Q$  and  $R$ . Thus corresponding to  $P_1$  we have  $Q + R + 2P_2$ ; similarly, corresponding to  $P_2$  we have  $Q + R + 2P_1$ . The point  $P$  in fact counts for four among the twenty-four coincidences. The points  $Q$  and  $R$  are not included, since  $P_1$  and  $P_2$  are on different branches of  $C$ . We have then twenty coincidences, which give generators of the surface in [3] touching the double curve  $C_8$ .

Taking a section of the ruled surface we have a curve  $C'$  of order 16, and in the correspondence between  $C$  and  $C'$

$$\alpha = 2, \quad \alpha' = 4, \quad p = 1, \quad \eta = 20, \quad \eta' = 12,$$

so that  $p' = 3$ , and  $C'$  is of genus 3.

The curve  $C_8^3$  has two double points and two simple points on  $G^*$ . Through either of the double points there passes one line meeting  $C_8$  in two other points not on  $G$ ; this is the generator, other than  $G$ , which passes through this point; it is represented on  $\Omega$  by one of the points  $Q, R$ .

We can find the genus of  $C_8^3$  and  $E_8^3$  in another way.

The chords of  $C$  form a locus  $M_3^8$  on which  $C$  is a quadruple curve; the quartic cone projecting  $C$  from  $P$  is a double surface on  $M_3^8$ , and  $P$  is a sextuple point. Making use of an argument similar to that of § 262, we find that there is a double surface  $F_2^{11}$  on  $M_3^8$ , but this is composed of the quartic cone and a double surface  $F_2^7$ .  $C$  is a simple curve on  $F_2^7$ ; a quadrisecant plane of  $C$  meets  $M_3^8$  in six chords of  $C$  and a conic passing through the four points of  $C$ ; the intersection of  $\Omega$  and  $F_2^7$  consists of  $C$  and an octavic curve, while  $\Omega$  meets the quartic cone in  $C$  and two lines  $PQ, PR$ .

Hence the chords of  $C$  which lie on  $\Omega$  form a ruled surface of order 16 with two double generators, a double curve of order 8 and the quadruple curve  $C$ . The section of this ruled surface by a solid is a curve of order 16 lying on a quadric and meeting all the generators in eight points; it has six quadruple points and ten double points. Its genus is therefore

$$105 - 56 - 36 - 10 = 3,$$

and therefore  $C_8^3$  and  $E_8^3$  are also of genus 3.

\* Cf. § 102.

266. Let us project a normal elliptic sextic from a point on one of its chords. We obtain a curve  $C$  with a double point  $P$ , and there are four lines through  $P$  which are double lines of quartic ruled surfaces whose generators join pairs of points of four  $g_2^1$ 's on  $C$ . Since there are  $\infty^3$  quadrics passing through  $C$  there are  $\infty^1$  containing one of these quartic ruled surfaces.

Now regard one of these as a section of  $\Omega$  by a [4]. The surface formed by the chords of  $C$  which lie on  $\Omega$  here breaks up into the quartic ruled surface and a ruled surface of order 12; every plane of  $\Omega$  meets the [4] containing  $C$  in a line meeting the former surface in two points and the latter in six. Hence  $C$  represents the generators of an elliptic sextic ruled surface in [3] contained in a linear complex; the double curve is  $C_2 + G + C_6$  and the bitangent developable  $E_2 + G + E_6$ .

The quartic surface sets up on  $C$  a (1, 1) correspondence of valency 1, while the ruled surface of order 12 sets up a (3, 3) correspondence of valency 1. This gives sixteen as the number of points of  $C$  at which two of the three generators of this latter surface coincide; but  $P$  is counted four times in this result.

Of the twelve tangents of  $C$  which lie on  $\Omega$  four belong to the quartic ruled surface and eight to the ruled surface of order 12. If we take the section  $C'$  by a solid we have a curve of order 12, and for the correspondence between  $C$  and  $C'$

$$\alpha = 2, \quad \alpha' = 3, \quad p = 1, \quad \eta = 12, \quad \eta' = 8,$$

so that  $p' = 2$ , and  $C'$  is of genus 2.

Thus the ruled surface in [3] has a double curve  $C_2 + G + C_6^2$  and a bitangent developable  $E_2 + G + E_6^2$ .

The plane of  $C_2$  contains  $G$ , and  $C_6$  passes through the two intersections of  $G$  and  $C_2$ ;  $C_6^2$  also meets  $G$  in two other points\*. There are two further intersections of  $C_6^2$  and  $C_2$ .

There is one other generator passing through either intersection of  $C_2$  and  $G$ ; this is the unique line, other than  $G$ , passing through the point, which is trisecant to  $C_6^2$ .

There are precisely similar statements for the bitangent developable.

*Surfaces with a directrix line which is not a generator*

267. If  $C$  is of the type III (A) it meets every plane  $\omega$  of  $\Omega$  through  $O$  in two points and every plane  $\rho$  of  $\Omega$  through  $O$  in four points; two chords of the curve passing through  $O$ . An arbitrary plane  $\rho$  of  $\Omega$  meets  $T$  in a line, and the  $\omega$ -plane through this line joins it to  $O$ . This contains one chord of the curve; there are eight others meeting the line. We thus

\* For this statement cf. § 102.

have in [3] a ruled surface with a directrix line  $R$ ; through any point of  $R$  there pass two generators, while any plane through  $R$  contains four generators; the double curve is  $R + C_8$ .

The pairs of points of  $C$  which lie in the  $\omega$ -planes through  $O$  form a  $g_2^1$ , and so the chords of  $C$  in these planes form a quartic ruled surface. The  $\rho$ -planes through  $O$  cut out on  $C$  sets of a  $g_4^1$ ; they thus set up a (3, 3) correspondence on  $C$  of valency 1, and there are sixteen points of  $C$  for which two of the three corresponding points coincide.

The curve  $C$  is touched by four  $\omega$ -planes and eight  $\rho$ -planes; each of these last planes contains two of the sixteen points just mentioned. The ruled surface formed by the chords of  $C$  which lie in the  $\rho$ -planes is of order 14; take  $C'$ , a section of it by a [3]. Then for the correspondence between  $C$  and  $C'$

$$\alpha = 2, \quad \alpha' = 3, \quad p = 1, \quad \eta = 16, \quad \eta' = 8,$$

so that  $p' = 3$ , and  $C'$  is of genus 3.

Thus the double curve is  $R + C_8^3$ . The generators of the ruled surface meet  $R$  and are trisecants of  $C_8^3$ . Since  $C$  has two trisecants  $C_8^3$  has two triple points;  $R$  is a chord of  $C_8^3$ , the points in which it meets  $C_8^3$  being represented on  $\Omega$  by the  $\omega$ -planes containing the two chords of  $C$  through  $O$ .

Any plane  $\omega$  of  $\Omega$  meets  $T$  in a line, and the plane  $\rho$  through this line joins it to  $O$ . This contains four points and therefore six chords of  $C$ ; there are three other chords meeting the line. Thus the bitangent developable of the surface is  $6R + E_3^0$ ; the planes of  $E_3^0$  are the planes of the pairs of generators which intersect in the points of  $R$ , and two of these planes pass through  $R$ , being represented on  $\Omega$  by the  $\rho$ -planes containing the two chords of  $C$  through  $O$ .

When  $C$  is of the type III (B) we have similarly a double curve  $6R + C_3^0$  and a bitangent developable  $R + E_3^3$ .  $R$  is a chord of  $C_3^0$ ;  $E_3^3$  has two triple planes and two of its planes pass through  $R$ . Through any point of  $R$  there pass four generators, while any plane through  $R$  contains two generators.

**268.** Consider again a normal elliptic sextic in [5] and a  $g_2^1$  thereon. We have a quartic ruled surface, and the planes of the directrix conics of this form a  $V_3^3$ . Now project on to a [4] from a point of this  $V_3^3$  which is not on a chord of the curve. The projection of  $V_3^3$  is a quadric point-cone\* and  $C$ , the projected curve, lies on this cone, meeting all the planes of one system in two points and all of the other system in four points; but each of these sets of four points consists of two pairs of the same  $g_2^1$ .

We can then regard this quadric point-cone as the section of  $\Omega$  by a tangent prime  $T$ , and  $C$  is evidently of the type III (A) or III (B).

Consider for definiteness a curve  $C$  of the type III (A). The four points in which  $C$  is met by a  $\rho$ -plane form two pairs of the same  $g_2^1$ ; hence the chords of  $C$  which lie in  $\rho$ -planes now form a quartic ruled surface  $Q_\rho$  and a ruled surface  $R_2^{10}$  of order ten on which  $C$  is a double curve; the chords of  $C$  in  $\varpi$ -planes form as before a quartic ruled surface  $Q_\varpi$ . The trisecants of  $C$  are generators of  $Q_\rho$  and double generators of  $R_2^{10}$ . The surface  $R_2^{10}$  sets up on  $C$  a symmetrical (2, 2) correspondence of valency zero, and we can deduce that  $R_2^{10}$  and  $Q_\rho$  have two common generators.

The number of points of  $C$  at which the two generators of  $R_2^{10}$  coincide is eight; also, of the twelve tangents of  $C$  which lie on  $\Omega$  four belong to  $Q_\varpi$ , four to  $Q_\rho$  and four to  $R_2^{10}$ . Hence, considering the correspondence between  $C$  and a section  $C'$  of  $R_2^{10}$  by a solid,

$$\alpha = 2, \quad \alpha' = 2, \quad p = 1, \quad \eta = 8, \quad \eta' = 4,$$

so that  $p' = 2$ .

A plane  $\rho$  of  $\Omega$  meets  $T$  in a line; the nine chords of  $C$  meeting this line consist of the chord in the plane  $\varpi$  joining the line to  $O$ , of two generators of  $Q_\rho$  and six of  $R_2^{10}$ . Thus the ruled surface has a double curve  $R + C_2 + C_6^2$ ;  $C_6^2$  has two double points through both of which  $C_2$  passes, while  $C_6^2$  and  $C_2$  have two further simple intersections.  $C_6^2$  meets  $R$  in two points, but  $R$  and  $C_2$  do not meet. The bitangent developable is still  $6R + E_3^0$ .

Similarly, if  $C$  is of the type III (B) we have in [3] an elliptic sextic ruled surface whose double curve is given by  $6R + C_3^0$  and bitangent developable by  $R + E_2 + E_6^2$ ; there are properties of this developable dual to those already given for the double curve of the ruled surface just mentioned.

269. There are still more special examples of surfaces in [3] belonging to the types III (A) and III (B).

Consider again the elliptic normal sextic curve in [5]. We can have sets of four points  $\alpha, \alpha', \beta, \beta'$  upon the curve such that  $\alpha$  and  $\alpha'$  form a pair of a  $g_2^1$  including also the pair  $\beta\beta'$ , while the pairs  $\alpha\beta$  and  $\alpha'\beta'$  belong to a second  $g_2^1$ \*. Consider the quartic surface formed by the first  $g_2^1$ ; the second  $g_2^1$  gives an involution of pairs of generators on this surface; the joins of pairs of points in which these pairs of lines meet any directrix conic of the surface thus pass through a fixed point. In fact the solids determined by the sets of four points such as  $\alpha, \alpha', \beta, \beta'$  (such a set being determined by one point) have a line in common lying on the  $V_3^3$  formed by the planes of the directrix conics.

\* We take the elliptic arguments so that

$$\alpha + \alpha' \equiv c, \quad \beta + \beta' \equiv c, \quad \alpha + \beta \equiv c + \frac{1}{2}\omega, \quad \alpha' + \beta' \equiv c + \frac{1}{2}\omega,$$

where  $\omega$  is a period of the elliptic functions. Any three of these congruences involve the fourth.

Let us then project on to a [4] from a point of this line. We obtain an elliptic sextic curve lying on a quadric point-cone, meeting all planes of one system in four points and all of the other system in two points; but the sets of four points are such that, if  $\alpha, \alpha', \beta, \beta'$  is one of them, the pairs  $\alpha\alpha'$  and  $\beta\beta'$  belong to one  $g_2^1$ , while the pairs  $\alpha\beta$  and  $\alpha'\beta'$  belong to another.

Suppose, for definiteness, that  $C$  is of the type III (A). The chords of  $C$  lying in the  $\omega$ -planes through  $O$  form as before a quartic ruled surface  $Q_\omega$ ; those which lie in the  $\rho$ -planes through  $O$  form two quartic ruled surfaces  $Q_\rho$  and  $Q_{\rho'}$  and a sextic ruled surface  $R_2^6$ . The trisecants of  $C$  are generators of each of the three surfaces  $Q_\rho, Q_{\rho'}, R_2^6$ . The surface  $R_2^6$  gives on  $C$  a (1, 1) correspondence of valency  $-1$ , so that this surface has two generators in common with each of the surfaces  $Q_\rho, Q_{\rho'}$  as well as the trisecants.

An arbitrary plane  $\rho$  meets  $T$  in a line; this meets  $Q_\rho$  in two points,  $Q_{\rho'}$  in two points,  $R_2^6$  in four points and is met by the chord of  $C$  lying in the  $\omega$ -plane joining it to  $O$ . Also it is easily seen that the surface  $R_2^6$  is elliptic.

We have thus in [3] a surface whose double curve is  $R + C_2 + D_2 + C_4^1$ ;  $C_2$  and  $D_2$  have two intersections through each of which  $C_4^1$  passes, while  $C_4^1$  meets each conic in two other points.  $R$  is a chord of  $C_4^1$ . The plane of either conic meets the surface in this conic counted twice and a pair of generators which intersect on  $R$ .

The bitangent developable is again  $6R + E_3^0$ .

Similarly we have a surface reciprocal to this when  $C$  is of the type III (B).

**270.** If  $C$  is of the type III (C) it meets every plane  $\omega$  of  $\Omega$  through  $O$  in two points and every plane  $\rho$  of  $\Omega$  through  $O$  in four points; it has a double point  $P$  and one of its chords passes through  $O$ .

An arbitrary plane  $\rho$  of  $\Omega$  meets  $T$  in a line; the plane joining this line to  $P$  meets  $\Omega$  again in a second line passing through  $P$ . The plane  $\omega$  joining the line to  $O$  contains one chord of  $C$ , and there are seven others meeting the line. Hence we have in [3] a ruled surface with a double generator  $G$ ; the double curve is  $R + G + C_7$ .

The chords of  $C$  which lie in the  $\omega$ -planes through  $O$  form a quartic ruled surface  $Q_\omega$ , while those which lie in the  $\rho$ -planes through  $O$  form a ruled surface  $R_2^{12}$  of order 12 on which  $C$  is a triple curve. The generators of  $R_2^{12}$  set up on  $C$  a symmetrical (3, 3) correspondence of valency 1; this shews that there are sixteen points of  $C$  at which two of the three generators of  $R_2^{12}$  coincide. Through  $P$  there pass two chords  $PQ, PR$  of  $C$  which lie on  $\Omega$ ; the point  $P$  is not included among the sixteen coincidences. To the point  $P_1$  on one branch of  $C$  there correspond three distinct points  $Q, R, P_2$ ,

while to the point  $P_2$  on the other branch there correspond three distinct points  $Q, R, P_1$ .

$C$  is touched by four  $\omega$ -planes and eight  $\rho$ -planes through  $O$ . Taking a section  $C'$  of  $R_2^{12}$  by a solid we have, for the correspondence between  $C$  and  $C'$ ,

$$\alpha = 2, \quad \alpha' = 3, \quad p = 1, \quad \eta = 16, \quad \eta' = 8,$$

so that  $p' = 3$ , and  $C'$  is of genus 3. Hence the double curve is  $R + G + C_7^3$ .

The plane of the two tangents of  $C$  at  $P$  meets the  $\omega$ -plane through  $OP$  in a generator of  $Q_\omega$  and the  $\rho$ -plane through  $OP$  in a generator of  $R_2^{12}$ . The plane  $\omega$  through this generator of  $R_2^{12}$  represents a point of intersection of  $C_7^3$  and  $G$ .  $C_7^3$  has also two double points on  $G$ ; these are represented on  $\Omega$  by the  $\omega$ -planes through the two double generators of  $R_2^{12}$ ,  $PQ$  and  $PR$ .

$R$  and  $G$  intersect. Also  $C_7^3$  meets  $R$ , the point of intersection being represented on  $\Omega$  by the  $\omega$ -plane which contains the chord of  $C$  through  $O$ .

An arbitrary plane  $\omega$  of  $\Omega$  meets  $T$  in a line; the plane  $\rho$  through this line joins it to  $O$  and contains six chords of  $C$ ; there are two other chords of  $C$  meeting the line. Hence the ruled surface has a bitangent developable  $6R + G + E_2$ . There is a plane of  $E_2$  passing through  $R$ ; it is represented on  $\Omega$  by the  $\rho$ -plane containing the chord of  $C$  passing through  $O$ .

Similarly, when  $C$  is of the type III (D) we have a double curve  $6R + G + C_2$  and a bitangent developable  $R + G + E_7^3$ .  $R$  and  $G$  intersect;  $C_2$  meets  $R$ .  $E_7^3$  has two double planes and one ordinary plane passing through  $G$  and one ordinary plane passing through  $R$ .

**271.** Consider again a normal elliptic sextic curve in [5]. Through any one of its chords there pass four secant planes; consider any one of these and the  $\infty^1$  others belonging to the same quartic ruled surface and forming a  $V_3^3$ . Projecting from a point of the chord on to a [4] we have an elliptic sextic curve  $C$  with a double point and lying on a quadric point-cone; it meets all the planes of one system in two points and all of the other system in four points. Regarding the cone as a section of  $\Omega$  by a tangent prime  $T$  we have clearly a curve  $C$  of one of the types III (C) and III (D); but here the four points of  $C$  in a plane of the cone consist of two pairs of the same  $g_2^1$ .

Suppose then that  $C$  is of the type III (C). The chords of  $C$  lying in the  $\omega$ -planes form a quartic ruled surface  $Q_\omega$ , while those lying in the  $\rho$ -planes form a quartic ruled surface  $Q_\rho$  together with a ruled surface  $R_2^8$  of order eight on which  $C$  is a double curve.

An arbitrary plane  $\rho$  meets  $T$  in a line; this line is met by one generator of  $Q_\omega$ , two of  $Q_\rho$  and five of  $R_2^8$ . Hence the ruled surface in [3] has a double curve  $R + G + C_2 + C_5$ .

The surface  $R_2^8$  sets up on  $C$  a symmetrical (2, 2) correspondence of valency zero.  $R_2^8$  and  $Q_\rho$  have two common generators. Also there are eight points of  $C$  at which the two generators of  $R_2^8$  coincide, and of the twelve tangents of  $C$  which lie on the point-cone four belong to  $Q_\omega$ , four to  $Q_\rho$  and four to  $R_2^8$ . Thus, if  $C'$  is the section of  $R_2^8$  by a solid, we have a correspondence between  $C$  and  $C'$  for which

$$\alpha = 2, \quad \alpha' = 2, \quad p = 1, \quad \eta = 8, \quad \eta' = 4,$$

so that  $p' = 2$  and  $C'$  is of genus 2.

Hence the double curve of the ruled surface is  $R + G + C_2 + C_5^2$ . The plane of  $C_2$  contains  $G$ , while  $C_5^2$  passes through their two intersections.  $C_5^2$  meets  $G$  in a third point and meets  $R$  in one point.

The bitangent developable is still  $6R + G + E_2$ .

Similarly, when  $C$  is of the type III (D) we have in [3] an elliptic sextic ruled surface whose double curve is  $6R + G + C_2$  and bitangent developable  $R + G + E_2 + E_5^2$ .

**272.** If  $C$  is of the type III (E) it meets every plane of  $\Omega$  through  $O$  in three points and has three chords passing through  $O$ . We have a surface in [3] with a double curve  $3R + C_6$  and a bitangent developable  $3R + E_6$ .

The planes of either system of  $\Omega$  which pass through  $O$  cut out a  $g_3^1$  on  $C$ , this has six double points; of the twelve tangents of  $C$  which lie on  $\Omega$  there are six in each system of planes. The chords of  $C$  which lie in either system of planes through  $O$  form a ruled surface  $R_2^9$ , of order nine, on which  $C$  is a double curve; we have thus a (2, 2) correspondence of valency 1 on  $C$ , so that there are six points of  $C$  at which the two generators of  $R_2^9$  coincide. Then, taking a section  $C'$  of  $R_2^9$ , there is a correspondence between  $C$  and  $C'$  for which

$$\alpha = 2, \quad \alpha' = 2, \quad p = 1, \quad \eta = 6, \quad \eta' = 6,$$

so that  $p' = 1$ , and  $C'$  is of genus 1. Hence  $C_6$  is an elliptic curve  $C_6^1$  and  $E_6$  is an  $E_6^1$ .

Of the two trisecants of  $C$  one is joined to  $O$  by a  $\omega$ -plane and the other by a  $\rho$ -plane. The  $\rho$ -plane through the first of these trisecants represents a triple plane of  $E_6^1$ , while the  $\omega$ -plane through the second trisecant represents a triple point of  $C_6^1$ .

$R$  is a trisecant of  $C_6^1$ ; the three points in which it meets  $C_6^1$  are represented by the  $\omega$ -planes through those three chords of  $C$  which pass through  $O$ . Similarly there are three planes of  $E_6^1$  passing through  $R$ .

**273.** If  $C$  is of the type III (F) it meets every plane of  $\Omega$  passing through  $O$  in three points; two of its chords pass through  $O$  and it has a



double point  $P$ . We now have a surface whose double curve is  $3R + G + C_5$  and bitangent developable  $3R + G + E_5$ .

The chords of  $C$  which lie in either system of planes of the cone form a ruled surface  $R_2^8$  on which  $C$  is a double curve. The planes of either system cut out a  $g_3^1$  on  $C$ ; there are six points of  $C$  at which the two generators of  $R_2^8$  coincide. Of the twelve tangents of  $C$  which lie on  $\Omega$  there are six in either system of planes. Taking then a section  $C'$  of  $R_2^8$  we have for the correspondence between  $C$  and  $C'$

$$\alpha = 2, \quad \alpha' = 2, \quad p = 1, \quad \eta = 6, \quad \eta' = 6,$$

whence  $p' = 1$ , so that  $C'$  is of genus 1. Hence we have a double curve  $3R + G + C_5^1$  and bitangent developable  $3R + G + E_5^1$ .

Since there are two chords of  $C$  passing through  $O$  there are two points of  $R$  at which two of the three generators which meet there lie in a plane with  $R$ . These two points are intersections of  $R$  and  $C_5^1$ ; the planes are planes of  $E_5^1$  passing through  $R$ .

There are two chords of  $C$  passing through  $P$  and lying on  $\Omega$ ; one is joined to  $O$  by a plane  $\omega$  and the other by a plane  $\rho$ .  $C_5^1$  has a double point and a simple point on  $G$ ;  $E_5^1$  has a double plane and a simple plane through  $G$ .

### *Surfaces with a directrix line which is also a generator*

274. If  $C$  is of the type IV (A) it passes through  $O$ , meeting every plane  $\omega$  of  $\Omega$  through  $O$  in two points other than  $O$  and every plane  $\rho$  of  $\Omega$  through  $O$  in three points other than  $O$ ; one of its trisecants passes through  $O$ .

An arbitrary plane  $\rho$  meets  $T$  in a line; the plane  $\omega$  through this line passes through  $O$  and contains three points and therefore three chords of  $C$ ; there are six further chords meeting the line. Thus the surface in [3] has a double curve  $3R + C_6$ ; similarly it has a bitangent developable  $6R + E_3$ .

To obtain a representation of  $C_6$  we consider a section of the ruled surface formed by the chords of  $C$  which lie in  $\rho$ -planes through  $O$  and do not pass through  $O$ . There are ten tangents of  $C$  lying on  $\Omega$ , this being the number of tangents of an elliptic quintic curve on a quadric which lie on the quadric. The planes  $\omega$  cut out a  $g_2^1$  on  $C$  (exclusive of  $O$ ) and thus four tangents of  $C$  lie in  $\omega$ -planes; the remaining six lie in  $\rho$ -planes. These  $\rho$ -planes cut out a  $g_3^1$  on  $C$  which is a symmetrical (2, 2) correspondence of valency 1; there are thus six points of  $C$  for which the two corresponding points of the  $g_3^1$  coincide.

If then  $C'$  denotes a section of the ruled surface formed by the chords

of  $C$  which lie in the  $\rho$ -planes and do not pass through  $O$  we have for the correspondence between  $C$  and  $C'$

$$\alpha = 2, \quad \alpha' = 2, \quad p = 1, \quad \eta = 6, \quad \eta' = 6,$$

so that  $p' = 1$ , and  $C'$  is of genus 1.

Thus the double curve is  $3R + C_6^1$ .

Similarly the bitangent developable is  $6R + E_3^0$ ; the chords of  $C$  which lie in  $\omega$ -planes and do not pass through  $O$  form a rational quartic ruled surface.

The second trisecant of  $C$  is joined to  $O$  by a  $\rho$ -plane; this represents a tritangent plane of the surface which passes through  $R$  and contains three generators concurrent in a triple point of  $C_6^1$ .

$R$  is itself a generator; through any point of  $R$  there pass two other generators, and there is one point of  $R$  at which these lie in a plane through  $R$ . This point of  $R$  is on  $C_6^1$ ;  $R$  is in fact a trisecant of  $C_6^1$ , the other generators being ordinary chords. There are two planes of  $E_3^0$  passing through  $R$ .

Similarly, if  $C$  is of the type IV (B) we have a surface in [3] whose double curve is  $6R + C_3^0$  and bitangent developable  $3R + E_6^1$ .

**275.** If  $C$  is of the type IV (C) it passes through  $O$ , meeting every  $\omega$ -plane of  $\Omega$  through  $O$  in two points other than  $O$  and every  $\rho$ -plane of  $\Omega$  through  $O$  in three points other than  $O$ , and has a double point  $P$ .

An arbitrary plane  $\rho$  of  $\Omega$  meets  $T$  in a line; the plane  $\omega$  through this line joins it to  $O$  and contains three points and therefore three chords of  $C$ . There are five further chords of  $C$  meeting the line; so that the double curve of the surface in [3] is  $3R + G + C_5$ . Similarly the bitangent developable is  $6R + G + E_2$ . Just as in the last article we find that  $C_5$  is an elliptic curve  $C_5^1$ .

There are two chords of  $C$  which pass through  $P$  and lie on  $\Omega$ ; one of these is  $PO$ ; let the other be  $PQ$ . The plane  $POQ$  is a  $\rho$ -plane.  $R$  and  $G$  intersect; their plane contains the generator represented by  $Q$ .  $C_5^1$  has a double point at the intersection of this generator with  $G$  and meets  $G$  in one other point.  $R$  is a chord of  $C_5^1$ . The  $\rho$ -plane through the tangent of  $C$  at  $O$  meets  $C$  in two other points; the  $\omega$ -planes which contain the chords of  $C$  joining these two points to  $O$  represent the two points of intersection of  $R$  and  $C_5^1$ .

Similarly, when  $C$  is of the type IV (D) we have in [3] a surface whose double curve is  $6R + G + C_2$  and bitangent developable  $3R + G + E_5^1$ .

**276.** If  $C$  is of the type V it has a double point at  $O$  and meets every plane of  $\Omega$  passing through  $O$  in two points other than  $O$ . An arbitrary plane of  $\Omega$  meets  $T$  in a line and the plane of the opposite system through this line joins it to  $O$ ; this contains three chords of  $C$  and there are three others meeting it, the projection of  $C$  from the line being a plane

sextic with a quadruple point and three double points. Hence the surface has a double curve  $6R + C_3$  and a bitangent developable  $6R + E_3$ ;  $C_3$  and  $E_3$  are rational as being represented by the pairs of two series  $g_2^1$  on  $C$ .  $R$  is not only a directrix but also a double generator; it is a chord of  $C_3^0$  and an axis of  $E_3^0$ .

*Surfaces whose generators belong to a linear congruence*

277. If  $C$  belongs to any of the types VI it has two directrices  $R$  and  $R'$ .

In the type VI (A) we have two double generators  $G$  and  $H$ ; the double curve is  $6R + G + H + R'$  and the bitangent developable  $R + G + H + 6R'$ .

In the type VI (B) we have three double generators  $G$ ,  $H$  and  $J$ ; the double curve is  $3R + G + H + J + 3R'$  and the bitangent developable  $3R + G + H + J + 3R'$ .

In the type VI (C) we have a triple generator  $G$ ; the double curve is  $3R + 3G + 3R'$  and the bitangent developable  $3R + 3G + 3R'$ .

If  $C$  belongs to any of the types VII there is a single directrix  $R$ .

In the type VII (A)  $R$  is a double generator and there are also two other double generators  $G$  and  $H$ ; the double curve is  $7R + G + H$ , which gives also the bitangent developable. The curve  $C$  is projected from a point of the cone into a plane sextic having two double points and a quadruple point with only three distinct tangents.

In the type VII (B) we have a double curve and bitangent developable  $6R + G + H + J$ . The curve  $C$  is projected from a point of the cone into a plane sextic with three double points and a triple point at which all the branches touch.

In the type VII (C) we have a double curve and bitangent developable  $6R + 3G$ ; the curve  $C$  is projected from a point of the cone into a plane sextic with an ordinary triple point and a triple point at which all the branches touch.

*The normal elliptic ruled surfaces of a given order  $n$*

278. In order to carry out the next steps in our investigation we require a knowledge of the properties of the normal elliptic ruled surfaces of the sixth order. We will then take this opportunity of obtaining some of the most fundamental properties of the normal elliptic ruled surfaces of any order  $n$ ; in particular, we shall investigate how many different types of surfaces there are which are projectively distinct from one another, and give some account of the curves which lie upon them. We shall also obtain methods for generating the surfaces by correspondences between two curves. The results are due originally to Segre\*.

\* See his paper, "Ricerche sulle rigate ellittiche di qualunque ordine," *Atti Torino*, 21 (1886), 868.

If we have an elliptic ruled surface of order  $n$  which is not a cone then, as is seen by an argument precisely similar to that used for a quintic surface in § 160, the surface belongs to a space of dimension  $n - 1$  at most. If the surface is contained in a space of dimension less than  $n - 1$  then we take a quadric through one of its generators; this quadric meets the surface again in a curve of order  $2n - 1$  and genus  $n$  which, just as in § 160, is the projection of a normal curve in  $[n - 1]$ . We thus prove, just as for quintic surfaces, that *an elliptic ruled surface of order  $n$  is normal in  $[n - 1]$ ; and any elliptic ruled surface of order  $n$  which belongs to a space of dimension less than  $n - 1$  is the projection of a normal surface in  $[n - 1]$ .*

279. Consider now a normal elliptic ruled surface  $F$  in  $[n - 1]$ . We shall suppose that  $F$  is not a cone, that it does not break up into separate surfaces, and that it is not contained in a space of dimension less than  $n - 1$ .

Any space which is contained in  $[n - 1]$  and meets  $F$  in an infinite number of points may contain only a number of generators of  $F$ , or else a number of generators together with a directrix curve of  $F$ ; it cannot contain two directrix curves of  $F$  for then it would contain the surface entirely.

If we have on  $F$  a curve of order  $\nu < n - 1$  this curve is contained in a space of dimension less than or equal to  $n - 2$ ; it is therefore a directrix curve of  $F$  and is elliptic. Also a curve of order  $n - 1$  must be a directrix curve of  $F$  and therefore elliptic; for if it met the generators of  $F$  in more than one point it would belong to  $[n - 1]$  and be a rational normal curve, and Zeuthen's formula shews that we cannot have a  $(1, k)$  correspondence between an elliptic curve and a rational curve. Hence *every curve on  $F$  of order  $\nu < n - 1$  is an elliptic curve and meets each generator in one point.*

A directrix curve of  $F$  of order  $\nu$  cannot be contained in a space of dimension less than  $\nu - 1$ . For if it were contained in a space  $[\nu - 2]$  this space, together with  $n - \nu - 1$  arbitrary generators, would determine a space  $[n - 3]$  meeting  $F$  in a composite curve of order  $n - 1$ . Then the pencil of primes through  $[n - 3]$  would give the generators of  $F$ , which is impossible since  $F$  is not rational. Hence *every curve of  $F$  of order  $\nu \leq n - 1$  is an elliptic normal curve.* This same argument shews that a curve of order  $\nu < n - 1$  cannot be contained in a space  $[\nu - 2]$  when it is composite, i.e. consists of a directrix curve and a certain number of generators. It is true in this case also that the curve belongs to a space  $[\nu - 1]$  and not to a space of higher dimension; for if it consists of  $\mu$  generators and an elliptic normal curve of order  $\nu - \mu$  this latter lies in a  $[\nu - \mu - 1]$  and this, together with the  $\mu$  generators, determines a  $[\nu - 1]$  containing the whole of the composite curve. Hence any curve of  $F$  of order  $\nu < n - 1$ , whether simple or composite, belongs to a space  $[\nu - 1]$ .

We cannot have on  $F$  two directrix curves  $C_\nu, C_{\nu'}$  of orders  $\nu$  and  $\nu'$  such that  $\nu + \nu' < n$ . For  $C_\nu$  is contained in a space  $S_{\nu-1}$  and  $C_{\nu'}$  is contained in a space  $S_{\nu'-1}$ , so that  $F$  is contained in a space  $[\nu - 1 + \nu' - 1 + 1]$  or  $[\nu + \nu' - 1]$ . Hence, since  $F$  is not contained in a space of less dimension than  $n - 1$ ,  $\nu + \nu' \geq n$ . Hence, *if  $F$  contains a curve of order less than  $\frac{1}{2}n$  it is the only such curve and is the minimum directrix.*

Similarly, if we have two curves  $C_\nu, C_{n-\nu}$  on the surface, the sum of whose orders is  $n$ , they cannot intersect. For two spaces  $S_{\nu-1}, S_{n-\nu-1}$  with a common point are contained in a space  $[n - 2]$ , and a prime cannot contain two directrix curves of  $F$ .

**280.** In any space  $[n - 1]$  we can at once construct a normal elliptic surface  $F$ , and indeed several types of surfaces. For take two spaces  $S_{m-1}$  and  $S_{n-m-1}$  which do not intersect; in  $S_{m-1}$  take a normal elliptic curve of order  $m$  and in  $S_{n-m-1}$  a normal elliptic curve of order  $n - m$ , these two curves having the same modulus. Then a (1, 1) correspondence between the two curves gives an elliptic ruled surface  $F$  of order  $n$ . If  $m$  is not equal to  $\frac{1}{2}n$  we can take the curve of order  $m$  to be the minimum directrix; clearly two surfaces  $F$  whose minimum directrices are not of the same order cannot be projectively equivalent, and two surfaces  $F$  whose minimum directrices are of the same order but have not the same modulus cannot be projectively equivalent.

If  $n$  is even and we choose  $m = \frac{1}{2}n$  the surface has two minimum directrices. In this case, since the two curves between which there is a (1, 1) correspondence are of the same order, we can obtain a special type of surface by making the correspondence belong to a projectivity\* between the two spaces  $S_{\frac{1}{2}n-1}$  and  $S'_{\frac{1}{2}n-1}$  containing the curves. Then, if a space  $[\frac{1}{2}n - 2]$  meets the first curve in  $\frac{1}{2}n$  points, the generators through these points meet the second curve in  $\frac{1}{2}n$  points which also lie in a space  $[\frac{1}{2}n - 2]$ ; hence these  $\frac{1}{2}n$  generators are contained in a space  $[n - 3]$ . Then, through this  $[n - 3]$ , we have a pencil of primes cutting out  $\infty^1$  curves of order  $\frac{1}{2}n$  on  $F$ ; hence this particular surface has  $\infty^1$  *minimum directrices of order  $\frac{1}{2}n$ , and through any point of  $F$  there passes one of them.*

Let us then divide the surfaces  $F$  into three classes, according to the order  $m$  of their minimum directrices:

$$(a) \quad m < \frac{n}{2},$$

$$(b) \quad m = \frac{n}{2},$$

$$(c) \quad m > \frac{n}{2}.$$

\* In other words, the (1, 1) correspondence between the two elliptic curves is *special*. Cf. § 164.

We prove that all surfaces of class (a) are generated by a (1, 1) correspondence between two elliptic normal curves of orders  $m$  and  $n - m$  such as we have already mentioned. We prove also that the surfaces of class (b) are those already considered, one with two minimum directrices and the other with  $\infty^1$  minimum directrices; as well as a third type of surface with only one minimum directrix. We prove further that in class (c) we have only one type of surface—that for which  $n$  is odd and  $m = \frac{1}{2}(n + 1)$ ; this being the most general type of surface of odd order  $n$ . For all surfaces  $F$ ,  $m < \frac{1}{2}(n + 1)$ .

These are the only possible types of elliptic normal ruled surfaces for which no two generators intersect.

**281.** Let us consider now a surface of the class (a); there is on the surface a minimum directrix  $\gamma^m$ , where  $m < \frac{1}{2}n$ . There can be no curve on the surface of order less than  $n - m$ ; we prove that the surface contains an infinity of curves of order  $n - m$ .

If there is on  $F$  a curve  $C_{n-m}$  then any prime containing it must meet  $F$  further in  $m$  generators, and these will meet  $\gamma^m$  in  $m$  associated points\*. Conversely, any set of  $m$  associated points of  $\gamma^m$  is contained in an  $[m - 2]$ , and any curve  $C_{n-m}$  is contained in an  $[n - m - 1]$ ; hence there is a prime containing any  $C_{n-m}$  and any set of  $m$  generators which meet  $\gamma^m$  in associated points. Hence, in order to obtain all the curves  $C_{n-m}$  on  $F$ , it will be sufficient to consider the system of primes which contain the generators of  $F$  passing through any given set of associated points of  $\gamma^m$ .

Let us take a set of associated points of  $\gamma^m$ , they lie in an  $[m - 2]$ ; thus the  $m$  generators which pass through them belong to a  $[2m - 2]$ . They cannot belong to a space  $[2m - 3]$ ; for we should then have a  $[2m - 2]$  containing  $\gamma^m$  and these  $m$  generators, i.e. a composite curve of  $F$  of order  $2m$  lying in a  $[2m - 2]$ , and this we have seen to be impossible if  $2m < n$ . Hence, through this set of  $m$  fixed generators of  $F$  there passes a linear system of  $\infty^{n-2m}$  primes, so that we have on  $F$  a linear system of  $\infty^{n-2m}$  curves  $C_{n-m}$ ; through  $n - 2m$  points of general position on  $F$  there passes one of the curves  $C_{n-m}$ .

To make this statement quite accurate we must verify that the curve  $C_{n-m}$ , which is the remaining part of the intersection of  $F$  with a prime containing the  $m$  fixed generators, does not break up. If it does break up it must consist of  $\gamma^m$  and  $n - 2m$  generators, so that we have a prime containing  $\gamma^m$ , the  $m$  fixed generators, and  $n - 2m$  other generators. But  $\gamma^m$ , the  $m$  fixed generators, and  $n - 2m - 1$  other generators arbitrarily chosen, determine a prime which contains them; hence the remaining

\* When we have an elliptic normal curve of order  $s$  in  $[s - 1]$  then the set of  $s$  points in which it is met by any prime  $[s - 2]$  is called a set of *associated points*.

generator in which the prime meets  $F$  is determined by the choice of the  $n - 2m - 1$  arbitrary generators. This proves that the prime which contains the  $m$  fixed generators and  $n - 2m$  arbitrary points of  $F$  meets  $F$  in a curve  $C_{n-m}$  which does not break up. Through  $n - 2m - 1$  arbitrary points of  $F$  there pass  $\infty^1$  curves  $C_{n-m}$  forming a pencil, all these curves having also one other point in common.

The surface  $F$  can be generated by a (1, 1) correspondence between  $\gamma^m$  and any one of the curves  $C_{n-m}$ .

If, in particular,  $m = \frac{1}{2}(n - 1)$ , we have a surface of odd order with a single minimum directrix of order  $\frac{1}{2}(n - 1)$ ; there are on this surface  $\infty^1$  curves of order  $\frac{1}{2}(n + 1)$ , which all pass through the same point.

282. Consider now surfaces  $F$  of the class (b); the surface is of even order  $n$ , and there is on it a minimum directrix  $\gamma^m$  of order  $m = \frac{1}{2}n$ . We enquire whether there are other directrices of this same order. Just as for surfaces of the class (a) such a curve is given by a prime containing a set of  $m$  fixed generators which meet  $\gamma^m$  in a set of  $m$  associated points.

Such a set of generators lies, as before, in a space  $[2m - 2]$ ; but we cannot now assert that they do not lie in a  $[2m - 3]$ . However, they certainly cannot lie in a  $[2m - 4]$ ; for then we should have a  $[2m - 3]$  or  $[n - 3]$  containing the composite curve of order  $n$  formed by these generators and  $\gamma^m$ .

If the  $m$  generators do lie in an  $[n - 3]$  then this  $[n - 3]$  is the base of a pencil of primes; these primes give on  $F$   $\infty^1$  curves  $C_m$  of order  $\frac{1}{2}n$ , one such curve passing through any given point of  $F$ . A set of generators which meets any one of these curves in associated points meets any other  $C_m$  in associated points also; the surface is that generated by a special (1, 1) correspondence between two normal elliptic curves of the same order.

In general the  $m$  generators do not lie in an  $[n - 3]$ ; they lie then in a prime, which meets  $F$  again in a curve  $C_m$  of order  $\frac{1}{2}n$ . The surface has then two minimum directrices; a prime through either of them meets  $F$  further in  $m$  generators, these generators meeting the other minimum directrix in a set of associated points. This is the most general surface of even order  $n$ , and is generated by a (1, 1) correspondence between two elliptic normal curves of the same order. This correspondence is *not* special.

When we take a set of  $m$  generators through associated points of  $\gamma^m$  it may happen that the prime which they determine contains  $\gamma^m$  itself. If this happens for one set of associated points of  $\gamma^m$  it must happen for all sets, and we have a surface  $F$  with one minimum directrix of order  $m = \frac{1}{2}n$ . Any  $m - 1$  generators of  $F$  determine an  $[n - 3]$ ; through this there pass the primes of a pencil, giving thus  $\infty^1$  curves  $C_{\frac{1}{2}n+1}$  on  $F$ . Each of these curves meets  $\gamma^m$  in one point; for the prime meets  $\gamma^m$  in  $m$  points of which  $m - 1$  are on the generators that we have chosen, the remaining

one being on  $C_{\frac{1}{2}n+1}$ . The prime containing  $\gamma^m$  and the generators through a set of associated points of  $\gamma^m$  meets a curve  $C_{\frac{1}{2}n+1}$  in  $m+1$  associated points, these including its intersection with  $\gamma^m$ . Hence we have the generation of a surface of this type. We take in  $[n-1]$  two spaces  $[\frac{1}{2}n-1]$  and  $[\frac{1}{2}n]$ ; these have a common point  $O$ . In  $[\frac{1}{2}n-1]$  we take a normal elliptic curve of order  $\frac{1}{2}n$  and in  $[\frac{1}{2}n]$  we take a normal elliptic curve of order  $\frac{1}{2}n+1$ ; these curves must have the same modulus and must both pass through  $O$ . We then place the curves in (1, 1) correspondence with  $O$  as a united point. The correspondence must be such that any set of  $\frac{1}{2}n$  points of the second curve, which correspond to a set of  $\frac{1}{2}n$  associated points of the first curve, forms, together with  $O$ , a set of  $\frac{1}{2}n+1$  associated points.

**283.** Just as for rational ruled surfaces we investigated the curves of lowest order on the surface by considering primes which contained the greatest possible number of generators, so we can proceed for the elliptic ruled surfaces.

If  $n$  is even we take  $\frac{1}{2}n-1$  generators; these determine an  $[n-3]$  which is the base of a pencil of primes. We have thus on the surface  $\infty^1$  curves of order  $\frac{1}{2}n+1$ ; any two of these curves have two intersections, and these two intersections must be in  $[n-3]$ . All the  $\infty^1$  curves of order  $\frac{1}{2}n+1$  pass through these two points, and the prime containing  $[n-3]$  and the generator through either of these points meets the surface further in a curve of order  $\frac{1}{2}n$ . Hence, if  $m$  is the order of the minimum directrix on a surface of even order  $n$ ,  $m < \frac{1}{2}n$ .

Similarly, if  $n$  is odd we take  $\frac{1}{2}(n-1)$  generators, so determining a prime  $[n-2]$ . This meets the surface further in a curve of order  $\frac{1}{2}(n+1)$ . Hence, if  $m$  is the order of the minimum directrix on a surface of odd order  $n$ ,  $m < \frac{1}{2}(n+1)$ .

This proves that the only surfaces belonging to the class (c) are those for which  $n$  is odd and  $m = \frac{1}{2}(n+1)$ .

**284.** If we consider now a surface belonging to the class (c), for which  $m = \frac{1}{2}(n+1)$ , any curve  $C_m$  on it belongs to a space  $[m-1]$ . Such a curve determines, together with any  $m-2$  fixed generators of  $F$ , a prime  $[2m-3]$ . This prime meets  $F$  in one other generator; hence, in order to obtain all the curves  $C_m$  on  $F$ , it is sufficient to consider the sections of  $F$  by primes through  $m-2$  fixed generators and one variable generator.

Now  $m-2$  fixed generators of  $F$  belong to a  $[2m-5]$ ; they cannot be contained in a  $[2m-6]$ , for then they would lie, together with a  $C_m$ , in the same  $[2m-6+m-1-(m-3)]$  or  $[n-3]$ , and no simple or composite curve of  $F$  of order  $m-2+m=n-1$  can lie in an  $[n-3]$ . Moreover, the  $[2m-5]$  containing these  $m-2$  fixed generators does not contain any other generators.



This space  $[2m - 5]$  and any other generator of  $F$  determine a prime  $[2m - 3]$ , and this prime meets  $F$  in a curve  $C_m$ . We have thus on  $F \infty^1$  curves  $C_m$ ; since, from the way in which we have obtained them, they are in (1, 1) correspondence with the generators of  $F$ , they form an elliptic family.

Any two of the curves  $C_m$  have one intersection. For consider the prime section of  $F$  formed by one curve  $C_m$ , the  $m - 2$  fixed generators, and one other generator. Any other curve of order  $m$  meets this prime in  $m$  points, of which  $m - 2$  are on the fixed generators and one on the other generator; the remaining intersection is then on  $C_m$ .

Hence, to generate this most general surface  $F$  of odd order, we take two elliptic curves of the same order  $m$  and the same modulus, and place them in (1, 1) correspondence with a united point.

If this correspondence is special, then a set of  $m - 1$  points on one of the curves which forms, with the united point, an associated set of points gives, on the other curve, a corresponding set of  $m - 1$  points which also forms, with the united point, an associated set of points. Thus the  $m - 1$  generators lie in a  $[2m - 4]$  or  $[n - 3]$ . Then the pencil of primes through  $[n - 3]$  gives on  $F \infty^1$  curves of order  $m$ , and these must all meet  $[n - 3]$  in the same point, since they all meet it in one point which is not on any of the  $m - 1$  generators. The generator through this point lies in a prime with  $[n - 3]$ , and this prime gives a curve of order  $m - 1$  on  $F$ . Hence, when we generate the most general surface of odd order  $2m - 1$  in the way described, the correspondence between the two curves must *not* be special.

**285.** If we take  $k$  generators of the elliptic normal surface  $F$  in  $[n - 1]$  then, to whatever class the surface may belong, a space  $[n - 3]$  containing these generators meets  $F$ , in general, in these generators and a number of isolated points. In order that this may be true it is necessary that the space to which the  $k$  generators belong should not contain a directrix curve of  $F$ , and this is certainly always true when  $k < m$ , the order of the minimum directrix. A prime through  $[n - 3]$  meets  $F$  in a curve of order  $n - k$ , and, of the  $n - k$  intersections of this curve with  $[n - 3]$ ,  $k$  lie on the generators, while the remaining  $n - 2k$  give isolated intersections of  $[n - 3]$  with  $F$ . Thus, if  $k < m$ , any  $S_{n-3}$  containing  $k$  generators of  $F$  meets  $F$  again in  $n - 2k$  isolated points.

**286.** In order to find, for the surface of class (c), how many curves  $C_m$  pass through a given point  $P$  of  $F$ , we have to obtain those primes which contain  $P$  and  $m - 2$  fixed generators, and which also contain another generator of  $F$ . Now the  $m - 2$  fixed generators and  $P$  determine a  $[2m - 4]$  or  $[n - 3]$ , and this, as we have just proved, meets  $F$  in two other points  $Q$  and  $Q'$ . Then the prime which contains  $[n - 3]$  and the generator through  $Q$  meets  $F$  in a  $C_m$  passing through  $P$ , and we have a second  $C_m$  passing

through  $P$  if we take the prime which contains  $[n-3]$  and the generator through  $Q'$ . Hence *through any point of  $F$  there pass two of the minimum directrices.*

The following result can also be obtained by arguments similar to those just used. *An elliptic ruled surface of order  $n$  with a minimum directrix of order  $m$  contains  $\infty^{2\mu-n}$  elliptic curves of order  $\mu$ , for all values of  $\mu$  for which  $n > \mu > n - m$ . Through  $2\mu - n$  points of general position on the surface there pass two of the curves. The only exception is for the surface of class (b) with  $\infty^1$  minimum directrices, when  $\mu = \frac{1}{2}n + 1$ .*

**287.** Suppose now that we have in  $[n-1]$  a normal elliptic ruled surface  $F$  of order  $n$ , on which two generators intersect. If we project from their point of intersection on to an  $[n-2]$  we obtain an elliptic ruled surface of order  $n-2$ , which is therefore a cone. Hence on  $F$  we have a line passing through the intersection of the two generators and meeting all the other generators. Since the surface  $F$  is not rational this line  $\lambda$  must be a *double line*. Through every point of  $\lambda$  there must pass two generators of  $F$ ; for otherwise  $\lambda$  would be a directrix and also a generator, one variable generator passing through each point of  $\lambda$ , or else  $\lambda$  would be a double generator. In the first case  $F$  would be rational and in the second case a prime section of  $F$  would be a curve  $C_{n-1}$  in  $[n-2]$ , the curve having a double point, which means again that  $F$  is rational. We have then a surface  $F$  with a double line  $\lambda$ ;  $\lambda$  is a double directrix, through each point of it there pass two generators of  $F$ .

Such a surface cannot contain a directrix curve of order  $\nu < n-2$ ; for, since such a curve is contained in  $[\nu-1]$ , the  $[\nu+1]$  containing this curve and  $\lambda$  would contain the whole of  $F$ . But there is an infinity of curves of order  $n-2$  on  $F$ ; such curves are all obtained by means of primes through two fixed intersecting generators, so that there are  $\infty^{n-4}$  such curves. Thus the surface  $F$  is generated by a (1, 2) correspondence between a line  $\lambda$  and an elliptic normal curve of order  $n-2$ . The curves of order  $n-2$  on  $F$  form a linear system; through  $n-4$  points of general position on  $F$  there passes *one* such curve. But any two of the curves intersect in  $n-4$  points, and through such a set of  $n-4$  points there passes a pencil of the curves.

### *The normal surfaces in [5] with directrix cubic curves*

**288.** We now proceed to study the elliptic sextic ruled surfaces in [3] as projections of normal surfaces in higher space. All elliptic sextic ruled surfaces are projections of normal surfaces in [5]. We project the normal surface  $F$  from a line  $l$  which does not meet it on to a [3]  $\Sigma$ , and we thus obtain one of the surfaces  $f$ . The planes joining  $l$  to the points of  $F$  meet  $\Sigma$  in the points of  $f$ , while the solids joining  $l$  to the generators of  $F$  meet  $\Sigma$  in the generators of  $f$ .

The general surface  $F$  has two elliptic plane cubic curves  $\Gamma_1$  and  $\Gamma_2$  on it and can be generated by means of a non-special (1, 1) correspondence between  $\Gamma_1$  and  $\Gamma_2^*$ .

The tangents of  $F$  form a four-dimensional locus; this can be regarded as the locus of  $\infty^3$  lines,  $\infty^2$  planes or  $\infty^1$  solids†; these  $\infty^1$  solids are, in fact, determined by the pairs of tangents of  $\Gamma_1$  and  $\Gamma_2$  at corresponding points. Thus any two tangent solids of  $F$  have a line in common meeting the planes of  $\Gamma_1$  and  $\Gamma_2$ .

We can set up a correspondence between the points  $B$  and  $C$  of  $\Gamma_2$  by saying that  $B$  and  $C$  shall correspond when the solid which is determined by  $l$  and the tangent to  $\Gamma_1$  at the point  $A$  on the generator through  $B$  meets the plane of  $\Gamma_2$  on the tangent at  $C$ . In this way we have a (6, 6) correspondence of valency zero, the six points  $C$  which correspond to a given point  $B$  lying on a conic. Now if a point  $B$  coincides with one of its corresponding points  $C$  there is a [4] through  $l$  containing the tangents of  $\Gamma_1$  and  $\Gamma_2$  at  $A$  and  $B$ , and conversely. When this is so the solid containing the tangents to  $\Gamma_1$  and  $\Gamma_2$  at a pair of corresponding points meets  $l$ . There are therefore twelve such solids; or the locus of tangents of  $F$  is an  $M_4^{12}$ .

289. Through a general point  $P$  of the space [5] there passes a unique line meeting the planes of  $\Gamma_1$  and  $\Gamma_2$ ‡; let the points of meeting be  $O_1$  and  $O_2$ . There is a finite number of chords of  $F$  passing through  $P$ ||; let one of these meet  $F$  in  $X$  and  $Y$ .

The solid through  $O_1O_2P$  and the generator of  $F$  which passes through  $X$  meets  $F$  further in at least five points, namely,  $Y$ , two points of  $\Gamma_1$  and two points of  $\Gamma_2$ . We therefore conclude¶ that this solid contains also the generator of  $F$  which passes through  $Y$ . The lines through  $O_2$  in the plane of  $\Gamma_2$  cut out a  $g_3^1$  on that curve; the generators of  $F$  through a set of this  $g_3^1$  meet  $\Gamma_1$  in three points forming also a set of a  $g_3^1$ ; and corresponding to each chord of  $F$  passing through  $P$  there are two of the three points of a set collinear with  $O_1$ . Now the lines through  $O_1$  in the plane of  $\Gamma_1$  cut out a second  $g_3^1$  on that curve, and there are just three pairs of points on  $\Gamma_1$  which are common to two linear series  $g_3^1$ \*\*.

Hence *there are three chords of  $F$  passing through a general point  $P$  of the space* [5].

Through any two points  $X, Y$  of  $F$  there pass two directrix quartic

\* § 282.

† § 51.

‡ This line is the intersection of the solid containing  $O$  and  $\Gamma_1$  with the solid containing  $O$  and  $\Gamma_2$ .

|| Cf. § 52.

¶ § 285.

\*\* Cf. the footnote to § 134.

curves  $Q_1$  and  $Q_2^*$ . Through any point  $P$  of  $XY$  there passes a second chord  $PX_1Y_1$  of  $Q_1$  and a second chord  $PX_2Y_2$  of  $Q_2$ ; thus the existence of one chord of  $F$  through  $P$  at once implies the existence of three. The four points  $X_1, Y_1, X_2, Y_2$  lie on another directrix quartic curve.

**290.** *Elliptic sextic ruled surfaces in [4].* If we project from  $P$  on to a [4] we see that a general elliptic sextic ruled surface in [4] has three double points; it contains two plane cubic curves and three plane quartic curves. Each of the plane quartics has double points at two of the double points of the surface.

If  $P$ , instead of occupying a general position, is at the vertex of a quadric cone which contains a directrix quartic  $Q$  (there are four such positions of  $P$  possible for each directrix quartic, and the locus of such positions will be a surface) we obtain in [4] an elliptic sextic ruled surface with a double conic. It can be obtained by taking two planes in [4] and then a conic in one of the planes and an elliptic cubic curve in the other, both these curves passing through the intersection of the planes; a (1, 2) correspondence between the conic and the cubic with a united point gives the surface.

If  $P$  is taken to lie in the plane of  $\Gamma_1$  or the plane of  $\Gamma_2$  we obtain in [4] an elliptic sextic ruled surface with a triple line. The surface is given by a (1, 3) correspondence between a line and an elliptic plane cubic, the line not meeting the plane of the cubic curve.

**291.** We have seen that there are three chords of  $F$  passing through a general point of [5]. If the point is such that two of the three chords coincide it will have to lie on some primal  $V_4$ . Instead of having three chords  $PXY, PX_1Y_1$  and  $PX_2Y_2$  we shall have only two chords  $PXY$  and  $PX_0Y_0$ , and one of the two directrix quartics through  $X_0$  and  $Y_0$  is such that its tangents at  $X_0$  and  $Y_0$  intersect. Conversely, if we take two points  $X_0$  and  $Y_0$  on a directrix quartic of  $F$  such that the tangents of the quartic at these two points intersect †, then, if any point is taken on  $X_0Y_0$ , there is no other chord of the quartic passing through it, the projection of the quartic from this point on to a plane having a tacnode. There is another directrix quartic of  $F$  passing through  $X_0$  and  $Y_0$  and there will be a second chord of this second quartic passing through any point on  $X_0Y_0$ . Thus any point on  $X_0Y_0$  is such that only two chords of  $F$  pass through it.  $V_4$  can therefore be defined as the locus of chords of  $F$  such as  $X_0Y_0$ .

We shall now examine some of the properties of  $V_4$  which will be of use in the sequel, and this affords an opportunity of introducing an idea

\* § 286.

† Each tangent of the quartic is met by four others.

of which we have not availed ourselves before—namely, we can regard the normal surface  $F$  as *the projection of another normal surface, in space of one higher dimension, from a point of itself*; the order of this other normal surface being one higher than that of  $F$ .

292. Take a general normal elliptic ruled surface  $F_0$  of the seventh order in [6]. There are on this surface  $\infty^1$  elliptic quartic curves; through any point of  $F_0$  there pass two of these curves, while any two of the curves have one intersection\*. The chords of  $F_0$  form in [6] a primal  $M_5$  whose order is the number of its intersection with an arbitrary line. But the projection of  $F_0$  from this line is an elliptic ruled surface of the seventh order in [4] which is known, by a general result †, to have seven double points. Hence the chords of  $F_0$  form an  $M_5^7$ . Now the projection of  $F_0$  on to [4] from a line meeting it is an elliptic sextic ruled surface, which we have seen to have three double points. Hence such a line meets three chords of  $F_0$  in points other than that in which it meets  $F_0$ . Thus  $F_0$  is a quadruple surface on  $M_5^7$ .

A general chord of  $F_0$  does not meet any others, because the projection of  $F_0$  from the chord is an elliptic quintic ruled surface in [4] which has no double points. But if the chord of  $F_0$  meets it in two points which are on the same directrix quartic there passes a second chord of  $F_0$  through every point of the first chord; the projection from the chord now gives the elliptic quintic ruled surface in [4] with a double line ‡. Thus through any point which lies in a solid containing a directrix quartic of  $F_0$  there pass two chords of  $F_0$ ; these solids, of which there are  $\infty^1$ , form a locus  $M_4$  which is a double locus on  $M_5^7$ . It can be shewn that if there is any point, not on  $F_0$  itself, through which there pass two chords of  $F_0$ , then the intersections of these chords with  $F_0$  are four points lying on the same directrix quartic of  $F_0$ . Thus  $M_4$  is the only double locus on  $M_5^7$ .

To find the order of  $M_4$  we take a prime section; this gives the quadrisecant planes of an elliptic curve  $C_7^1$  of the seventh order in [5]. There are

\* §§ 284, 286.

† A ruled surface of order  $n$  and genus  $p$  in [4] has  $\frac{1}{2}(n-2)(n-3) - 3p$  double points. See Tanturri, *Atti Torino*, 35 (1900), 441. Some of our former work gives particular cases of this general result. For  $n = 4$  and  $p = 0$  we see that a rational ruled quartic surface in [4] has one double point and therefore that there is one chord of a rational normal quartic ruled surface passing through a general point of [5] (§ 81). For  $n = 5$  and  $p = 0$  the rational quintic ruled surface in [4] has three double points and the chords of the normal surface in [6] form an  $M_5^3$  (§ 124). For  $n = 5$  and  $p = 1$  the elliptic quintic ruled surface in [4] has no double points (§ 160). For  $n = 6$  and  $p = 0$  the rational sextic ruled surface in [4] has six double points and the chords of the normal surface form an  $M_5^6$  (§ 174). For  $n = 6$  and  $p = 1$  the elliptic sextic ruled surface in [4] has three double points and there are three chords of the normal surface through a point of [5] (§ 289).

‡ § 164.

$\infty^1$  such planes, two passing through each point of the curve\*; they form an  $M_3$  which is a prime section of  $M_4$ , and the order of  $M_3$  is simply the number of quadrisecant planes of  $C_7^1$  which meet a general plane of [5]. But this number is 7, as is seen on making use of a general result due to Giambelli†. Thus we have an  $M_3^7$  on which  $C_7^1$  is a double curve, and so the solids containing the directrix quartics of  $F_0$  form an  $M_4^7$  on which  $F_0$  is a double surface.

Now let us project  $F_0$  from a point  $O$  of itself on to [5]; we obtain the normal elliptic sextic ruled surface  $F$ .

A plane through  $O$  meets  $M_4^7$  in five points other than  $O$ ; it meets  $M_5^7$  in a curve of the seventh order with a quadruple point at  $O$  and these five points for double points; this curve of the seventh order will then be of genus 4. The plane meets [5] in a line, and the fact that any line through  $O$  in the plane meets the curve again in three points corresponds to the fact that there are three chords of  $F$  passing through each point of the line in which the plane meets [5]. Now the lines through  $O$  give a pencil of  $\infty^1$  sets of three points upon a curve of genus 4; thus there are, by a result originally due to Riemann,  $2(3+4-1)$  or twelve positions of the line for which two of the three points coincide; which is the same as saying that twelve tangents can be drawn to the curve from the quadruple point. Hence, given any line in [5] there are twelve points on it such that two of the three chords of  $F$ , which pass through any one of them, coincide.

Hence the locus  $V_4$  of such points is of order 12; we denote it by  $V_4^{12}$ . It must not be confused with  $M_4^{12}$ , the locus of tangents of  $F$ .

**293.** There are two directrix quartics of  $F_0$  passing through  $O$ ; these give on projection the cubics  $\Gamma_1$  and  $\Gamma_2$  on  $F$ . The directrix quartics of  $F_0$  which do not pass through  $O$  give  $\infty^1$  of the directrix quartics on  $F$ ; there are  $\infty^2$  elliptic quintic curves on  $F_0$  which pass through  $O$  and these give on projection  $\infty^2$  elliptic quartic curves on  $F$ .

Consider now a point  $U$  of [5] which is the vertex of a quadric cone containing a directrix quartic of  $F$ . In general, this quartic is the projection of a directrix quintic of  $F_0$  which passes through  $O$ ; then the line  $OU$  meets  $\infty^1$  chords of this quintic and so lies on  $M_5^7$ . It meets  $M_4^7$  in one point other than  $O$ ‡. Then a plane through  $OU$  meets  $M_5^7$  in  $OU$

\* For the projection of  $C_7^1$  from a point of itself on to a [4] is an elliptic sextic curve having two trisecants (cf. § 253).

† *Memorie Torino* (2), 59 (1909), 489.

‡ Each quartic on  $F_0$  meets the quintic in two points; the [3] containing the quartic meets the [4] containing the quintic in the line joining the two points. We have thus on the quintic a symmetrical (2, 2) correspondence; two points of the quintic corresponding when they lie on the same directrix quartic of  $F_0$ . The chords of the quintic meeting  $OU$  generate a cubic ruled surface and give a symmetrical (1, 1) correspondence. Both these symmetrical correspondences are of valency one and they have *one* common pair of points, the quintic being elliptic.

and a sextic curve; this sextic has a triple point at  $O$  and double points at those four points in which the plane meets  $M_4^7$  and which themselves do not lie on  $OU$ . It is therefore of genus 3 and ten tangents can be drawn to it from the triple point. Hence, if in [5] we take a line  $l$  which passes through the vertex  $U$  of a quadric cone containing a directrix quartic of  $F$ , the line will only meet  $V_4^{12}$  in ten points other than  $U$ .

A similar argument can be used when the quartic on  $F$  is the projection of a quartic on  $F_0$  which does not pass through  $O$ ; the same result is obtained.

Thus the vertices of the quadric cones which contain the directrix quartics of  $F$  are *double points* on  $V_4^{12}$ ; the locus of the vertices will be a *double surface* on  $V_4^{12}$ .

The point  $U$  is a *quadruple point* of  $M_4^{12}$ , since there are *four* tangents of  $F$  passing through it.

Take now a line through  $O$  lying in the solid containing one of the two directrix quartics of  $F_0$  which pass through  $O$ . This line lies on  $M_4^7$  and any plane through the line will meet  $M_4^7$  in four further points\*. This same plane meets  $M_5^7$  in the line, counted doubly, and a quintic curve; this quintic curve has a double point at  $O$  and four other double points. It is therefore elliptic, and six tangents can be drawn to it from its double point. Hence, if we take a line  $l$  meeting the plane of one of the cubic curves on  $F$  it will only meet  $V_4^{12}$  in six points other than its intersection with the plane; or the planes of  $\Gamma_1$  and  $\Gamma_2$  are *sextuple planes* on  $V_4^{12}$ .

The planes of  $\Gamma_1$  and  $\Gamma_2$  are also sextuple planes on  $M_4^{12}$ ; through a general point of either plane there pass six tangents of  $F$ .

Through any point of  $V_4^{12}$  there pass two chords  $XY$  and  $X_0Y_0$  of  $F$ . The two quartic curves of  $F$  which pass through  $X$  and  $Y$  coincide, while one of those through  $X_0$  and  $Y_0$  is such that its tangents at  $X_0$  and  $Y_0$  intersect; the other quartic through  $X_0$  and  $Y_0$  is the same as the one through  $X$  and  $Y$ . Now consider a line  $l$  which lies in the [3] determined by two generators  $g$  and  $g'$  of  $F$ ; there are twelve points of  $V_4^{12}$  on  $l$  and through each of these there passes a transversal to  $g$  and  $g'$ . We now enquire how many of these twelve transversals give a pair of points  $X$  and  $Y$  on  $g$  and  $g'$ ; the remaining ones will give pairs of points  $X_0$  and  $Y_0$ .

Consider the quartics of  $F$  passing through a point of  $g$ ; there are  $\infty^1$  of them and any two intersect in one further point. They thus form an elliptic family, since each curve may be put into correspondence with its second intersection with a fixed curve of the family. Now we can regard this family as  $\infty^1$  pairs of curves, since if any point is taken on  $g'$  there are two and only two curves of the family passing through it. Thus these

\* A solid through  $O$  meets  $M_4^7$  in an elliptic curve of the seventh order with a double point at  $O$ ; but if the solid contains a line through  $O$  on  $M_4^7$  it meets it further in an elliptic sextic curve which meets the line in  $O$  and in one other point.

pairs, associated with the points of  $g'$ , form a rational family; there are therefore four double elements, just as a  $g_2^1$  on an elliptic curve has four double points. Hence, given a point  $X$  on  $g$  there are four points  $Y$  on  $g'$  such that the two quartics of  $F$  which pass through  $X$  and  $Y$  coincide.

Now we can set up a correspondence between the points  $P$  and  $Q$  of  $g'$ , saying that the points  $P$  and  $Q$  correspond when the two directrix quartics through  $P$  and  $Q'$  coincide,  $Q'$  being the intersection of  $g$  with the transversal to  $g$  and  $l$  from  $Q$ . We have thus a (4, 4) correspondence, and therefore eight coincidences. Hence there are eight pairs of points  $X$  and  $Y$  on  $g$  and  $g'$  such that  $XY$  meets  $l$  and the two directrix quartics of  $F$  which pass through  $X$  and  $Y$  coincide.

This argument must be slightly modified if  $l$  meets the plane of one of the curves  $\Gamma_1, \Gamma_2$ . Suppose that  $l$  meets the plane of  $\Gamma_1$  in a point  $O$ , we have a line through  $O$  meeting  $g$  in a point  $A$  and  $g'$  in a point  $A'$ ,  $A$  and  $A'$  being the intersections of  $\Gamma_1$  with  $g$  and  $g'$ . Then through  $A$  and a given point of  $g'$  there passes only one directrix quartic of  $F$ , the other being the degenerate one consisting of  $\Gamma_1$  and  $g'$ . Hence the four points of  $g'$ , which are such that the two directrix quartics which are determined by  $A$  and any one of them coincide, must all be at the point  $A'$ . In the correspondence between the points  $P$  and  $Q$  of  $g'$ , when  $P$  is at  $A'$  the four corresponding points  $Q$  are also at  $A'$ . Hence  $A'$  counts four times among the eight points of  $g'$  for which  $P$  coincides with one of its corresponding points  $Q$ . There are four further coincidences.

We have then the following: if a line  $l$  is taken to meet the plane of one of the directrix cubics of  $F$  in a point  $O$  and to lie in the solid containing a pair of generators  $g, g'$  of  $F$  (which will meet the cubic in a pair of points collinear with  $O$ )  $l$  will meet  $V_4^{12}$  in six points other than  $O$ ; if the transversals are drawn from these points to  $g$  and  $g'$ , four of them give pairs of points  $X, Y$  on  $g, g'$ , while the remaining two give pairs of points  $X_0, Y_0$ .

**294.** Take now a line  $l$  of general position in [5] and project  $F$  from  $l$  on to a [3]  $\Sigma$ . A prime through  $l$  meets  $F$  in an elliptic sextic curve of which nine chords meet  $l$ . Thus the chords of  $F$  which meet  $l$  meet  $F$  in a curve  $C_{18}$  of order 18. Since the solid containing  $l$  and an arbitrary generator of  $F$  meets  $F$  further in four points\*,  $C_{18}$  meets each generator in four points.

Thus after projection we obtain in  $\Sigma$  a general elliptic ruled surface  $f$  of the sixth order with a double curve  $C_9$  of the ninth order meeting each generator in four points.  $C_9$  is the projection of  $C_{18}$  from  $l$ .

Now there is a (3, 1) correspondence between the points of  $C_9$  and the points of  $l$ , two points corresponding when the line joining them is a chord



of  $F$ , and the twelve points in which  $l$  meets  $V_4^{12}$  are branch-points of the correspondence. Then Zeuthen's formula shews that  $C_9$  is a curve  $C_9^4$  of genus 4. Further, there is a (1, 2) correspondence between  $C_9^4$  and  $C_{18}$ , and the correspondence has twelve branch-points on  $C_9^4$ , these arising from the twelve intersections of  $l$  with  $M_4^{12}$  or the twelve points of  $l$  which lie on tangents of  $F$ . A second application of Zeuthen's formula now shews that  $C_{18}$  is a curve  $C_{18}^{13}$  of genus 13. Since  $C_{18}^{13}$  meets each generator of  $F$  in four points it is of genus  $19 - x$ , where  $x$  is the number of its double points\*. Thus  $C_{18}^{13}$  has six double points. Let  $A$  be one of these. Then there are two chords  $AB$  and  $AC$  of  $C_{18}^{13}$  passing through  $A$  and meeting  $l$ ; this forces the chord  $BC$  to meet  $l$  also, and  $B$  and  $C$  are double points of  $C_{18}^{13}$ . Thus the six double points lie in two planes passing through  $l$ , each plane containing three of them. On projecting we have two triple points on  $C_9^4$ .

The (1, 4) correspondence between  $\Gamma_1$  and  $C_{18}^{13}$  shews that there are twenty-four generators of  $F$  touching  $C_{18}^{13}$ ; hence there are also twenty-four generators of  $f$  touching  $C_9^4$  (cf. § 256).

The prime determined by  $l$  and the plane of  $\Gamma_1$  contains three generators of  $F$ , as also does that determined by  $l$  and the plane of  $\Gamma_2$ . Thus  $f$  has two tritangent planes.

$f$  is generated by two elliptic cubics in (1, 1) correspondence.

295. There are  $\infty^2$  directrix quartic curves on  $F$ , and through any two points of  $F$  there pass two of these. There are four quadric cones through any such curve, and we may specialise the position of  $l$  so that it passes through the vertex  $U$  of a quadric cone containing a directrix quartic  $C_4^1$ . The chords of  $F$  which meet  $l$  now meet  $F$  in the points of  $C_4^1$  and a curve  $C_{14}$  which meets every generator of  $F$  in three points; when we project from  $l$  on to  $\Sigma$  we obtain a surface  $f$  whose double curve is  $C_2 + C_7$ .

Since  $U$  is a double point on  $V_4^{12}$  there are ten branch-points of the (3, 1) correspondence between  $C_7$  and  $l$ ; hence  $C_7$  is a  $C_7^3$ . The (1, 2) correspondence between  $C_7^3$  and  $C_{14}$  only has eight branch-points since  $U$  is a quadruple point on  $V_4^{12}$ ; hence  $C_{14}$  is a  $C_{14}^9$ .

$C_{14}^9$  meets each generator in three points and is therefore of genus  $11 - x$ , where  $x$  is the number of its double points; it has therefore two double points  $A$  and  $B$ . Through  $A$  there pass two chords  $AA'$  and  $AA''$  of  $C_{14}^9$  both meeting  $l$ ;  $A'A''$  will then also meet  $l$ , and  $C_4^1$  passes through  $A'$  and  $A''$ . Similarly we have two intersections  $B'$  and  $B''$  of  $C_4^1$  and  $C_{14}^9$ . There are in all eight intersections of  $C_4^1$  and  $C_{14}^9$  †, the other four lie

\* § 17 *supra*.

† Take a [4] through  $C_4^1$ ; it meets  $C_{14}^9$  in fourteen points and  $F$  in  $C_4^1$  and two generators. Since  $C_{14}^9$  meets each of these generators three times it will have to meet  $C_4^1$  eight times.

in pairs on two lines through  $U$ . Hence, projecting on to  $\Sigma$ ,  $C_7^3$  has two double points and two simple points on  $C_2$  (cf. § 259).

The (1, 3) correspondence between  $\Gamma_1$  and  $C_{14}^9$  shews that there are sixteen generators of  $F$  touching  $C_{14}^9$ ; there must then be sixteen generators of  $f$  touching  $C_7^3$  (cf. § 257).

This surface is generated by a conic and an elliptic cubic in (1, 2) correspondence with a united point. The pairs of points of the cubic corresponding to the points of the conic form a  $g_2^1$  and are therefore collinear with a fixed point\* of the curve. Since one of these lines passes through the point of the conic which gives rise to it the planes joining the points of the conic to the pairs of corresponding points on the cubic touch a quadric cone; this is part of the bitangent developable of the surface.

**296.** We may further choose  $l$  to join the vertex  $U$  of a quadric cone containing a directrix quartic  $C_4^1$  to the vertex  $V$  of a quadric cone containing a directrix quartic  $D_4^1$ .  $C_4^1$  and  $D_4^1$  have two intersections  $X$  and  $Y$ . The chords of  $F$  which meet  $l$  meet  $F$  in  $C_4^1$ ,  $D_4^1$  and a curve  $C_{10}$  which meets each generator of  $F$  in two points. Thus on projection we have a surface  $f$  with a double curve  $C_2 + D_2 + C_5$ .

$U$  and  $V$  are both double points of  $V_4^{12}$ , so that  $l$  meets  $V_4^{12}$  in eight other points which are the branch-points of the (3, 1) correspondence between  $C_5$  and  $l$ . Hence  $C_5$  is a  $C_5^2$ . Since  $U$  and  $V$  are quadruple points of  $M_4^{12}$  there are four branch-points of the (1, 2) correspondence between  $C_5^2$  and  $C_{10}$  so that  $C_{10}$  is a  $C_{10}^5$ . There are eight generators of  $F$  touching  $C_{10}^5$  and eight generators of  $f$  touching  $C_5^2$ .

The curve  $C_{10}^5$ , meeting each generator in two points, has no double points. The lines  $XU$  and  $XV$  meet  $C_4^1$  and  $D_4^1$  again in points  $U'$  and  $V'$ , the line  $U'V'$  meets  $l$  and  $U'$  and  $V'$  must lie on  $C_{10}^5$ ; we have similarly  $U''$  on  $C_4^1$  and  $C_{10}^5$  and  $V''$  on  $D_4^1$  and  $C_{10}^5$  arising from  $Y$ . There are in all six intersections of  $C_{10}^5$  with  $C_4^1$  or  $D_4^1$ ; the remaining four will lie on two lines through  $U$  or  $V$  respectively. Thus, on projection,  $C_2$  and  $D_2$  have two intersections through which  $C_5^2$  passes;  $C_5^2$  has two other intersections with each conic (cf. § 261).

To generate this surface take two conics in (2, 2) correspondence with two united points  $P$  and  $Q$ ; to  $P$  regarded as a point of one conic correspond two points of the other conic one of which is  $P$ , and similarly for  $Q$ .

The lines joining the pairs of points of one conic which correspond to the points of the other touch a conic; the planes joining these lines to

\* If we use elliptic arguments for the points of the cubic curve it is known that a  $g_2^1$  is given by  $a + \beta \equiv c$ , where  $a$  and  $\beta$  are the arguments for a pair of points of the  $g_2^1$  and  $c$  is a constant; different values of  $c$  give  $\infty^1 g_2^1$ 's on the curve. If then we take the point of the curve whose elliptic argument is  $\gamma \equiv -c$  we have  $a + \beta + \gamma \equiv 0$  and each pair of the  $g_2^1$  is collinear with  $\gamma$ :

the points of the other conic from which they arise touch a quadric cone, two lines passing through the points which give rise to them. Thus this surface has two quadric cones belonging to its bitangent developable.

**297.** Take now  $l$  to lie in the solid determined by a pair of generators  $g, g'$  of  $F$ . A prime through  $l$  meets this solid in a plane containing a transversal of  $l, g, g'$ ; it meets  $F$  in an elliptic sextic curve which meets  $g$  and  $g'$  at their intersections with the transversal and which has eight other chords that meet  $l$ . Hence the chords of  $F$  which meet  $l$  meet  $F$  in  $g, g'$  and a curve  $C_{16}$  of order 16; this last curve meets each generator of  $F$  in four points.

The solid  $gg'l$  meets  $F$  further in two points  $X$  and  $Y$ . Through  $X$  there passes a transversal of  $g$  and  $l$  and also a transversal of  $g'$  and  $l$ , so that  $X$  is a double point of  $C_{16}$ . Similarly  $Y$  is a double point of  $C_{16}$ . Associated with  $X$  we have intersections of  $C_{16}$  with each of  $g$  and  $g'$ , namely, the points in which  $g$  and  $g'$  are met by the plane  $XL$ . Similarly the plane  $YL$  meets  $g$  and  $g'$  in points which lie on  $C_{16}$ . The remaining points of intersection of  $C_{16}$  with  $g$  and  $g'$ , two on each generator, lie on two transversals of  $l, g, g'$ . Hence when we project from  $l$  on to  $\Sigma$  we obtain a surface  $f$  with a double generator  $G$  and a double curve  $C_8, C_8$  having two double points and two simple points on  $G$  (cf. § 265).

There are twelve points on  $l$  for which two of the three chords of  $F$  which pass through them coincide; we have seen\* that for eight of the twelve points the two chords of  $C_{16}$  coincide, while for the remaining four points it is a chord of  $C_{16}$  which coincides with the transversal to  $g$  and  $g'$ . Thus there are eight branch-points of the (2, 1) correspondence between  $C_8$  and  $l$ , so that  $C_8$  is a  $C_8^3$ . There are twelve tangents of  $C_{16}$  meeting  $l$  so that the (1, 2) correspondence between  $C_8^3$  and  $C_{16}$  has twelve branch-points and  $C_{16}$  is a  $C_{16}^{11}$ . It has only the two double points  $X$  and  $Y$ .

There are twenty generators of  $F$  touching  $C_{16}^{11}$  as is seen at once by the application of Zeuthen's formula to the (1, 4) correspondence between  $\Gamma_1$  and  $C_{16}^{11}$ ; there will then be twenty generators of  $f$  touching  $C_8^3$  (cf. § 265).

$f$  is generated by two elliptic plane cubics in (1, 1) correspondence, the correspondence being specialised to give the double generator.

**298.** Take the two generators  $g$  and  $g'$  of  $F$ ; the solid which they determine meets  $F$  in two further points  $X$  and  $Y$ . Take  $C_4^1$ , one of the two directrix quartics through  $X$  and  $Y$ , meeting  $g$  in  $Z$  and  $g'$  in  $Z'$ . Then the four points  $XYZZ'$  must be coplanar, since the solid  $gg'$  cannot contain  $C_4^1$ .

\* § 293.

Through  $X$  there passes a transversal of  $g$  and  $g'$ ; but the only lines which are trisecant to  $F$  are those lying in the planes of its directrix cubics, hence we may say that  $X$  is on  $\Gamma_1$  and  $Y$  on  $\Gamma_2$ .

The lines  $XZ$  and  $YZ'$  meet in a point  $U$ ; by suitably selecting the points where  $g$  and  $g'$  meet  $\Gamma_1$  we can cause  $U$  to be the vertex of a quadric cone through  $C_4^1$ .

This being done, take a line  $l$  through  $U$  which lies in the solid  $gg'$ . The chords of  $F$  which meet  $l$  meet  $F$  in  $g, g', C_4^1$  and a curve  $C_{12}$  meeting each generator of  $F$  in three points. Through  $X$  there passes a transversal of  $g'$  and  $l$  meeting  $g'$  in  $X'$  say, and the lines  $XZ$  and  $X'Z$  will meet  $l$ ;  $C_{12}$  must pass through  $X'$  and  $X$ . Similarly, we have a point  $Y'$  on  $g$ ;  $C_{12}$  will pass through  $Y'$  and  $Y$ .

Thus on projecting from  $l$  we have in  $\Sigma$  a surface  $f$  whose double curve consists of a conic, a double generator meeting the conic in two points and a sextic curve  $C_6$  passing through these two points (cf. § 266).  $C_6$  meets the double generator in two further points, these are projections of pairs of intersections of  $C_{12}$  with  $g$  and  $g'$ .

There are ten points of  $V_4^{12}$  on  $l$ , other than  $U$ ; six of these give rise to coincidences of the two chords of  $C_{12}$  which pass through them, while the remaining four give a coincidence of a chord of  $C_{12}$  with a transversal of  $g$  and  $g'$ . Thus the (2, 1) correspondence between  $C_6$  and  $l$  has six branch-points, so that  $C_6$  is a  $C_6^2$ . There are eight tangents of  $C_{12}$  meeting  $l$  so that the (1, 2) correspondence between  $C_6^2$  and  $C_{12}$  has eight branch-points and  $C_{12}$  is a  $C_{12}^7$ . It has no double points, and it touches twelve generators of  $F$ .

This surface  $f$  is generated by a (2, 1) correspondence between an elliptic cubic and a conic with one united point. The conic is part of the double curve. If  $P$  is the united point of the correspondence the plane of the cubic meets that of the conic in a line through  $P$ ; this line will meet the cubic again in two points  $P', P''$  and the conic in one point  $Q$ ; the points  $P'$  and  $P''$  must be the two points of the cubic which correspond to the point  $Q$  of the conic.

The pairs of points of the cubic which correspond to the points of the conic form a  $g_2^1$  on the curve; thus these pairs must be on lines through a fixed point of the curve, in fact through  $P$ . One of them  $PP'P''$  passes through its corresponding point  $Q$ , so that the planes joining the points of the conic to the corresponding pairs of points on the cubic touch a quadric cone which is part of the bitangent developable of the surface.

**299.** We shall now suppose the position of  $l$  to be specialised so that  $l$  lies in a solid  $K'$  containing a directrix quartic  $C_4^1$  of  $F$ . Then the projected surface  $f$  has a directrix line  $R$ , the intersection of  $\Sigma$  and  $K'$ . Any plane in  $K'$  passing through  $l$  meets  $C_4^1$  in four points, while a prime through

$K'$  meets  $F$  further in two generators; hence through any point of  $R$  there pass four generators, while any plane through  $R$  contains two generators, and  $f$  is of the type III (B).

A prime through  $l$  meets  $F$  in an elliptic sextic curve, nine of whose chords meet  $l$ ; but this sextic has four intersections with  $C_4^1$  which lie in a plane through  $l$ , and six of the nine chords arise from these. Through any point of  $C_4^1$  there pass three of its chords meeting  $l$ . Hence the chords of  $F$  which meet  $l$  meet  $F$  in the curve  $C_4^1$  counted three times and in a curve  $C_6$ . This curve meets each generator of  $F$  in one point; for the solid determined by  $l$  and the generator meets  $F$  in four further points of which three are on  $C_4^1$ . Thus we have an elliptic curve  $C_6^1$ , and since there are eight tangents of  $C_4^1$  meeting  $l$  there are four tangents of  $C_6^1$  meeting  $l$ , and its projection from  $l$  is a twisted cubic  $C_3^0$ .

To generate this surface we take a line and an elliptic cubic curve, placing them in (1, 4) correspondence with a united point. The sets of four points on the cubic curve which correspond to the points of the line  $R$  form a  $g_4^1$ ; they can therefore be cut out by conics through two fixed points  $P_1, P_2$  of the curve and two other fixed points  $Q_1$  and  $Q_2$  in the plane of the curve. One conic is the line pair  $Q_1Q_2, P_1P_2$ , and this gives rise to a tritangent plane of the surface passing through  $Q_1Q_2$ . The plane of the cubic curve is a second tritangent plane of the surface. Since a  $g_4^1$  and a  $g_2^1$  on an elliptic curve have two pairs of points in common\* there are two planes passing through  $R$  which also belong to that part of the bitangent developable other than the planes through  $R$ ; these join points of  $R$  to pairs of points on the cubic.

**300.** It may happen, however, that the  $g_4^1$  on the cubic curve is such that each of its sets of four points consists of two pairs of a  $g_2^1$ ; this will happen if  $l$  passes through the vertex of a quadric cone containing  $C_4^1$ . We can see this geometrically thus: take two points  $P_1, P_2$  of the cubic, the line joining them meeting the curve again in  $P_3$ . Then take  $Q_1$  and  $Q_2$  on a line through  $P_3$  so that they separate  $P_3$  and its polar line in regard to the cubic harmonically. Then the conics through  $P_1, P_2, Q_1, Q_2$  cut out on the curve a  $g_4^1$  whose sets of four points consist each of two sets of the  $g_2^1$  given by the lines through  $P_3$ . The polar line of  $P_3$  in regard to the cubic curve is also the polar of  $P_3$  in regard to all the conics.

The planes joining the points of  $R$  to the corresponding pairs of the  $g_2^1$  touch a quadric cone with vertex  $P_3$  (since there is a united point). This, as well as the planes of the pencil  $R$ , is part of the bitangent developable of the surface. The remaining part has two double planes, one is the plane of the cubic and the other passes through  $Q_1Q_2$ ; these also touch the quadric cone.

\* Cf. the footnote to § 134.

**301.** Now we may further suppose that  $l$  passes through the vertices of two quadric cones containing  $C_4^1$ . To generate the projected surface in  $\Sigma$  we now take a line and an elliptic cubic in (1, 4) correspondence; but if  $S, T, U, V$  are the four points of the cubic corresponding to a point of the line the pairs  $ST, UV$  must belong to one  $g_2^1$  and the pairs  $SU, TV$  to another  $g_2^1$ . The existence of such sets of points on the cubic curve is clear. For take any point  $X$  of the curve and draw from  $X$  two tangents touching the curve in  $Y$  and  $Z$ . If a line is drawn through  $Y$  meeting the curve again in  $S$  and  $T$  and  $ZS, ZT$  meet the curve again in  $U, V$  then  $Y, U, V$  are collinear\* and the sets of four points  $S, T, U, V$  vary in a series such as we require; different sets of the series are given by the different lines  $YST$  through  $Y$ . There are now two quadric cones belonging to the bitangent developable of the surface.

**302.** The pencil of [4]'s through the solid  $K'$  containing  $C_4^1$  meet  $F$  further in pairs of generators; the solids determined by these pairs of generators meet  $K'$  in planes forming a developable. In general, no plane of this developable will contain a given line in  $K'$ ; but we can choose  $l$  to lie in one of the planes. Then the projected surface  $f$  will have a double generator and will be of the type III (D). The chords of  $F$  which meet  $l$  now meet  $F$  in  $C_4^1$  counted three times,  $g, g'$  and another directrix quartic  $D_4^1$ .

To generate this surface we again take a line and an elliptic cubic in (1, 4) correspondence with a united point  $P$ , but the correspondence must be specialised so that two of the three points of the cubic which correspond to  $P$  and do not coincide with it are collinear with it.

**303.** There is certainly a plane of the developable passing through the vertex of a quadric cone containing  $C_4^1$ ; hence we can take  $l$  to pass through it also. Then the  $g_4^1$  on the elliptic cubic curve is such that each of its sets of four points consists of two sets of a  $g_2^1$ , and the surface has a quadric cone as part of its bitangent developable.

**304.** Now let us take  $l$  to meet the plane of  $\Gamma_1$  in a point  $O$ . A prime passing through  $l$  meets  $F$  in an elliptic sextic curve with a trisecant through  $O$ ; there are six chords of this curve, besides the trisecant, which meet  $l$ . Hence the chords of  $F$  which meet  $l$  meet  $F$  in  $\Gamma_1$  counted twice and a curve  $C_{12}$ . The curve  $C_{12}$  meets each generator of  $F$  in two points, as is seen at once by considering the solid through  $l$  and the generator.

\* This follows from the fact that all cubic curves through eight points of a plane have a ninth point in common.

We have, on projection from  $l$ , a surface  $f$  with a directrix line  $R$ , the intersection of  $\Sigma$  with the solid  $l\Gamma_1$ . Through any point of  $R$  there pass three generators, while any plane through  $R$  contains three generators;  $f$  is of the type III (E). It has a double curve  $C_6$ , the projection of  $C_{12}$ .

The point  $O$  is a sextuple point on  $V_4^{12}$ ,  $l$  having six other intersections with  $V_4^{12}$ . Hence the (3, 1) correspondence between  $C_6$  and  $l$  has six branch-points, so that  $C_6$  is a  $C_6^1$ .  $O$  is also a sextuple point on  $M_4^{12}$ , there being six tangents of  $\Gamma_1$  passing through  $O$  and six other tangents of  $F$  meeting  $l$ ; hence the (1, 2) correspondence between  $C_6^1$  and  $C_{12}$  has six branch-points and  $C_{12}$  is a  $C_{12}^4$ .

Since  $C_{12}^4$  meets each generator of  $F$  twice it must have three double points. These will lie in a plane through  $l$  and give on projection a triple point of  $C_6^1$ . Also  $C_{12}^4$  meets  $\Gamma_1$  in six points lying two on each of three lines through  $O$ , so that  $R$  is a trisecant of  $C_6^1$ .  $C_6^1$  has a triple point not lying on  $R$ .

This surface is generated by a line and an elliptic cubic in (1, 3) correspondence.

**305.** Suppose now that  $l$ , as well as meeting the plane of  $\Gamma_1$  in a point  $O$ , lies in a solid containing a pair of generators of  $F$ . Then the chords of  $F$  which meet  $l$  meet  $F$  in  $g, g'$ ,  $\Gamma_1$  counted twice and a curve  $C_{10}$  which meets every generator of  $F$  in two points. The solid  $gg'$  meets  $F$  further in a point  $X$  of  $\Gamma_1$  and a point  $Y$  of  $\Gamma_2$ ;  $C_{10}$  has a double point at  $Y$ , and meets  $\Gamma_1$  in four points lying two on each of two lines through  $O$ .

We thus have on projection a surface  $f$  whose double curve is  $3R + G + C_5$ ;  $C_5$  has a double point on  $G$  and meets it in one other point, and meets  $R$  in two points.  $R$  and  $G$  intersect (cf. § 273).

The line  $l$  meets  $V_4^{12}$  in six points other than  $O$ ; we have seen (cf. the end of § 293) that four of these six points are branch-points of the (2, 1) correspondence between  $C_5$  and  $l$ , so that  $C_5$  is an elliptic curve  $C_5^1$ . There are six tangents of  $\Gamma_1$  passing through  $O$ , so that there are six other tangents of  $F$  meeting  $l$ ; the (1, 2) correspondence between  $C_5^1$  and  $C_{10}$  has six branch-points. Thus  $C_{10}$  is a  $C_{10}^4$ . It has the one double point  $Y$ .

To generate this surface we again take a line and an elliptic cubic in (1, 3) correspondence; but the correspondence must be specialised to give the double generator.

**306.** Choosing again  $l$  to meet the plane of  $\Gamma_1$  in a point  $O$  we can further secure that the solid  $l\Gamma_1$  contains a generator  $g$  of  $F$ . Then the projected surface  $f$  has a directrix  $R$ —the intersection of this solid with  $\Sigma$ —which is also a generator; through any point of  $R$  there pass three other generators, while any plane through  $R$  contains two other generators. Thus  $f$  is of the type IV (B).

A prime through  $l$  meets  $F$  in an elliptic sextic curve; there is a tri-

secant of this curve meeting  $l$  and the plane containing  $l$  and the trisecant meets the curve in another point. There are three chords of the curve which meet  $l$  and do not lie in this quadrisecant plane.

Thus the points of  $F$  which lie on chords meeting  $l$  and not on  $g$  or  $\Gamma_1$  lie on a curve  $C_6$  of the sixth order. This is an elliptic curve  $C_6^1$  meeting each generator of  $F$  in one point.

To generate the surface in  $\Sigma$  we take a line and an elliptic cubic in (1, 3) correspondence; they have a point of intersection, but it is not a united point.

**307.** Take now a pair of generators  $g, g'$  of  $F$  and let the [4] determined by  $g, g'$  and  $\Gamma_1$  meet  $F$  further in a generator  $h$ . Then the two solids  $gg'$  and  $\Gamma_1 h$  have a plane of intersection; let us project  $F$  from a line  $l$  in this plane which does not meet the surface.

The solid  $l\Gamma_1 h$  meets  $\Sigma$  in a line  $R$  which is a directrix and also a generator of  $f$ ; through any point of  $R$  there pass three other generators, while any plane through  $R$  contains two other generators; there is also a double generator—the intersection of  $\Sigma$  with the solid  $lgg'$ . Thus  $f$  is of the type IV (D).

A prime through  $l$  meets  $F$  in an elliptic sextic curve; there is a trisecant of this curve meeting  $l$ , while the plane of  $l$  and the trisecant meets the curve in a fourth point. There is another chord of the curve meeting  $l, g$  and  $g'$ . There are two other chords of the curve meeting  $l$ , and the points of  $F$ , not on  $g, g', h$  or  $\Gamma_1$ , which lie on chords of  $F$  meeting  $l$  are on an elliptic quartic directrix  $C_4^1$ . This projects into a conic meeting  $R$ ; the double curve of  $F$  is  $6R + G + C_2$ .

To generate  $f$  we take a line and an elliptic cubic in (1, 3) correspondence, the correspondence being specialised to give a double generator. The line meets the cubic but there is no united point. Or we may take a line and a conic in (2, 3) correspondence with a united point, specialising to obtain the double generator.

**308.** Now suppose that  $l$  meets the plane of  $\Gamma_1$  in a point  $O_1$  and the plane of  $\Gamma_2$  in a point  $O_2$ . Then the projected surface has two directrices  $R$  and  $R'$ , and through any point of either of these pass three generators meeting the other. The lines in the plane of  $\Gamma_2$  through  $O_2$  meet  $\Gamma_2$  in sets of a  $g_3^1$ , and the generators of  $F$  through these sets of points meet  $\Gamma_1$  in the sets of a  $g_3^1$ . There is a second  $g_3^1$  on  $\Gamma_1$  cut out by lines through  $O_1$ , and since two series  $g_3^1$  on an elliptic curve have three pairs of points in common we conclude that there are three pairs of generators of  $F$  which determine solids containing  $l$ . The surface  $f$  has then three double generators—it is of the type VI (B).

To generate  $f$  we take two lines in (3, 3) correspondence; the correspondence being specialised to give three double generators.



**309.** There is in [5] an elliptic sextic ruled surface  $F$  with only one directrix cubic  $\Gamma$ ; any three generators which meet  $\Gamma$  in three collinear points lie in a [4] containing  $\Gamma$  and conversely any [4] containing  $\Gamma$  meets  $F$  again in three generators which meet  $\Gamma$  in three collinear points\*.

Now project this surface on to a solid  $\Sigma$  from a line  $l$  meeting the plane of  $\Gamma$  in a point  $O$ . The solid  $l\Sigma$  meets  $\Sigma$  in a line  $R$  which is a directrix of the projected surface  $f$ ; through any point of  $R$  there pass three generators of  $f$  lying in a plane through  $R$ . Thus  $f$  is of the type VII (B).

To generate the surface we take a line and an elliptic quartic in (1, 3) correspondence with a united point  $P$ , the three points of the quartic which correspond to a point of the line lying in a plane through the line. The pencil of planes through the line is related to the range of points on the line; to the point  $P$  corresponds the plane through the line and the tangent at  $P$ .

To a point of the line corresponds a plane through the line meeting the quartic in three points other than  $P$ ; these three points are joined by three chords meeting the line. Conversely, through any point of the line pass two chords of the quartic, and the planes determined by these and the line give two corresponding points on the line. Thus we have a (2, 3) correspondence between the points of the line; but  $P$  counts twice among the five coincidences. There are thus three double generators of  $f$ .

**310.** There exists also in [5] an elliptic sextic ruled surface  $F$  with  $\infty^1$  directrix cubics on it. Then the solid determined by any two generators contains a third\*.

Take such a solid containing three generators  $g, g', g''$ . The planes of the cubic curves meet this solid in transversals of  $g, g', g''$ , i.e. in lines of a regulus. An arbitrary line  $l$  in the solid will thus meet the planes of two of the cubic curves. The planes of the cubic curves form a  $V_3^3$  †.

Projecting from  $l$  on to  $\Sigma$  we have a surface  $f$  with two directrices and a triple generator; through any point of either directrix there pass three generators meeting the other. To generate it we take a (3, 3) correspondence between two lines specialised to give the triple generator; the surface is of the type VI (C).

If, however,  $l$  is chosen to touch the quadric surface determined by the regulus the projected surface will have a triple generator and a single directrix  $R$ ; through any point of  $R$  there pass three generators lying in a plane through  $R$ . The surface is of the type VII (C). To generate this surface we take a line and an elliptic cubic in (1, 3) correspondence; to any point of the line correspond the three intersections of the cubic

\* § 282.

† The locus formed by the  $\infty^1$  planes which meet three lines in [5] in related ranges of points.

with a plane through the line. The plane of the cubic contains the triple generator.

*The normal surface in [5] with a double line*

**311.** There exists in [5] a normal elliptic sextic ruled surface  $F$  with a double line  $\lambda$ ; through every point of  $\lambda$  there pass two generators of  $F^*$ .

The tangent solids of  $F$  all pass through  $\lambda$  and meet an arbitrary [3] in the tangent lines of an elliptic quartic curve. We have thus  $\infty^1$  solids forming a line-cone  $M_4^8$ , so that there are eight tangents of  $F$  meeting an arbitrary line  $l$ .

There is on  $F$  a linear system of  $\infty^2$  directrix quartic curves; through two general points of  $F$  there passes one of these curves, while any two of them intersect in two points through which there passes a pencil of the curves\*.

There is a finite number of chords of  $F$  passing through a general point of [5]; it is at once seen that this number is 2, for if we take one chord of  $F$  through a general point of [5] there is a directrix quartic (and in general only one) passing through its two intersections with  $F$ , and there will be a second chord of this quartic passing through the point of the first chord with which we started.

There are points of [5] for which the two chords of  $F$  coincide; the locus of such points is a primal  $V_4$ . To study this primal  $V_4$  we regard  $F$  as the projection of a normal elliptic ruled surface  $F_0$  of the seventh order in [6] from a point  $O$  of itself.  $V_4$  can be also defined as the locus of the chords of  $F$  which are such that the tangents to the directrix quartic, which passes through the two intersections of the chord with  $F$  at these two points, intersect.

**312.** If we take the general ruled surface of the seventh order in [6] we cannot obtain, as a projection, the sextic surface with a double line. But there is in [6] a normal elliptic ruled surface  $F_0$  of the seventh order having an elliptic cubic curve as a directrix †; if we project this on to [5] from a point  $O$  of the cubic curve  $\gamma$  we obtain the sextic ruled surface  $F$  with a double line.

The chords of  $F_0$  form an  $M_5^7$  on which  $F$  is a quadruple surface and  $\gamma$  a quintuple curve.

There is on  $F_0$  a pencil of elliptic quartic curves with a common point of intersection †; the cubic  $\gamma$  taken with the generator  $g_0$  through this intersection belongs to the pencil of curves. If  $F_0$  is projected from a general point of itself we obtain the general elliptic sextic ruled surface which has two directrix cubic curves; one of these is the projection of  $\gamma$  and the other the projection of that quartic curve of  $F_0$  which passes

\* § 287.

† § 281.

through the centre of projection. If  $F_0$  is projected from the point common to the pencil of quartics we obtain the sextic surface with  $\infty^1$  cubic curves; if  $F_0$  is projected from a point of  $g_0$  we obtain the sextic surface with only one cubic curve. If, however, we project from a point  $O$  of  $\gamma$  we obtain the sextic surface  $F$  with a double line.

The locus of the solids which contain the quartic curves on  $F_0$  is a  $V_4^3$ , the projection on to [5] from the point common to all the solids being the  $V_3^3$  formed by the planes of the  $\infty^1$  cubic curves on the resulting sextic surface\*. This is a double locus on  $M_5^7$ ; it contains  $F_0$ , but  $F_0$  is not a double surface on  $V_4^3$ .

A plane through a point  $O$  of  $\gamma$  meets  $M_5^7$  in a curve of the seventh order with a quintuple point at  $O$ . It meets  $V_4^3$  in two points other than  $O$ , which are double points of the curve. The curve is therefore of genus 3, and eight tangents can be drawn to it from the quintuple point.

Hence, given an elliptic ruled surface  $F$  of the sixth order in [5] which has a double line  $\lambda$ , the primal which is the locus of points of [5] which are such that the two chords of  $F$  which pass through them coincide is  $V_4^8$  of order eight.

**313.** The  $\infty^1$  quartic curves on  $F_0$  give after projection a pencil of quartics on  $F$ ; this pencil on  $F$  has two base-points, one the projection of the single base-point of the pencil on  $F_0$  and the other the intersection of the [5] containing  $F$  with the generator of  $F_0$  through  $O$ . But, in general, a quartic curve on  $F$  is the projection of a quintic curve on  $F_0$  which passes through  $O$ ; on  $F_0$  there are  $\infty^3$  quintic curves, such that through three general points of  $F_0$  there pass two of them†. Thus through  $O$  we have  $\infty^2$  quintic curves on  $F_0$  giving after projection  $\infty^2$  quartic curves on  $F$ .

If  $U$  is the vertex of a quadric cone containing a directrix quartic of  $F$ , the line  $OU$  meets  $\infty^1$  chords of  $F_0$  and lies on  $M_5^7$ , the quartic being regarded as the projection of a quintic of  $F_0$  which passes through  $O$ . The line  $OU$  does not meet  $V_4^3$  except in  $O$ ; thus a plane through  $OU$  meets  $M_5^7$  further in a sextic curve having a quadruple point at  $O$  and two double points. This curve is of genus 2 and six tangents can be drawn to it from  $O$ . Hence if in [5] we take a line  $l$  passing through the vertex of a quadric cone which contains a directrix quartic of  $F$  this line will only meet  $V_4^8$  in six points other than  $U$ ; so that  $U$  is a *double point* on  $V_4^8$ . The same result holds when the quartic on  $F$  is the projection of a quartic on  $F_0$ .

The point  $U$  is a *quadruple point* on the locus  $M_4^8$  formed by the tangents of  $F$ , since four tangents of the quartic curve pass through  $U$ .

The solids joining  $\gamma$  to the generators of  $F_0$  are, like the solids containing the quartic curves of  $F_0$ , double solids on  $M_5^7$ . If then we take a line passing through  $O$  and lying in one of these solids it is a double line on

\* § 310.

† § 286.

$M_5^7$ ; a plane through it meets  $M_5^7$  further in a quintic curve with a triple point at  $O$ . This curve has also two double points at the intersections, other than  $O$ , of the plane with  $V_4^3$ . It is therefore of genus 1 and four tangents can be drawn to it from its triple point. Hence if in [5] we take a line which meets the plane containing  $\lambda$  and a generator of  $F$  it will only meet  $V_4^3$  in four points other than its intersection with the plane. Hence the planes joining  $\lambda$  to the generators of  $F$  are *quadruple planes* on  $V_4^3$ .

**314.** Let us consider now the projection of  $F$  from a line  $l$  of general position on to a solid  $\Sigma$ . The projection of  $\lambda$  is a line  $R$ ; through any point of  $R$  there pass two generators of the projected surface  $f$ , while any plane of  $\Sigma$  passing through  $R$ , being the intersection of  $\Sigma$  with a [4] through  $\lambda$  and  $l$ , contains four generators of  $f$ . This shews that  $f$  belongs to the type III (A).

A [4] through  $l$  meets  $F$  in an elliptic sextic curve with a double point; there are eight chords of this curve which meet  $l$ . Hence the chords of  $F$  which meet  $l$  meet  $F$  in the points of a curve  $C_{16}$  of order 16.

Take any generator  $g$  of  $F$ ; there is a generator  $g'$  which meets it. The [4] determined by  $l$  and the plane  $gg'$  meets  $F$  further in a quartic curve which meets each generator of  $F$  in one point; the solid  $gl$  meets this curve in three points not on  $g$ . We thence conclude that the solid determined by  $l$  and any generator of  $F$  meets  $F$  in three further points. Whence also the curve  $C_{16}$  meets each generator of  $F$  in three points, and therefore the line  $\lambda$  in four points. These four intersections with  $\lambda$  consist in fact of two double points\*.

Hence  $f$  has a double curve  $C_8$  meeting each generator in three points, as well as a double directrix  $R$ . The chords of  $F$  which meet  $l$  meet  $\Sigma$  in the points of  $C_8$ , so that there is a (2, 1) correspondence between the points of  $C_8$  and the points of  $l$ . The eight intersections of  $l$  with  $V_4^3$  are the branch-points of the correspondence, so that, by Zeuthen's formula, we have a  $C_8^3$  of genus 3. Further, there is a (1, 2) correspondence between the points of  $C_8^3$  and the points of  $C_{16}$ ; this also has eight branch-points, since  $l$  meets  $M_4^8$  in eight points. Hence we have a  $C_{16}^9$ .

Since  $C_{16}^9$  meets each generator of  $F$  in three points it must, by Segre's formula for the genus of a curve on a ruled surface, have six double points which are not double points of  $F$ . These double points lie three in each of two planes through  $l$ , so that  $C_8^3$  has two triple points.

The (1, 3) correspondence between a directrix quartic of  $F$  and  $C_{16}^9$

\* The planes of pairs of intersecting generators of  $F$  are projected from  $\lambda$  by solids forming a quadric line-cone; there are two of these solids meeting  $l$ . The pairs of generators contained in these solids meet  $\lambda$  in its intersections with  $C_{16}$ ;  $C_{16}$  has double points at both these points. The plane of the two tangents, at either of these double points, of  $C_{16}$  meets  $l$ . The points project into the two intersections of  $R$  and  $C_8^3$ ; the two generators of  $f$  which pass through either of these points lie in a plane through  $R$ , this plane also containing the tangent of  $C_8^3$ .

shews that  $C_{16}^9$  touches sixteen generators of  $F$ ; whence also  $C_8^3$  touches sixteen generators of  $f$  (cf. § 267).

To generate this surface we take in  $\Sigma$  a line  $R$  and an elliptic quartic in (1, 2) correspondence. The pairs of points of the quartic which correspond to the points of  $R$  give the lines of a regulus; neither of the two lines of the regulus which meet  $R$  must do so in the point of  $R$  which gives rise to it. The planes joining the points of  $R$  to the pairs of points of the quartic which correspond to them give a developable  $E_3^0$  which has two planes passing through  $R$ ; this is part of the bitangent developable of  $f$ .

**315.** Now assume that  $l$ , instead of being of general position, passes through the vertex  $U$  of a quadric cone containing a directrix quartic  $C_4^1$  of  $F$ . Then the chords of  $F$  which meet  $l$  meet  $F$  in  $C_4^1$  and in a curve  $C_{12}$  meeting each generator of  $F$  in two points and having two double points on  $\lambda$ . After projection we have a surface  $f$  whose double curve is  $R + C_2 + C_6$ .

Since  $U$  is a double point of  $V_4^8$  there are six branch-points in the (2, 1) correspondence between  $C_6$  and  $l$ ; thus  $C_6$  is a  $C_6^2$ . Then the (1, 2) correspondence between  $C_6^2$  and  $C_{12}$  has four branch-points, since there are four tangents of  $C_4^1$  passing through  $U$  and therefore four tangents of  $C_{12}$  meeting  $l$ . Hence  $C_{12}$  is a  $C_{12}^5$ . It touches eight generators of  $F$ .

Since  $C_{12}^5$  meets every generator of  $F$  in two points it must have two double points,  $X$  and  $Y$ , not on  $\lambda$ . Through  $X$  there pass two chords  $XX'$  and  $XX''$  of  $C_{12}^5$  which meet  $l$ ;  $X'$  and  $X''$  are on  $C_4^1$  and  $X'X''$  passes through  $U$ , and similarly for  $Y'$  and  $Y''$ .  $C_{12}^5$  meets  $C_4^1$  in four further points lying on two lines through  $U$ . Thus  $C_6^2$  has two double points and two simple points on  $C_2$  (cf. § 268).

To generate this surface we take a line and a conic in (2, 2) correspondence. The pairs of points of the conic which correspond to the points of the line are such that their joins touch another conic; the planes joining the points of the line to their corresponding pairs on the conic form a cubic developable which is part of the bitangent developable of the surface. There are two planes of this developable passing through the line.

**316.** Now suppose that  $l$  passes through the vertex  $U$  of a quadric cone containing a directrix quartic  $C_4^1$  and through the vertex  $V$  of a quadric cone containing a directrix quartic  $D_4^1$ . The chords of  $F$  which meet  $l$  meet  $F$  in  $C_4^1$ ,  $D_4^1$  and a curve  $C_8^1$  meeting each generator of  $F$  once and having two double points on  $\lambda$ .

$C_4^1$  and  $D_4^1$  intersect in two points  $X$ ,  $Y$ .  $UX$  and  $UY$  meet  $C_4^1$  again on  $C_8^1$ , while  $VX$  and  $VY$  meet  $D_4^1$  again on  $C_8^1$ . Thus the projected surface has a double curve  $R + C_2 + D_2 + C_4^1$ ;  $C_2$  and  $D_2$  have two intersections through which  $C_4^1$  passes, while  $C_4^1$  meets  $R$  in two points.  $C_8^1$  meets each of  $C_4^1$  and  $D_4^1$  in six points of which two are already specified; thus  $C_4^1$  has

two intersections with each of  $C_2$  and  $D_2$  other than those already mentioned (cf. § 269). The surface is generated by the lines which meet  $C_2$ ,  $D_2$  and  $R$ .

**317.** The planes of the pairs of generators which intersect in the points of  $\lambda$  form a three-dimensional locus. Let us choose  $l$  to meet a plane of this locus—say the plane which contains two generators  $g$  and  $g'$  of  $F$ . Then projecting from  $l$  on to  $\Sigma$  we have a surface  $f$  with a directrix line  $R$  through every point of which there pass two generators, while a plane through  $R$  contains four generators; but here there is also a double generator and  $f$  is of the type III (C).

Consider the chords of  $F$  which meet  $l$ . A prime through  $l$  meets  $F$  in an elliptic sextic curve with a double point; there are eight chords of this curve meeting  $l$ , but one of these is in the plane  $gg'$ . There are seven others, so that the chords of  $F$  which meet  $l$  meet  $F$  in  $g$ ,  $g'$  and a curve  $C_{14}$  of order 14.  $C_{14}$  meets each generator of  $F$  in three points.

The [4]  $lgg'\lambda$  meets  $F$  further in two generators. Hence the solid  $lgg'$  meets  $F$  further in two points  $X$  and  $Y$ . Through  $X$  there pass transversals to  $l$  and  $g$  and to  $l$  and  $g'$ , so that  $X$  is a double point of  $C_{14}$ ; similarly for  $Y$ .  $C_{14}$  has a double point on  $\lambda$ .

After projection we obtain a surface  $f$  whose double curve is  $R + G + C_7$ ;  $R$  and  $G$  intersect,  $C_7$  has two double points and one ordinary point on  $G$  and meets  $R$  in one point (cf. § 270).

There are eight branch-points in the (2, 1) correspondence between  $C_7$  and  $l$  so that  $C_7$  is a  $C_7^3$ ; also there are eight branch-points in the (1, 2) correspondence between  $C_7^3$  and  $C_{14}$  so that  $C_{14}$  is a  $C_{14}^9$ . It has then just the three double points mentioned. It is touched by sixteen generators of  $F$ .

To generate this surface we take a line and an elliptic quartic and place them in (1, 2) correspondence. The correspondence must, however, be specialised to give the double generator; the pairs of points of the quartic which correspond to the points of the line form a  $g_2^1$  and the lines joining them form a regulus; one of the points in which the regulus meets the line must lie on the corresponding line of the regulus. The planes joining the points of the line to the corresponding elements of the regulus touch a quadric cone which is part of the bitangent developable of the surface. There is one tangent plane of the quadric cone passing through the line.

**318.** Now suppose that  $l$  not only meets the plane  $gg'$  but also passes through the vertex  $U$  of a quadric cone which contains a directrix quartic  $C_4^1$  of  $F$ . Then  $C_4^1$  will meet  $g$  and  $g'$  and pass through  $X$  and  $Y$ , the line  $UX$  meeting  $g$  and the line  $UY$  meeting  $g'$ . There is a [4] containing  $C_4^1$ ,  $g$  and  $g'$ . The chords of  $F$  which meet  $l$  now meet  $F$  in  $g$ ,  $g'$ ,  $C_4^1$  and a curve  $C_{10}$  meeting every generator of  $F$  in two points.  $C_{10}$  passes through  $X$  and  $Y$ ; it has a double point on  $\lambda$ .

Thus on projection we have a surface  $f$  with a double curve  $R + G + C_2 + C_5$ ; the plane of  $C_2$  contains  $G$  and  $C_5$  passes through their two intersections.  $C_2$  and  $C_5$  have two other intersections, these arise from the four points of intersection, other than  $X$  and  $Y$ , of  $C_4^1$  and  $C_{10}$ .  $C_5$  has a further intersection with  $G$  and meets  $R$  (cf. § 271).

Since  $U$  is a double point of  $V_4^8$  there are six branch-points in the (2, 1) correspondence between  $C_5$  and  $l$ ; hence  $C_5$  is a  $C_5^2$ . As there are four tangents of  $C_4^1$  passing through  $U$  there are four tangents of  $C_{10}$  meeting  $l$ ; thus the (1, 2) correspondence between  $C_5^2$  and  $C_{10}$  has four branch-points, so that  $C_{10}$  is a  $C_{10}^5$ . It has no double points other than the one on  $\lambda$ , and touches eight generators of  $F$ .

To generate this surface we take a line and a conic in (2, 2) correspondence; but the correspondence must be specialised to give the double generator, to the point of the line in the plane of the conic must correspond two points of the conic collinear with it. Then the planes joining the points of the line to the pairs of points of the conic which correspond to them touch a quadric cone which is part of the bitangent developable of the surface.

**319.** The planes which project the generators from  $\lambda$  form a quartic line-cone of three dimensions. Let us choose  $l$  to meet a plane of this cone, the plane joining  $\lambda$  to a generator  $g$ . Then the solid  $l\lambda g$  meets  $\Sigma$  in a line  $R$  which is a directrix and also a generator of the projected surface  $f$ ; any plane through  $R$  contains three other generators, while through any point of  $R$  there pass two other generators. Thus  $f$  is of the type IV (A).

Any prime through  $l$  meets  $F$  in an elliptic sextic curve with a double point, the plane of  $l$  and this double point meeting the curve again. There are six other chords of the curve meeting  $l$ . Hence the points of intersection of  $F$ , other than  $g$  and  $\lambda$ , with its chords which meet  $l$  lie on a curve  $C_{12}$ . By considering the intersection of  $F$  with a [4] through the solid determined by  $l$  and an arbitrary generator we see that  $C_{12}$  meets each generator of  $F$  in two points; it will then have four intersections with  $\lambda$ , in fact it has a double point on  $\lambda$  and meets it in two other points.

The plane  $\lambda g$  is a quadruple plane of  $V_4^8$ , so that there are only four branch-points in the (2, 1) correspondence between  $C_6$  and  $l$ ; thus  $C_6$  is an elliptic curve  $C_6^1$ . There are six\* tangents of  $C_{12}$  meeting  $l$ , so that the (1, 2) correspondence between  $C_6^1$  and  $C_{12}$  has six branch-points; whence  $C_{12}$  is a  $C_{12}^4$ . It touches six generators of  $F$ .

Since  $C_{12}^4$  meets each generator of  $F$  in two points it must have three double points not on  $\lambda$ ; these lie in a plane through  $l$  and give rise on projection to a triple point of  $C_6^1$ .

To generate this surface we take in  $\Sigma$  a line and an elliptic quartic in

\* The number of tangents of an elliptic quartic in [3] meeting a line which meets the curve.

(1, 2) correspondence; they have a point of intersection but it is not a united point. The pairs of points of the quartic which correspond to the points of the line form a  $g_2^1$  and the lines joining the pairs form a regulus. When we join the points of the line to the corresponding lines of the regulus we have a developable of the third class which is part of the bitangent developable of the surface. This developable has two planes passing through the line.

**320.** Now let us choose  $l$  to meet the plane of two intersecting generators  $g$  and  $g'$  and also to meet the plane through  $\lambda$  and a generator  $h$ . The solid  $l\lambda h$  meets  $\Sigma$  in a line  $R$  which is a directrix and also a generator of  $f$ ; through any point of  $R$  there pass two further generators, while any plane through  $R$  contains three further generators. Also the surface has a double generator—the intersection of  $\Sigma$  with the solid  $lgg'$ .

A [4] through  $l$  meets  $F$  in an elliptic sextic curve with a double point, the plane through this point and  $l$  meeting the curve again. Also there is a chord of the curve meeting  $g$ ,  $g'$  and  $l$ . There are five further chords of the curve meeting  $l$ . Hence the points of  $F$  other than  $g$ ,  $g'$ ,  $h$  which lie on chords of  $F$  meeting  $l$  are on a curve  $C_{10}$ .  $C_{10}$  meets each generator in two points and has two intersections with  $\lambda$ . The solid  $lgg'$  meets  $F$  in two further points, one of which is on  $h$ ; through the other there pass transversals to  $l$  and  $g$  and to  $l$  and  $g'$ , so that it is a double point of  $C_{10}$ .

The projected surface  $f$  is of the type IV (C) and has a double curve  $3R + G + C_5$ .  $R$  meets  $G$  and is a chord of  $C_5$ ;  $C_5$  has a double point on  $G$  and meets it in one other point (cf. § 275).

The plane  $\lambda h$  being a quadruple plane on  $V_4^8$  the (2, 1) correspondence between  $C_5$  and  $l$  has only four branch-points; thus  $C_5$  is an elliptic curve  $C_5^1$ . There are six tangents of  $C_{10}$  meeting  $l$ , so that the (1, 2) correspondence between  $C_5^1$  and  $C_{10}$  has six branch-points; whence  $C_{10}$  is a  $C_{10}^4$ . It has only one double point and touches six generators of  $F$ .

To generate this surface we take a line and an elliptic quartic in (1, 2) correspondence. The correspondence must, however, be specialised to give the double generator; one of the points of the line gives rise to the two intersections of the quartic with one of its chords through that point. The line meets the quartic, but there is no united point. The planes joining the points of the line to the pairs of points of the quartic which correspond to them now touch a quadric cone, which is part of the bitangent developable of the surface. There is one tangent plane of the cone passing through the line.

**321.** If we choose  $l$  to lie in the solid through  $\lambda$  and a pair of intersecting generators  $g$  and  $g'$  this solid meets  $\Sigma$  in a line  $R$  which is a directrix and also a double generator of  $f$ ; through any point of  $R$  there pass two further generators, while any plane through  $R$  contains two further generators, so that  $f$  is of the type V.



To generate  $f$  we take an elliptic quartic and one of its chords and place them in (2, 1) correspondence without any united points. The pairs of points of the quartic corresponding to the points of the line give the lines of a regulus and the planes joining the points of the line to the corresponding pairs of points on the quartic form a cubic developable  $E_3^0$  which is part of the bitangent developable of the surface, two planes of  $E_3^0$  passing through the line.

**322.** The directrix quartics of  $F$  are all given by primes through two fixed intersecting generators. It is thus clear that an arbitrary line  $l$  does not lie in a solid with any one of these quartic curves. Let us, however, choose  $l$  to lie in a solid  $K'$  containing a directrix quartic.

Then the projected surface  $f$  has two directrices;  $R$ , the intersection of  $\Sigma$  and  $K'$ , and  $R'$ , the intersection of  $\Sigma$  with the solid  $l\lambda$ . Through any point of  $R$  there pass four generators of  $f$  all meeting  $R'$ , while through any point of  $R'$  there pass two generators of  $f$  meeting  $R$ . The pairs of intersecting generators of  $F$  give planes meeting  $K'$  in the lines of a regulus;  $l$  meets two of these lines so that  $f$  has two double generators. Thus  $f$  is of the type VI (A).

To generate this surface we take two lines  $R$  and  $R'$  in (2, 4) correspondence, the correspondence being specialised to give two double generators.

**323.** The quadric line-cone which projects the  $\infty^1$  pairs of intersecting generators of  $F$  from  $\lambda$  has another system of generating solids; each solid of this other system contains two generators of  $F$  which do not intersect.

If we now choose  $l$  to lie in one of these latter solids we have on projection a surface with a directrix  $R$  which is also a double generator; through any point of  $R$  there pass two generators lying in a plane through  $R$ . The surface  $f$  is of the type VII (A).

To generate the surface we take in [3] an elliptic quartic and one of its chords. These determine a quadric surface; we place the quartic and the line in (2, 1) correspondence without united points so that the two points of the quartic corresponding to any point of the line are on the same generator of the quadric, this generator belonging to the opposite system to the line. The range of points on the line is related projectively to the system of generators which meet the line. There are two generators of the quadric which meet the line in the points which correspond to them; these are the two double generators of the sextic surface.

**324.** A table shewing the different types of elliptic sextic ruled surfaces in ordinary space is given on p. 309. There are thirty-four different types. We have already obtained twenty-eight of them; the others will be found at the end of Chapter VI.

## SECTION III

SEXTIC RULED SURFACES WHICH ARE NEITHER  
RATIONAL NOR ELLIPTIC*The normal sextic curve of genus 2*

**325.** The sextic curve of genus 2 is normal in [4] and cannot lie in a space of higher dimension. The pairs of points\* of the  $g_2^1$  give  $\infty^1$  chords of the curve forming a rational ruled surface  $\phi$ , the order of  $\phi$  being the number of its generators which meet an arbitrary plane. Now the solids passing through the plane cut out a  $g_6^1$  on the curve  $C$ , and on a curve of genus 2 a  $g_6^1$  and a  $g_2^1$  have three pairs of common points †. Hence  $\phi$  is a cubic ruled surface ‡.

The projection of  $C$  from one of its chords on to a plane is a quartic with one double point; thus each chord of  $C$  is met by one other chord, and there are  $\infty^2$  quadrisecant planes of  $C$ .

The projection of  $C$  from a point of itself on to a solid is a quintic curve of genus 2; this lies on a quadric surface and its trisecants form a regulus thereon. Hence those quadrisecant planes of  $C$  which pass through a point of the curve form one system of planes of a quadric point-cone. The other system of planes of the cone cuts out the  $g_2^1$  on  $C$ ; thus the generators of  $\phi$  meet all the quadrisecant planes of  $C$ . These quadrisecant planes are in fact none other than the planes of the  $\infty^2$  conics on  $\phi$ .

If we take two points  $P$  and  $Q$  of  $C$  we have two quadric point-cones with vertices  $P$  and  $Q$ ; these have in common the quadrisecant plane through  $PQ$  and the cubic ruled surface  $\phi$ .

There are  $\infty^3$  trisecant planes of  $C$ ; through any point of [4] there pass  $\infty^1$  of these. Consider, in particular, a point of [4] which is the intersection of two chords of  $C$ ; the projection of  $C$  from such a point on to a solid is a sextic of genus 2 with two double points. The trisecants of this sextic form both systems of generators of a quadric surface. Hence the trisecant planes of  $C$  which pass through an intersection of two of its chords and do not contain either chord are the planes of a quadric point-cone.

There are  $\infty^3$  quadrics containing  $C$ ; we have noticed already  $\infty^2$  point-cones among these. There is also a line-cone whose vertex is the directrix of  $\phi$ . This directrix is a chord of  $C$ , and the quadrisecant planes containing the pairs of points of the  $g_2^1$  all contain this directrix.

**326.** The projection of  $C$  from a line on to a plane is a sextic with eight double points, while the projection from a line which meets it is a quintic with four double points. Hence the chords of  $C$  form a locus  $M_3^8$  on which

\* There is one, and only one,  $g_2^1$  on a curve of genus 2.

† Cf. the footnote to § 134.

‡ Cf. the footnote to § 19.

$C$  is a quadruple curve. There is also a double surface on  $M_3^8$ , the locus of points ( $\infty^2$  in aggregate) through which pass two chords of  $C$ .

Let us take a plane section of  $M_3^8$  and consider the (5, 2) correspondence between it and  $C^*$ . There are fourteen tangents of  $C$  meeting any plane; thus there are fourteen branch-points on the plane section of  $M_3^8$  and fifty-six branch-points on  $C$ . Then Zeuthen's formula† shews that the plane sections of  $M_3^8$  are of genus 14, and they therefore have seven double points. There is thus on  $M_3^8$  a double surface  $F_2^7$  of order 7.

This double surface contains the directrix of  $\phi$ . It also contains  $C$ ;  $C$  is a simple curve on  $F_2^7$ , the section of  $M_3^8$  by a quadrisecant plane of  $C$  consists of six chords of  $C$  and a conic of  $\phi$ .

There are curves  $C$  for which  $\phi$  degenerates into a cubic cone. The intersection of a cubic cone in [4] (i.e. a cone projecting a twisted cubic from a point  $V$  outside its space) with a quadric threefold not passing through  $V$  is a normal sextic curve of genus 2. The  $g_2^1$  is given by the generators of the cubic cone, and there are six tangents of  $C$  passing through  $V$ . The projection of  $C$  from a point of itself gives a quintic curve of genus 2 lying on a quadric cone.

### *Sextic curves of genus 2 which lie on quadrics*

**327.** Following our usual division of the curves on a quadric  $\Omega$  in [5] into different classes, we consider five classes of sextic curves of genus 2 as follows:

I. The normal sextic  $C$  in [4].

II.  $C$  lies on the quadric point-cone in which  $\Omega$  is met by a tangent prime at a point  $O$ .

(A)  $C$  meets every  $\omega$ -plane of  $\Omega$  through  $O$  in two points and every  $\rho$ -plane of  $\Omega$  through  $O$  in four points, a chord of  $C$  passing through  $O$ .

(B)  $C$  meets every  $\omega$ -plane of  $\Omega$  through  $O$  in four points and every  $\rho$ -plane of  $\Omega$  through  $O$  in two points, a chord of  $C$  passing through  $O$ .

(C)  $C$  meets every plane of  $\Omega$  through  $O$  in three points, two chords of  $C$  passing through  $O$ .

III.  $C$  again lies in a tangent prime of  $\Omega$  but now passes through the point of contact  $O$ .

(A)  $C$  meets every  $\omega$ -plane of  $\Omega$  through  $O$  in two points other than  $O$  and every  $\rho$ -plane of  $\Omega$  through  $O$  in three points other than  $O$ .

(B)  $C$  meets every  $\omega$ -plane of  $\Omega$  through  $O$  in three points other than  $O$  and every  $\rho$ -plane of  $\Omega$  through  $O$  in two points other than  $O$ .

IV.  $C$  lies on a quadric surface in a [3] through which pass two tangent primes of  $\Omega$ .

(A)  $C$  meets the generators of one system each in two points and the generators of the other system each in four points, and has a double point.

\* Cf. § 262.

† § 16.

(B)  $C$  meets the generators of both systems each in three points, and has two double points.

V.  $C$  lies on a quadric cone; the section of  $\Omega$  by a [3] which touches it.

(A)  $C$  has a double point at the vertex and one other double point, meeting every generator in two points other than the vertex.

(B)  $C$  meets every generator in three points and has two double points.

**328.** We have in fact already met with the classes I, II and III in § 325. Regarding  $C$  as the normal curve in [4] we can have non-specialised quadrics containing it, from which arises I. The quadric point-cones which project the surface  $\phi$  from points of itself not on  $C$  give II (A) and II (B), while those projecting it from points on  $C$  give III (A) and III (B). The quadric point-cones which have their vertices at intersections of two chords of  $C$  occur in the type II (C).

*The sextic ruled surfaces in [3] whose plane sections are curves of genus 2*

**329.** We regard a non-degenerate quadric containing the normal curve  $C$  as a prime section of a quadric  $\Omega$  in [5]. Then  $C$  will represent the generators of a ruled surface in [3] belonging to a linear complex.

Through any point of  $C$  there pass four of its chords which lie on  $\Omega$ ; the chords of  $C$  lying on  $\Omega$  form a ruled surface of order 16—the intersection of  $\Omega$  and  $M_3^8$ —on which  $C$  is a quadruple curve. The complete intersection of  $\Omega$  and  $F_2^7$  is of order 14; it consists of  $C$  and a curve of order eight. Thus the ruled surface of order 16 has a quadruple curve of order six and a double curve of order eight. A section of this ruled surface by a solid will be a curve of order 16 with six quadruple points and eight double points, the curve lying on a quadric and meeting each generator in eight points. Projecting this curve into a plane curve of order 15 we see that its genus is

$$91 - 2 \cdot 21 - 6 \cdot 6 - 8 = 5,$$

so that the double curve and bitangent developable of the ruled surface in [3] are also of genus 5.

Any plane of  $\Omega$  meets the [4] containing  $C$  in a line which is met by eight chords of  $C$ ; hence the double curve of the ruled surface is  $C_8^5$  and the bitangent developable  $E_8^5$ .

There are no triple points or tritangent planes because  $C$  has no trisecants.

This ruled surface in [3] of the sixth order and with plane sections of genus 2 cannot contain any directrix lines, conics or cubic curves unless they are multiple curves. But there are on the surface  $\infty^1$  plane quartics of genus 2 each with a double point, these being the intersections of the surface with those planes which contain two intersecting generators.

There is a finite number of these curves passing through a general point  $A$  of the surface; this number is simply the number of planes of the bitangent developable which pass through  $A$  and do not contain the generator through  $A$ . But of the eight planes of  $E_8^5$  passing through  $A$  there are four which contain the generator; hence there are four plane quartics passing through a general point of the surface.

**330.** Suppose now that  $C$  is of the type II (A); we regard the quadric cone containing  $C$  as the section of  $\Omega$  by a tangent prime. Then we have in [3] a sextic ruled surface with a directrix line  $R$ ; through any point of  $R$  there pass two generators, while any plane through  $R$  contains four generators.

The chords of  $C$  lying in the  $\varpi$ -planes join the pairs of the  $g_2^1$ ; they form the cubic ruled surface  $\phi$ . The chords lying in the  $\rho$ -planes therefore form a ruled surface of order 13 on which  $C$  is a triple curve. The intersection of  $F_2^7$  with the quadric point-cone is made up of  $C$ , the directrix of  $\phi$ , and a curve of order seven which is a double curve on the latter ruled surface. The section of  $\phi$  by an arbitrary solid is a twisted cubic lying on a quadric; the other ruled surface gives a curve of order 13 lying on a quadric, meeting all generators of one system in seven points and all of the other system in six points; the curve has six triple points and seven double points. This curve is of genus

$$66 - 21 - 15 - 18 - 7 = 5.$$

Hence the chords of  $C$  lying in the  $\varpi$ -planes form a rational cubic ruled surface  $\phi$  and those lying in the  $\rho$ -planes form a ruled surface of order 13, the genus of whose prime sections is 5.

Any arbitrary plane  $\rho$  of  $\Omega$  meets the [4] containing  $C$  in a line and the plane  $\varpi$  containing this line joins it to  $O$ ; there are seven other chords of  $C$  meeting the line, all of them belonging to the ruled surface of order 13.

Hence the double curve of the ruled surface in [3] is  $R + C_7^5$ ;  $R$  and  $C_7^5$  have one intersection, represented on  $\Omega$  by the plane  $\varpi$  which contains the chord of  $C$  passing through  $O$ . Also the bitangent developable is  $6R + E_2$ , there being a plane of  $E_2$  containing  $R$ .

Similarly, when  $C$  is of the type II (B) we have a double curve  $6R + C_2$  and a bitangent developable  $R + E_7^5$ ;  $C_2$  has one intersection with  $R$ , while  $E_7^5$  has one plane passing through  $R$ .

**331.** If  $C$  is of the type II (C) we have in [3] a ruled surface with a directrix line  $R$  through any point of which there pass three generators, while any plane through  $R$  contains three generators.

The chords of  $C$  lying in either system of planes form a ruled surface of order eight on which  $C$  is a double curve, and there will be no other double curve. A prime section of this surface gives a curve of order eight lying on a

quadric, meeting all generators of one system in three points and all of the other system in five points, the curve having six double points. The genus of such a curve is

$$21 - 3 - 10 - 6 = 2.$$

An arbitrary plane of  $\Omega$  meets the [4] containing  $C$  in a line, and the plane of the opposite system through this line joins it to  $O$ . This contains three chords of  $C$ ; there are five others meeting the line.

Hence the double curve of the ruled surface in [3] is  $3R + C_5^2$  and the bitangent developable is  $3R + E_5^2$ .  $C_5^2$  has two intersections with  $R$  represented on  $\Omega$  by the  $\varpi$ -planes containing the two chords of  $C$  which pass through  $O$ ;  $E_5^2$  has two planes passing through  $R$  represented by the  $\rho$ -planes through the same two chords of  $C$ .

**332.** Let  $C$  be of the type III (A). Then we have in [3] a ruled surface with a directrix line  $R$  which is also a generator; through any point of  $R$  there pass two other generators, while any plane through  $R$  contains three other generators.

The  $\varpi$ -planes cut out the  $g_2^1$  on  $C$ . The ruled surface formed by the chords of  $C$  which lie on  $\Omega$  here splits up into three parts: the quintic cone of genus 2 projecting  $C$  from  $O$ , the rational cubic ruled surface  $\phi$  and a ruled surface of order eight formed by the chords lying in the  $\rho$ -planes and not passing through  $O$ ; the prime sections of this latter surface are of genus 2 as in § 331.

An arbitrary plane  $\rho$  meets the [4] containing  $C$  in a line; the plane  $\varpi$  through this line joins it to  $O$  and contains three chords of  $C$ . There are five other chords of  $C$  meeting the line and all belonging to the ruled surface of order eight. Hence the double curve of the ruled surface in [3] is  $3R + C_5^2$ .  $R$  and  $C_5^2$  have two intersections: taking that set of the  $g_3^1$  cut out by the  $\rho$ -planes which contains  $O$ , these intersections are represented on  $\Omega$  by the  $\varpi$ -planes containing the chords of  $C$  which join  $O$  to the two other points of the set.

An arbitrary plane  $\varpi$  meets the [4] containing  $C$  in a line; the plane  $\rho$  through this line joins it to  $O$  and contains six chords of  $C$ . There are two other chords of  $C$  meeting the line, both belonging to  $\phi$ . Hence the bitangent developable of the ruled surface in [3] is  $6R + E_2$ . There is one plane of  $E_2$  passing through  $R$ : taking the set of the  $g_2^1$  containing  $O$ , this plane is represented on  $\Omega$  by the  $\rho$ -plane containing the chord joining the pair of points.

Similarly, if  $C$  is of the type III (B) we have in [3] a ruled surface with a double curve  $6R + C_2$  and a bitangent developable  $3R + E_5^2$ .  $R$  and  $C_2$  have one intersection, while there are two planes of  $E_5^2$  passing through  $R$ .

**333.** In type IV we have two directrices  $R$  and  $R'$ .

In IV (A) the double curve is  $6R + G + R'$  and bitangent developable

$R + G + 6R'$ . Through any point of  $R$  pass four generators lying in a plane through  $R'$ , while through any point of  $R'$  pass two generators lying in a plane through  $R$ .

In IV (B) the double curve is  $3R + G + H + 3R'$  and bitangent developable  $3R + G + H + 3R'$ . Through any point of either directrix pass three generators in a plane through the other.

In type V we have a single directrix  $R$ .

In V (A) this is also a double generator; the double curve is  $7R + G$  and bitangent developable  $7R + G$ ; the plane section is a sextic curve with a double point and a quadruple point at which two of the four branches have the same tangent.

In V (B) the double curve and bitangent developable are  $6R + G + H$ . Through any point of  $R$  there pass three generators all lying in a plane through  $R$ . The plane sections are sextic curves with two double points and a triple point at which all three branches have the same tangent.

**334.** The ten different types of sextic ruled surfaces in [3] which we have obtained are shewn in tabular form on p. 310.

*A sextic ruled surface in [4] whose prime sections are curves of genus 2, and the ruled surfaces in [3] derived from it by projection*

**335.** Take in [4] a plane and a line  $R$  which do not intersect. Take in the plane a quartic with one double point  $P$ ; then relate the range of points on  $R$  to the pencil of lines through  $P$ ; each of these lines meets the quartic in two further points, and if these pairs are joined to the corresponding points on  $R$  the joining lines generate a ruled surface on which  $R$  is a double line and  $P$  a double point. The ruled surface is of the sixth order and its prime sections are of genus  $2^*$ .

The plane of the two generators intersecting in a point of  $R$  necessarily passes through  $P$ ; we have thus  $\infty^1$  planes through  $P$ . These planes, as joining the points of a range to lines of a related pencil, generate a quadric point-cone, with vertex  $P$ , containing the ruled surface. The plane  $PR$  is a plane of the opposite system of this cone.

A solid through the plane of two intersecting generators meets the surface further in a plane quartic with a double point at  $P$ ; we have thus  $\infty^1$  plane quartics. Through any one of these quartics there pass  $\infty^1$  solids joining it to the points of  $R$ , each meeting the surface in two intersecting generators. The planes of these  $\infty^1$  quartics are the opposite system of

\* The genus of the prime sections is the same as the genus of the plane quartic. The order of the surface is  $1 \cdot 2 + 4 \cdot 1 = 6$ ; there being a (1, 2) correspondence between the line and the quartic; cf. § 19.

This surface in [4] is mentioned by C. V. Hanumanta Rao, *Proc. Lond. Math. Soc.* (2), 19 (1919), 249.

planes of the quadric point-cone. Through any point of the surface there passes one of the quartic curves.

**336.** Now project this surface  $F$  from a point  $O$  on to a [3]  $\Sigma$ . A solid through  $O$  meets  $F$  in a sextic curve of genus 2 with a double point on  $R^*$ ; there are seven chords of this curve passing through  $O$ . There are  $\infty^1$  chords of  $F$  passing through  $O$  and they meet  $F$  in a curve  $C_{14}$  of order 14. Since a plane through a generator  $g$  of  $F$  meets  $F$  in three further points† the curve  $C_{14}$  meets each generator of  $F$  in three points.

A solid through  $O$  which contains  $P$  meets  $F$  in a sextic curve of genus 2 with two double points‡, one at  $P$  and one on  $R$ . There are six chords of this curve passing through  $O$ . We conclude that  $C_{14}$  has a double point at  $P$ . It has also § a double point on  $R$ .

We can now || calculate that  $C_{14}$  is of genus 14.

On projecting from  $O$  we obtain in  $\Sigma$  a ruled surface whose double curve consists of a directrix and a curve of the seventh order meeting each generator in three points. This curve is in (1, 2) correspondence with  $C_{14}$  and its genus can be calculated by Zeuthen's formula when we know the number of branch-points of the correspondence, i.e. the number of tangents of  $C_{14}$  which pass through  $O$ .

Now the tangent solids ¶ of  $F$  all pass through  $R$ ; there are ten of these solids passing through any point of [4], since a plane quartic of genus 2 is of class ten. Hence there will be ten tangents of  $C_{14}$  passing through  $O$ . Whence the double curve of the projected surface is of genus 5.

Hence on projection we have a surface of the type II (A) with a double curve  $R + C_7^5$ .  $R$  and  $C_7^5$  have one intersection; the tangent of  $C_7^5$  at this point lies in the plane containing  $R$  and the two generators of the surface which intersect there.

To generate the surface in  $\Sigma$  we take a line and a plane quartic with one double point, placing them in (1, 2) correspondence. The bitangent planes of the surface which do not pass through the line join the points of a range

\* This curve has  $\infty^1$  quadrisecants; it lies on a quadric, meeting all generators of one system in four points and all of the other system in two points.

† The solid through the plane and the generator which intersects  $g$  meets  $F$  further in one of the plane quartics; this has three intersections with the plane which do not lie on  $g$ .

‡ This curve lies on a quadric cone with vertex  $P$ , meeting every generator in two points other than  $P$ .

§ The solids joining  $R$  to the pairs of intersecting generators of  $F$  form a pencil of solids through the plane  $RP$ ; there will be one of them passing through  $O$ . The point of  $R$  which is the intersection of the two generators contained in this solid is a double point of  $C_{14}$ .

|| See § 17. Observe that no deduction is made for the double points of  $C_{14}$ ; this is because the double points of  $C_{14}$  are also double points of the ruled surface.

¶ See § 51.



to the lines of a related pencil and therefore touch a quadric cone. There is one tangent plane of this cone passing through the line, viz. the plane joining the line to the double point of the quartic curve.

**337.** We have projected the surface  $F$  from a general point  $O$  of [4]; we now proceed to consider the surfaces obtained in  $\Sigma$  by choosing special positions of the centre of projection  $O$ . Let us then choose  $O$  to lie on the  $M_3^4$  formed by the planes joining  $R$  to the generators of  $F$ , i.e.  $O$  lies in the plane containing  $R$  and a generator  $g$ . Then on projection we have in  $\Sigma$  a surface  $f$  with a directrix line  $R$  which is also a generator. Through any point of  $R$  there pass two other generators, while any plane through  $R$  contains three other generators. Hence  $f$  belongs to the type III (A).

A prime through  $O$  meets  $F$  in a sextic of genus 2 with a double point on  $R$ , and the line joining  $O$  to the double point meets the curve again in a point on  $g$ . There are five other chords of this curve passing through  $O$ , so that the chords of  $F$  passing through  $O$  meet  $F$  in a curve  $C_{10}$  of order ten. The plane joining an arbitrary generator of  $F$  to  $O$  meets  $g$ , and meets  $F$  only in two further points; hence every generator of  $F$  meets  $C_{10}$  twice.  $C_{10}$  has a double point at  $P$  but no other double points, and is of genus 7.  $R$  meets  $C_{10}$  in two distinct points; the lines joining these points to  $O$  meet  $C_{10}$  again in its intersections with  $g$ .

From a point on a plane quartic of genus 2 we can draw eight tangents to the curve, so that there are eight tangents of  $C_{10}$  passing through  $O$ . This shews that the projection of  $C_{10}$  is a quintic curve of genus 2.

Hence the surface  $f$  has a double curve  $3R + C_5^2$ .  $R$  and  $C_5^2$  have two intersections.

To generate this type of surface in  $\Sigma$  we take a plane quartic with one double point and a line which meets it, placing the line and the curve in (1, 2) correspondence without a united point. The bitangent planes of the surface which do not contain the line touch a quadric cone, one of whose tangent planes contains the line.

**338.** Let us now choose  $O$  to lie on the quadric point-cone containing  $F$ . Then  $O$  lies in the plane of two intersecting generators and also in the plane of a directrix quartic of  $F$ . This latter plane will meet  $\Sigma$  in a directrix  $R'$  of the projected surface, while we have as before a directrix  $R$ . Through any point of  $R$  there pass two generators, while through any point of  $R'$  there pass four generators. The projected surface has a double generator and is of the type IV (A).

To generate this surface we place two lines  $R$  and  $R'$  in (4, 2) correspondence, the correspondence being specialised to give the double generator.

Now let us further specialise the position of  $O$  and choose it in the plane  $PR$ ; this plane contains two generators of  $F$ . Any solid through this plane meets  $F$  further in two generators intersecting in a point of  $R$ . Hence on projection we obtain in  $\Sigma$  a surface  $f$  with a directrix line  $R$  which is also a double generator, any plane through  $R$  meeting  $f$  further in two generators which intersect on  $R$ :  $f$  is of the type V (A). The other double generator is the intersection of  $\Sigma$  with the other plane of the point-cone containing  $O$ .

To generate this surface we take a plane quartic with a double point  $P$  and a line  $R$  passing through  $P$ ; we then set up a (1, 2) correspondence between  $R$  and the curve without any united point. There is a double generator passing through  $P$  and lying in the plane of the quartic.

*The sextic ruled surfaces of genus 2 which are normal in [3]*

**339.** Although we have not obtained all the surfaces in [3] as projections of this surface  $F$  in [4], we cannot construct any other ruled surfaces in [4] which are of the sixth order and have prime sections of genus 2. We thus meet abruptly the fact that *the normal space for a ruled surface is not uniquely determined by its order and the genus of its prime sections*. The normal space is uniquely determined for rational and elliptic ruled surfaces of all orders, but for ruled surfaces of given order, whose sections are of genus greater than or equal to 2, there are different normal spaces.

We have obtained ten different kinds of sextic ruled surfaces in [3] whose plane sections are of genus 2; of these four can be obtained as the projections of a single normal surface in [4], but the other six are normal in the space [3] itself. On these latter surfaces the plane sections form a complete linear system of curves; such a system of curves is not contained in any linear system, of greater freedom, of curves of the same order.

**340.** For those surfaces which are not obtainable by projection from [4] we here give methods of generation.

To obtain a surface of the type I we take two plane quartics of genus 2 and place them in (1, 1) correspondence with two united points, the curves having the same moduli. It is clear that the plane of either curve is met in eight points by the double curve of the ruled surface so generated. The other curve meets this plane in two points which are not united points; through either of these points passes a generator of the surface to the corresponding point on the first curve, this meets the first curve in three further points which are points of the double curve. We thus have six points of the double curve; to these must be added the double point of the quartic and the intersection of the two generators.

For a surface of the type II (B) we take a line and a conic with one intersection and place them in (2, 4) correspondence with a doubly united

point. Through any point of the line pass four generators, while any plane through the line contains two generators.

For a surface of the type II (C) we take a line and a plane quartic of genus 2, placing them in (1, 3) correspondence with a united point. Through any point of the line there pass three generators, while any plane through the line contains three generators. The double curve of the ruled surface so generated (apart from the triple line) meets the plane of the quartic in five points; through the united point there pass two lines to corresponding points on the quartic, these meet the quartic each in two other points which are points of the double curve. We have thus four points of the double curve, and to these must be added the double point of the quartic.

For a surface of the type III (B) we take a line and a conic with one intersection, placing them in (2, 3) correspondence with a united point. The ruled surface has the line as a generator, through any point of the line pass three other generators, while any plane through the line contains two other generators.

A surface of the type IV (B) is generated by two lines in (3, 3) correspondence, the correspondence being specialised to give two double generators.

To obtain a surface of the type V (B) we take a line and a plane quartic of genus 2 with one intersection, placing them in (1, 3) correspondence with a united point and so that the triads of points on the quartic corresponding to the points of the line are all collinear with the united point. This is secured by relating the range of points on the line to the pencil of lines through the united point; the united point considered as a point of the range corresponding to the tangent to the quartic. This tangent is a double generator of the ruled surface; there is a second double generator passing through the double point of the quartic.

**341.** If, on a ruled surface in space of any number of dimensions, we have a curve of order  $\nu$  and genus  $\pi$  meeting each generator in two points, and having no double points which are not also double points of the ruled surface, then

$$\nu - \pi = n - 2p + 1,$$

where  $n$  is the order and  $p$  the genus of the ruled surface. If the curve, of order  $\nu$  and genus  $\pi$ , is the projection of a normal curve in higher space, the ruled surface is also the projection of a normal ruled surface in higher space\*; we have already made use of this idea to obtain the normal space for elliptic ruled surfaces.

If the ruled surface is a sextic ruled surface whose prime sections are of genus 2 we have

$$\nu - \pi = 3,$$

\* Cf. Segre, *Math. Ann.* 34 (1889), 1.

so that the curve is certainly normal in [3] if  $\nu > 2\pi - 2$ , i.e. if  $\pi < 5$  and  $\nu < 8$ . Hence, *if a ruled surface in [3], of order 6 and genus 2, is the projection of a normal ruled surface in [4], there cannot be on the surface a curve of order less than eight (and of genus less than 5) meeting each generator in two points.*

If we attempt to obtain such a curve, e.g. by means of a quadric containing at least five generators, the attempt is bound to fail. If the surface has a directrix  $R$  the quadric must not contain  $R$ , for then the residual curve of intersection would not meet each generator twice.

*The sextic ruled surfaces whose plane sections are of genus greater than 2*

**342.** A sextic ruled surface in [4] whose prime sections are of genus 3 is necessarily a cone\*; thus such a ruled surface is normal in [3] and the question of projection from higher space will not arise, nor will it arise *a fortiori* for the surfaces with prime sections of genus greater than 3. It remains then to consider such surfaces as curves on the quadric  $\Omega$ .

The sextic curve of genus 3 is necessarily contained in [3], so that the sextic ruled surface will have two directrices—distinct or coincident. Now the sextic curves of genus 3 lying on a quadric surface are of two kinds:

(A) the curve meets all generators of one system in four points and all of the other system in two points;

(B) the curve meets every generator in three points and has a double point.

In the first case we have a ruled surface with a fourfold directrix and a double directrix; the double curve is  $6R + R'$  and the bitangent developable is  $R + 6R'$ . The surface is generated by two lines  $R$  and  $R'$  placed in (2, 4) correspondence.

In the second case we have a ruled surface with two triple directrices and a double generator; the double curve and bitangent developable are both  $3R + G + 3R'$ . The surface is generated by two lines  $R$  and  $R'$  in (3, 3) correspondence, the correspondence being specialised to give the double generator.

In either case the two directrices may coincide. We can have on a quadric cone:

(A) a curve with a double point at the vertex, meeting every generator in two other points;

(B) a curve meeting every generator in three points and having a double point.

In (A) the double curve and bitangent developable are both  $7R$ ; the directrix itself being a double generator. Through any point of  $R$  there pass two other generators lying in a plane through  $R$ . The surface is

\* Cf. § 160.

generated by a line and a plane quintic in (1, 2) correspondence with a united point, the quintic having a triple point on  $R$ .

In (B) the double curve and the bitangent developable are both  $6R + G$ ; through any point of  $R$  there pass three generators lying in a plane through  $R$ . To generate this surface we take a line and a plane quartic without double points and place them in (1, 3) correspondence with a united point, the three points of the quartic which correspond to any point of the line lying on a line through the united point. The tangent to the quartic at this point is a double generator of the surface.

**343.** On a quadric in [3] we can have a sextic of genus 4; this is the complete intersection of the quadric with a cubic surface and meets every generator in three points. We can thus have in [3] a sextic ruled surface with plane sections of genus 4; the surface has two directrices  $R$  and  $R'$ , through any point of either there pass three generators lying in a plane through the other. The double curve and bitangent developable are both  $3R + 3R'$ . The surface is generated by two lines  $R$  and  $R'$  in (3, 3) correspondence.

Here also we can have a surface for which the two directrices coincide; this is represented by the sextic curve which is the intersection of a cubic surface and a quadric cone. The double curve and bitangent developable are both  $6R$ ; through any point of  $R$  there pass three generators lying in a plane through  $R$ .

The surface is generated by a plane quintic with a tacnode and a line passing through the tacnode, these being in (3, 1) correspondence with a doubly united point. The three points of the quintic corresponding to any point of the line are collinear with the tacnode, and the tangent at the tacnode is a generator of the ruled surface.

## CHAPTER V

### DEVELOPABLE SURFACES

#### *Introduction*

**344.** We have already mentioned the construct in three-dimensional space which is known as a *developable*\*. It is formed by a singly infinite system of planes and can thus be regarded as the dual of a twisted curve; for this reason many properties of developables have been known for a long time. The *class* of the developable is the number of its planes which pass through an arbitrary point.

We may regard the developable in another way. Just as the line joining two points of a curve is called a chord of the curve, so the line of intersection of two planes of a developable is called an *axis* of the developable. As the two points of the curve tend to coincide, the chord tends to a limiting position called a tangent of the curve, so, when the two planes of the developable tend to coincide, the axis tends to a limiting position called a *generator* of the developable. Just as there is a tangent at every point of the curve, so there is a generator in every plane of the developable, leaving aside for the moment the consideration of possible singular planes. We have thus  $\infty^1$  generators forming a *developable surface*. It then appears that a developable surface is merely a particular case of a ruled surface; except for a finite number of generators which are torsal the tangent planes to a ruled surface are different for different points of the same generator, but on a developable surface all the generators are torsal with the same tangent plane at different points of the same generator. We shall in future use the same term developable to denote either the singly infinite system of planes or the surface formed by the generators.

Further, just as we derive osculating planes of a curve, so we can derive a singly infinite system of points forming a curve  $\Gamma$  on the developable. If we consider the generator  $g$  in the plane  $\alpha$ , then the limiting position of the point of intersection of  $g$  with another plane  $\beta$  of the developable, as  $\beta$  approaches  $\alpha$ , is a point of  $\Gamma$ . The generators of the developable touch  $\Gamma$  and the planes of the developable osculate  $\Gamma$ ;  $\Gamma$  is called the "edge of regression" or the "cuspidal edge" of the developable.

Every developable is formed by the osculating planes of some curve  $\Gamma$ , and conversely the osculating planes of every curve  $\Gamma$  form a developable. Any numerical relation connecting singularities of a developable gives another numerical relation connecting corresponding singularities of a

\* See e.g. § 21.

twisted curve and *vice versa*; these relations have been well known since Cayley's generalisation of Plücker's formulæ for plane curves to curves in space\*.

**345.** If we project a twisted curve  $\Gamma$  on to a plane from an arbitrary point  $O$  we obtain a plane curve, and every osculating plane of  $\Gamma$  passing through  $O$  gives on the plane a tangent at a point of inflection on the plane curve; the number of these is equal to the class of the developable formed by the osculating planes of  $\Gamma$ . If  $O$  lies on a tangent of  $\Gamma$  the plane curve loses two of these inflections and acquires a cusp.

The section of a developable by an arbitrary plane  $\omega$  gives a system of lines enveloping a curve; this curve is the locus of points in which  $\omega$  is met by the generators of the developable and has cusps at the points where  $\omega$  meets the cuspidal edge. If, however,  $\omega$  contains a generator the curve loses two of these cusps and acquires an inflection; this inflection is at the point of contact of the generator in  $\omega$  with the cuspidal edge and the generator is itself the inflectional tangent. If there are  $r$  generators of the developable meeting an arbitrary line in space, the section by a plane  $\omega$  containing a generator is a curve of order  $r - 1$ ; this will meet the generator in  $r - 4$  points other than the inflection. Thus given a skew curve  $\Gamma$  of which  $r$  tangents meet an arbitrary line any one tangent is met by  $r - 4$  others. We say that  $\Gamma$  is of rank  $r$ .

We have then on a developable surface not only a cuspidal edge but a nodal curve; this nodal curve meets every generator in  $r - 4$  points if  $r$  is the order of the surface. Any plane through a generator meets the surface further in a curve of order  $r - 1$  passing through these  $r - 4$  points (which are fixed for all positions of the plane) and having an inflection at the point of contact of the generator with the cuspidal edge; but when the plane is the tangent plane the residual intersection will only be of order  $r - 2$ ; it passes through the  $r - 4$  points in which the generator meets the nodal curve and touches the generator at its point of contact with the cuspidal edge.

**346.** Suppose that we have in [3] a curve  $\Gamma$  of order  $n$  and genus  $p$ ; then it can be proved, either by means of a correspondence between the planes of a pencil or by projecting  $\Gamma$  on to a plane and using Plücker's equations, that the rank of  $\Gamma$  is  $r = 2n + 2p - 2$ . A reduction may have to be made for certain singularities of  $\Gamma$ ; in particular we must subtract 1 for each cusp.

The curves that we shall meet with will have three kinds of singularities:

- (a) cusps, or stationary points;
- (b) inflections, or points at which the tangent is stationary;
- (c) points at which the osculating plane is stationary.

The last two types of singularity do not affect the rank of the curve; so

\* *Papers*, 8 (1871), 72.

that for a curve of order  $n$  and genus  $p$  the rank is  $r = 2n + 2p - 2 - \kappa$ , where  $\kappa$  is the number of cusps of the curve.

The tangents at the inflections are stationary generators of the developable of which  $\Gamma$  is the cuspidal edge; a plane meets the developable in a curve having cusps not only at the intersections of the plane with  $\Gamma$  but also at the intersections of the plane with the tangents to  $\Gamma$  at its inflections.

Each tangent of  $\Gamma$  is met by  $r - 4$  other tangents, so that we have a nodal curve on the developable. Hence the double curve of the developable surface may consist of three parts: the cuspidal edge, the nodal curve and the tangents at the inflections of the cuspidal edge. Similarly the bitangent developable may also consist of three parts: the original developable itself, the nodal developable and the pencils of planes through the inflectional tangents of the cuspidal edge.

347. The simplest curve in [3] is the twisted cubic; the tangents of this form a developable surface of the fourth order and no other curve can give a developable of so low an order. For a developable of the fifth order the only possible cuspidal edge is the rational quartic with one cusp\*. For a developable of the sixth order the cuspidal edge must be such that

$$2n + 2p - \kappa = 8,$$

and there are three possible curves:

- (a) A rational quartic,
- (b) A rational quintic with two cusps,
- (c) A rational sextic with four cusps.

Thus all developables of the sixth order are rational, and the cuspidal edge lies on a quadric surface.

It may be remarked in passing that for an elliptic curve of order  $n$  with  $\kappa$  cusps the order of the developable of tangents is  $2n - \kappa$ . Thus the lowest order possible for an elliptic developable is eight, this developable being formed by the tangents of the curve of intersection of two quadrics. Hence all developables of order less than eight are rational, or, to use Cayley's expression, planar developables†.

348. The expression  $r = 2n + 2p - 2 - \kappa$

for the rank of a curve of order  $n$  and genus  $p$  with  $\kappa$  cusps is true whatever the dimension  $m$  of the space in which the curve lies; it gives the number of tangents of the curve which meet an arbitrary  $[m - 2]$  of general position.

If the cuspidal edge  $\Gamma$  of a developable in [3] is not normal we can consider a normal curve  $\Gamma_0$ , in higher space, of which  $\Gamma$  is the projection; the developable surface formed by the tangents of  $\Gamma$  is then the projection

\* The number of cusps of a rational curve of order  $n$  in [3] cannot exceed the integral part of  $\frac{4}{3}(n - 3)$ ; Veronese, *Math. Ann.* 19 (1882), 209.

† Cf. Schwarz, "De superficiebus in planum explicabilibus primorum septem ordinum," *Journal für Math.* 64 (1865), 1.



of the developable surface formed by the tangents of  $\Gamma_0$ , the centre of projection meeting as many tangents of  $\Gamma_0$  as there are cusps of  $\Gamma$ .

Since a rational quartic in [3] is the projection of a normal rational quartic in [4] it follows that the developable of tangents of a rational quartic in [3] is the projection of the developable of tangents of a normal rational quartic in [4]. By selecting different positions for the point of projection in [4] we are able to obtain different types of developable surfaces of the sixth order in [3], while if we take the point of projection to lie on a tangent of the normal curve we obtain in [3] the developable of the fifth order formed by the tangents of a rational quartic with a cusp.

A very complete investigation of the properties of a rational quartic in [3] as derived by projection from the normal curve is given by Marletta\*. When, however, he is examining the details of the nodal curves of some sextic developables he simply obtains the results by geometry in three dimensions; we shall try to obtain all our results systematically by projection.

For a normal rational quartic curve  $C$  in [4] the order, or the number of points of  $C$  which lie in an arbitrary solid, is 4. The first rank, or the number of tangents of  $C$  which meet an arbitrary plane, is 6; the second rank, or the number of osculating planes of  $C$  which meet an arbitrary line, is 6; the class, or third rank, or the number of osculating solids of  $C$  which pass through an arbitrary point, is 4. These numbers are well known†. If then we take the section by a solid  $\Sigma$  of the surface formed by the tangents of  $C$  we obtain a rational curve of the sixth order with cusps at the four points where  $\Sigma$  meets  $C$ . The tangents of this curve are the lines in which  $\Sigma$  meets the osculating planes of  $C$ , and they form a developable surface of the sixth order. By selecting different positions of the cutting solid  $\Sigma$  we obtain different types of sextic developables in ordinary space.

We shall give subsequently a complete list of the developables of the sixth order with their double curves, bitangent developables and cuspidal edges. Of the ten different classes which we finally obtain all but one are obtainable from the rational normal quartic curve by projection and section. The surfaces have been considered by Chasles, Schwarz and Cayley; but no complete account of them seems to have been published. •

We must mention, for completeness, two other methods of studying developables, although we shall not pursue these lines of investigation here.

The generators of a developable form a ruled surface which may be the projection of a normal ruled surface, not itself developable, in higher space. For example: all developable surfaces of the sixth order are rational and can therefore be obtained by projection from the normal rational sextic ruled surfaces in [7].

The planes of a developable may be the projections of a system of

\* *Annali di Matematica* (3), 8 (1903), 97–128.

† See e.g. Clifford, *Collected Papers* (London, 1882), 314.

planes, in higher space, forming a three-dimensional locus. For example: a rational developable of class  $n$  can always be obtained by projection from a rational normal locus, of order  $n$ , of  $\infty^1$  planes in  $[n + 2]$ .

**349.** We know how to represent the generators of a ruled surface of order  $n$  and genus  $p$  in [3] by the points of a curve  $C$  of order  $n$  and genus  $p$  lying on a quadric primal  $\Omega$  in [5]. There are  $2(n + 2p - 2)$  torsal generators of the ruled surface\*, giving  $2(n + 2p - 2)$  points of  $C$  at which the tangents lie on  $\Omega$ . Hence, since every generator of a developable surface is a torsal generator, the generators of a developable in [3] are represented on  $\Omega$  by a curve  $C$  all of whose tangents also lie on  $\Omega$ .

Suppose that we have a curve  $C$ , of order  $n$ , genus  $p$  and with  $k$  cusps, lying on a quadric primal  $\Omega$  in space of any number of dimensions. Then the number of tangents of  $C$  which lie on  $\Omega$  is found at once by considering the section of the figure by a prime  $\Pi$ . For the surface formed by the tangents of  $C$  meets  $\Pi$  in a curve  $C'$ , of order  $2n + 2p - 2 - k$ , which has cusps at the  $n$  points of intersection of  $C$  with  $\Pi$ . These cusps are all on the section  $\omega$  of  $\Omega$  by  $\Pi$ , so that  $C'$  and  $\omega$  have

$$2(2n + 2p - 2 - k) - 2n = 2(n + 2p - 2) - 2k$$

further intersections.

The tangent of  $C$  which passes through any one of these further intersections of  $C'$  and  $\omega$  lies entirely on  $\Omega$ , since it meets it in two points at its point of contact with  $C$  and in one point where it meets  $\omega$ . Conversely, a tangent of  $C$  which lies on  $\Omega$  must pass through one of these further intersections of  $C'$  and  $\omega$ ; it does not pass through a cusp of  $C'$ , since the choice of  $\Pi$  is arbitrary. Hence the number of tangents of  $C$  which lie on  $\Omega$  is  $2(n + 2p - 2) - 2k$ .

**350.** In order to obtain the properties of developable surfaces in [3] we shall make use of properties of curves on a quadric  $\Omega$  in [5], these curves being such that all their tangents also lie on  $\Omega$ . We first obtain the number of osculating planes of such a curve which lie on  $\Omega$ .

Let us suppose that  $\Omega$  is a quadric primal in space of any number of dimensions, and that on  $\Omega$  there lies a curve  $C$ , of order  $n$ , genus  $p$  and having  $k$  cusps, whose tangents all lie on  $\Omega$ . The tangents of  $C$  form a surface, and the tangent planes of this surface at the different points of a tangent of  $C$  all coincide with the corresponding osculating plane of  $C$ . But, if we have a surface on a quadric, the tangent plane of the surface at any point is contained in the tangent prime of the quadric at the same point. Hence the tangent primes of  $\Omega$ , at the different points of a tangent of  $C$ , all contain the osculating plane of  $C$ . Thus each osculating plane of  $C$  touches  $\Omega$  at every point of the tangent of  $C$ , and if it should meet  $\Omega$  in any points which are not on this tangent it will lie on  $\Omega$  completely.

\* Cf. § 32.

Consider now the section by a prime  $\Pi$ . The surface formed by the tangents of  $C$  meets  $\Pi$ , as before, in a curve  $C'$  of order  $2n + 2p - 2 - k$ , but this curve now lies entirely upon  $\omega$ . Hence, by the result just proved, the number of tangents of  $C'$  lying on  $\omega$  is

$$2 \{(2n + 2p - 2 - k) + 2p - 2\} - 2n = 2(n + 4p - 4) - 2k,$$

since  $C'$  has  $n$  cusps and is of genus  $p$ .

Now the tangents of  $C'$  are the lines in which  $\Pi$  is met by the osculating planes of  $C$ . Hence, if a tangent of  $C'$  lies on  $\omega$ , the corresponding osculating plane of  $C$  will lie on  $\Omega$ , and conversely. Therefore the number of osculating planes\* of  $C$  which lie on  $\Omega$  is  $2(n + 4p - 4) - 2k$ .

**351.** If  $\Omega$  is in [5] the points of  $C$  represent the generators of a developable in [3]; these generators all touch a curve  $\Gamma$ . A point  $P$  of  $C$  represents a generator  $g$  of the developable; through the tangent of  $C$  at  $P$  there pass two planes of  $\Omega$ , one of each system. The point of [3] which is represented on  $\Omega$  by the  $\omega$ -plane is the point of contact of  $g$  and  $\Gamma$ , while the plane of [3] which is represented on  $\Omega$  by the  $\rho$ -plane is the corresponding osculating plane of  $\Gamma$ .

A cusp of  $\Gamma$  is represented on  $\Omega$  by a  $\omega$ -plane which osculates  $C$ , and a stationary osculating plane of  $\Gamma$  is represented on  $\Omega$  by a  $\rho$ -plane which osculates  $C$ . The tangent at an inflection of  $\Gamma$  is represented on  $\Omega$  by a cusp of  $C$ .

Suppose that  $\Gamma$  has  $\kappa$  cusps and  $\kappa'$  stationary osculating planes; then the number of osculating planes of  $C$  which lie on  $\Omega$  must be  $\kappa + \kappa'$ . But, by the Cayley-Plücker formulae for a twisted curve,

$$\kappa + \kappa' = 2(n + 4p - 4) - 2i,$$

where  $\Gamma$  is of order  $n$ , genus  $p$  and has  $i$  inflections. But  $C$  is of order  $n$ , genus  $p$  and has  $i$  cusps; thus the number of osculating planes of  $C$  which lie on  $\Omega$  is in agreement with that given by § 350.

**352.** If we have a quadric primal  $\Omega$  in space of any number of dimensions, and on it a curve  $C$  whose tangents also lie on  $\Omega$ , then the tangent primes of  $\Omega$  at the different points of the tangent of  $C$  at a point  $P$  all contain the osculating plane of  $C$  at  $P$ . The tangent prime of  $\Omega$  at  $P$  itself contains the osculating solid of  $C$  at  $P$ , and so meets  $C$  in only  $n - 4$  other points, if  $n$  is the order of  $C$ . We have then, corresponding to each point  $P$  of  $C$ , a set of  $n - 4$  points which are the intersections of  $C$ , other than  $P$  itself, with the tangent prime of  $\Omega$  at  $P$ ; if we have a point  $P$  such that one of the set of  $n - 4$  points is  $P$  itself, the osculating plane of  $C$  will lie on  $\Omega$ . The tangent prime of  $\Omega$  at a point  $P$  of  $C$  meets  $C$ , in general, in four points at  $P$ ; if it should happen to meet  $C$  in five points at  $P$  then the osculating plane of  $C$  at  $P$  lies on  $\Omega$ †.

\* Cf. Baker, *Proc. Edin. Math. Soc.* (2), 1 (1927), 19.

† Baker, *loc. cit.*

*The developable surface of the fourth order*

**353.** The tangents of a twisted cubic in [3] form a developable surface of the fourth order. The section by an arbitrary plane is a rational plane quartic with three cusps. A plane through a tangent of the cubic curve meets the curve in one further point; the section of the developable consists of the tangent and a rational plane cubic. This cubic has an inflection with the generator of the developable as the inflectional tangent, and has a cusp at the remaining intersection of the plane with the twisted cubic. An osculating plane of the twisted cubic meets the developable in the tangent counted twice and a conic, the conic touching the tangent at its point of contact with the cubic curve.

No two tangents of the cubic can intersect.

Given a quartic curve  $C$  in [4] there is one quadric  $Q$  which contains  $C$  and all its tangents; if  $Q$  is regarded as a prime section of  $\Omega$  the points of  $C$  represent the generators of a developable of the fourth order in [3]. The tangents of a twisted cubic belong to a linear complex.

Any three tangents of  $C$  have one transversal; this transversal must lie on  $Q$  and  $Q$  can, in fact, be defined as the locus of transversals of triads of tangents of  $C$ . The tangents of  $C$  form a developable surface lying on  $Q$ ; this is of the sixth order and meets every line of  $Q$  in three points. A plane  $\rho$  of  $\Omega$  meets the [4] containing  $Q$  in a line which is met by three tangents of  $C$ ; this means that there are three points of the cuspidal edge of the developable in an arbitrary plane of [3]. A plane  $\omega$  of  $\Omega$  meets the [4] containing  $Q$  in a line which is met by three tangents of  $C$ ; this means that there are three planes of the developable passing through an arbitrary point of [3].

The fact that any three osculating planes of a twisted cubic meet in a point coplanar with their points of osculation is quite clear on the quadric  $\Omega$ . Take three tangents  $t_1, t_2, t_3$  of  $C$ ; then the planes  $\rho_1, \rho_2, \rho_3$  of  $\Omega$  passing through them represent three arbitrary osculating planes of the twisted cubic. The point of intersection of these three planes is represented on  $\Omega$  by the unique plane  $\omega_0$  which meets each of the planes  $\rho_1, \rho_2, \rho_3$  in lines, and this is clearly the plane  $\omega_0$  which contains the transversal  $t_0$  of  $t_1, t_2, t_3$ . Then the plane  $\rho_0$  through  $t_0$  meets the four planes  $\omega_0, \omega_1, \omega_2, \omega_3$  all in lines and so represents a plane passing through the three points where the planes osculate the twisted cubic and also through the point of intersection of the three osculating planes;  $\omega_1, \omega_2, \omega_3$  are the  $\omega$ -planes of  $\Omega$  which contain  $t_1, t_2, t_3$ .

**354.** There is a result due to Brill and Nöther\* which states that the curves of order  $n$  and genus  $p$  in [ $r$ ] depend upon

$$n(r+1) - (p-1)(r-3)$$

parameters, provided that

$$n \geq r + \frac{rp}{r+1}.$$

\* Brill and Nöther, *Math. Ann.* 7 (1874), 308. Nöther, *Journal für Math.* 93 (1882), 281. Segre, *Math. Ann.* 30 (1887), 207.

Thus in [4] there are  $\infty^{21}$  rational normal quartic curves. Through each of these curves there passes a quadric containing the curve and all its tangents; hence, since there are only  $\infty^{14}$  quadrics in [4], there must be, on any given quadric in [4],  $\infty^7$  rational normal quartic curves all of whose tangents also lie on the quadric. Hence, given a quadric primal  $\Omega$  in [5], there are  $\infty^5$  prime sections  $Q$  on each of which there are  $\infty^7$  of these quartic curves. Hence there must be  $\infty^{12}$  developable surfaces of the fourth order in ordinary space.

This is also seen at once since a developable surface is completely determined by its cuspidal edge, and Brill and Nöther's result shews that there are  $\infty^{12}$  twisted cubics in [3]\*.

*The developable surface of the fifth order*

355. The developable surface of the fifth order is formed by the tangents of a rational quartic curve, in a [3]  $\Sigma$ , having a cusp. Each tangent is met by one other tangent.

The quartic curve  $\Gamma$  is the projection of a normal curve  $C$  in [4] from a point  $O$  on one of its tangents. The tangent touches  $C$  in a point  $P$  and meets  $\Sigma$  in the cusp  $K$  of  $\Gamma$ . The developable surface of the sixth order formed by the tangents of  $C$  is projected, from  $O$ , into the developable surface of the fifth order formed by the tangents of  $\Gamma$ .

The tangent solid of  $Q$  at  $O$  contains the osculating plane of  $C$  at  $P$ , and therefore meets  $C$  in one other point  $P'$ . The osculating solid† of  $C$  at  $P'$  passes through  $O$  and meets  $\Sigma$  in a plane which has four-point contact with  $\Gamma$  at the point  $W$  which is the projection of  $P'$ . Thus there is a point  $W$  of  $\Gamma$  at which the osculating plane is stationary. The intersection of  $Q$  with its tangent solid at  $O$  is an ordinary quadric cone; the line  $OP$  is a generator of this cone, the tangent plane of the cone along this generator being the osculating plane of  $C$  at  $P$ . This plane meets  $\Sigma$  in the cuspidal tangent of  $\Gamma$  at  $K$ . The quadric cone also has  $OP'$  as a generator. Hence the intersection of  $\Sigma$  with this quadric cone is a conic touching the cuspidal tangent of  $\Gamma$  at  $K$  and passing through  $W$ . The tangent of this conic at  $W$  lies in the stationary osculating plane of  $\Gamma$ .

This conic is the nodal curve of the developable surface of the fifth order, for every generator of the quadric cone on  $Q$  meets three tangents of  $C$ , and therefore meets two tangents other than  $OP$ . Conversely, every

\* For this particular result see also Reye, *Die Geometrie der Lage* (2) (Stuttgart, 1907), 200.

† If any point of [4] is taken there are four osculating solids of  $C$  passing through it; the four points in which they osculate  $C$  are the intersections of  $C$  with the polar solid of the point in regard to  $Q$  (Clifford, *Papers*, 312–313). For a point  $O$  on  $Q$  the polar solid is the tangent solid at  $O$ ; if  $O$  is on the tangent at a point  $P$  of  $C$  the four points of intersection of the solid with  $C$  consist of  $P$  counted three times and  $P'$ . The osculating solids of  $C$  are the tangent solids of  $Q$  at the points of  $C$ .

line through  $O$  meeting two further tangents of  $C$  must lie on  $Q$  and therefore in its tangent solid at  $O$ .

The pairs of tangents of  $C$  which meet the lines of  $Q$  passing through  $O$  touch  $C$  in pairs of points belonging to an involution. The double points of this involution are  $P$  and  $P'$ . The joins of the pairs of points form a cubic ruled surface\*, and the solids which contain the pairs of tangents of  $C$  are the tangent solids of this ruled surface; they touch a quadric line-cone whose vertex is the directrix of the surface, this directrix passing through  $O$ . But these solids meet  $\Sigma$  in planes, each of which contains two tangents of  $\Gamma$ ; hence the nodal developable consists of the tangent planes of a quadric cone. The osculating planes of  $\Gamma$  at  $K$  and  $W$  are tangent planes of this cone. The generators of the cone are the intersections of  $\Sigma$  with the planes joining  $O$  to the generators of the cubic ruled surface; the tangent of  $\Gamma$  at  $W$  is a generator of the cone, and there is a generator passing through  $K$ .

**356.** Given a rational normal quintic curve  $C$  in [5] there are three linearly independent quadric fourfolds which contain  $C$  and all its tangents. There are  $\infty^9$  linearly independent quadrics containing  $C$ , and each such quadric contains six tangents of  $C$ . The tangents of  $C$  form a developable surface of order eight on which  $C$  is a cuspidal curve; if then a quadric is made to contain  $C$  and seven tangents it will necessarily contain the developable surface. Thus of the  $\infty^9$  quadrics containing  $C$  there are  $\infty^2$  containing all its tangents.

Conversely, given a quadric  $\Omega$  in [5] there exist rational normal quintic curves on it such that all their tangents also lie on  $\Omega$ . The rational normal quintic curves  $C$  in [5] are  $\infty^{32}$  in aggregate, whereas quadrics are  $\infty^{20}$  in aggregate; hence, as there are for each curve  $\infty^2$  quadrics containing its tangents, there must be on a given quadric  $\infty^{14}$  curves  $C$  all of whose tangents also lie on the quadric.

**357.** Suppose then that we have on the quadric  $\Omega$  in [5] a normal rational quintic curve  $C$  all of whose tangents also lie on  $\Omega$ . If the points of  $\Omega$  represent the lines of a space  $S_3$  the points of  $C$  represent the generators of a developable surface of the fifth order.

The tangents of  $C$  form a developable surface  $R_2^8$  of the eighth order lying on  $\Omega$ ; hence the ruled surface formed by the chords of  $C$  which lie on  $\Omega$ , in all of order 12, breaks up into  $R_2^8$  and a quartic ruled surface  $R_2^4$ . There is one † generator of  $R_2^4$  through each point of  $C$ , and  $R_2^4$  is formed by the joins of pairs of an involution on  $C$ . This involution has two double points  $x$  and  $y$ ; the tangents of  $C$  at  $x$  and  $y$  are common generators of  $R_2^4$  and  $R_2^8$ .

\* See the footnote to § 19.

† See § 352.

The tangent prime of  $\Omega$  at any point  $P$  of  $C$  meets  $C$  in four points at  $P$  and in another point  $P'$ , the line  $PP'$  is the generator of  $R_2^4$  through  $P$ . The tangent prime of  $\Omega$  at  $P'$  meets  $C$  four times at  $P'$ , the remaining intersection of  $C$  with this tangent prime being  $P$ . But at  $x$  or  $y$  the tangent prime of  $\Omega$  has five intersections with  $C$  and does not meet it elsewhere; the osculating planes of  $C$  at  $x$  and  $y$  lie on  $\Omega^*$ . We must then regard these as the two trisecant planes of  $C$  lying on  $\Omega$ ; they are of opposite systems†.

The tangent plane of  $R_2^4$  at a point  $P$  of  $C$  is the plane containing the tangent of  $C$  at  $P$  and the generator  $PP'$  of  $R_2^4$ . Thus at  $x$  or  $y$  the tangent plane of  $R_2^4$  is the osculating plane of  $C$ . The plane of the directrix conic of  $R_2^4$ , which passes through  $x$ , thus meets  $\Omega$  in the directrix conic and also in its tangent at  $x$ , and so lies on  $\Omega$  entirely; similarly the plane of the directrix conic of  $R_2^4$  which passes through  $y$  also lies on  $\Omega$  entirely. The planes of the conics on  $R_2^4$  form a  $V_3^3$  whose complete intersection with  $\Omega$  consists of  $R_2^4$  and these two planes.

Suppose then that  $\Omega$  contains  $\omega_x$  (the osculating plane of  $C$  at  $x$ ),  $\rho_y$  (the osculating plane of  $C$  at  $y$ ),  $\rho_x$  (the plane of the directrix conic of  $R_2^4$  which passes through  $x$ ) and  $\omega_y$  (the plane of the directrix conic of  $R_2^4$  which passes through  $y$ ). The osculating plane of  $C$  at  $x$  or  $y$  must be of the opposite system to the plane of the directrix conic of  $R_2^4$  passing through that point since they have in common a tangent of this conic.

Any plane of  $\Omega$  of general position meets  $R_2^8$  in four points and  $R_2^4$  in two points. The developable surface in  $S_3$  is formed by the tangents of a rational quartic  $\Gamma$  of which there are four osculating planes passing through an arbitrary point of  $S_3$ .  $\Gamma$  has a cusp, represented on  $\Omega$  by  $\omega_x$ , and a stationary osculating plane represented on  $\Omega$  by  $\rho_y$ . The nodal curve is a conic whose plane is represented on  $\Omega$  by  $\rho_x$  and the nodal developable is a quadric cone whose vertex is represented on  $\Omega$  by  $\omega_y$ .

**358.** We have seen that on a quadric  $\Omega$  in [5] there are  $\infty^{14}$  rational normal quintic curves all of whose tangents lie on  $\Omega$ . In other words the quintic developables in  $S_3$  are  $\infty^{14}$  in aggregate. This can be verified in other ways. For rational quartics in  $S_3$  are  $\infty^{16}$  in aggregate and two conditions are necessary for such a curve to have a cusp. If, for example, the curve is regarded as the projection of a normal curve in [4], the point of projection must lie on the surface formed by the tangents of the normal curve, and this imposes two conditions on the point. Hence rational quartics in  $S_3$  which have cusps are  $\infty^{14}$  in aggregate, and since a developable is completely determined by its cuspidal edge the result is verified.

Again, any rational quartic in  $S_3$  which has a cusp is the intersection of a pencil of quadrics having stationary contact. Now the quadrics in  $S_3$  are  $\infty^9$  in aggregate, so that pencils of quadrics are  $\infty^{16}$  in aggregate,

\* Cf. § 350.

† Cf. § 88.

just as there are  $\infty^{16}$  lines in [9]. Then pencils of quadrics which touch are  $\infty^{15}$ , while pencils of quadrics having stationary contact are  $\infty^{14}$ ; two conditions being necessary for two quadrics to have stationary contact\*.

*The projections of the developable formed by the tangents of a normal rational quartic curve*

**359.** Any surface in [4] can be projected from a point  $O$  on to a solid  $\Sigma$ . There are  $\infty^1$  chords of the surface passing through  $O$ , and these form a cone of lines meeting the surface in a curve†. Two points of this curve which are collinear with  $O$  may be called associated points; the tangents to the curve at a pair of associated points intersect, their plane passing through  $O$ ; the osculating planes of the curve at a pair of associated points have a line in common, the solid containing them passing through  $O$ .

We can of course project ruled surfaces in this way.

Consider, in particular, the developable formed by the tangents of a curve  $C$  in [4]; we have a cone of lines through  $O$ , each line meeting two generators of the developable. These lines meet the developable in points lying on a certain curve; the points of this curve lie on the tangents of  $C$ , while the tangents of the curve lie in the osculating solids of  $C$ .

**360.** Consider now a rational normal quartic curve  $C$  in [4]; its tangents form a rational developable  $F$  of order '6. There are  $\infty^5$  quadric threefolds containing  $C$ , each of these containing four tangents of  $C$ ; there is a unique quadric  $Q$  containing  $F$ . Let us project  $F$  on to a solid  $\Sigma$  from a point  $O$ .

There are  $\infty^1$  chords of  $F$  passing through  $O$ ; a solid through  $O$  meets  $F$  in a rational sextic with four cusps and six apparent double points; thus the chords of  $F$  passing through  $O$  meet  $F$  in a curve  $C_{12}$  of order 12.

The plane joining  $O$  to a tangent of  $C$  is met by two other tangents; for the projection of  $C$  from one of its tangents on to a plane is a conic, and there are two tangents of this conic passing through any point of the plane. Hence  $C_{12}$  meets every generator of  $F$  in two points.

**361.** There are four points of  $C$  for which the osculating solids contain  $O$ ; these are, in fact, the intersections of  $C$  with the polar solid of  $O$  in regard to  $Q$ ‡. Take one of these points  $W$ , and the tangent  $w$  of  $C$  at  $W$ .

The projection of  $C$  from  $w$  on to a plane is a conic; the osculating solid at  $W$  meets the plane in a tangent of the conic passing through the pro-

\* Salmon, *Geometry of Three Dimensions*, 1 (Dublin, 1914), 208.

The quintic developable is studied algebraically by Cayley, *Papers*, 1 (1850), 491; 2 (1864), 275 and 518; Schwarz, *Journal für Math.* 64 (1865), 4-9; Dino, *Giornale di Matematiche* (1), 3 (1865), 100 and 133. For geometrical treatment see Chasles, *Comptes Rendus*, 54 (1862), 322 and 719; Cremona, *ibid.* 604-608; d'Ovidio, *Giorn. di Mat.* (1), 3 (1865), 107, 184 and 214.

† Lines through a point of [4] are  $\infty^3$  and one condition is necessary for a line to meet a surface in [4].

‡ See the footnote to § 355.



jection of  $O$ ; there is one other tangent passing through the projection of  $O$ . Hence the plane  $wO$  only meets one tangent of  $C$  other than  $w$ ; suppose that it meets this other tangent in a point  $A$ . Then  $OA$  meets  $w$  in a point  $B$ , and  $A$  and  $B$  are a pair of associated points on  $C_{12}$ . The tangents of  $C_{12}$  at  $A$  and  $B$  intersect and they both lie in the osculating solid of  $C$  at  $W$ .

This osculating solid of  $C$  is the tangent solid of  $Q$  at  $W$ ; it meets  $Q$  in a quadric cone whose vertex is  $W$  and of which  $WA$  and  $WB$  are generators. The osculating plane of  $C$  at  $W$  is common to the tangent solids of  $Q$  at all the points of  $w$  and touches the cone along  $w^*$ . The solid meets  $F$  in  $w$  counted three times and in a cubic curve lying on the cone; this cubic touches  $w$  at  $W$  and passes through  $A$ . There is one chord of this cubic passing through  $O$ , and we thus have two points of  $C_{12}$ ; the solid meets  $C_{12}$  also in  $A$  and  $B$ , touching it at each of these points; the remaining intersections of the solid with  $C_{12}$  consist simply of  $W$  counted six times. The six chords of  $C_{12}$  lying in the osculating solid at  $W$  consist of the chord of the cubic, the line  $OAB$  counted twice and the line  $OW$  counted three times. This solid meets  $\Sigma$  in an osculating plane of the nodal curve.

Thus on projection we have in  $\Sigma$  a rational quartic curve  $C_4$ ; the tangents of  $C_4$  form a developable of the sixth order with a nodal curve  $C_6$  of the sixth order. There are four points  $W_1, W_2, W_3, W_4$  of  $C_4$  at which the osculating plane is stationary; the nodal curve passes through these four points and has there the same osculating planes as  $C_4$ . The tangent of  $C_4$  at any one of these points meets  $C_6$  in one further point, the corresponding stationary plane touching  $C_6$  at this point†.

**362.** There are four planes through  $O$  which contain tangents of  $C$  and meet  $C$  again‡. Consider then a plane through  $O$  containing a tangent  $t$  which touches  $C$  in  $B$  and meeting  $C$  again in a point  $S$ . There are no tangents of  $C$  meeting this plane other than  $t$  and the tangent at  $S$ .  $OS$  meets  $t$  in a point  $T$  and is a stationary generator of the cone of chords of  $F$ ; it will meet  $\Sigma$  in a point which is a cusp of the nodal curve  $C_6$ .  $S$  and  $T$  are the only points of  $C_{12}$  in the plane  $Ot$ .

Consider the solid joining  $O$  to the osculating plane of  $C$  at  $B$ . It meets  $Q$  in a quadric cone whose vertex is somewhere on  $t$ , while it meets  $F$  in  $t$  counted twice and a rational quartic, this quartic touching  $t$  at  $B$  and having a cusp at  $S$ . There are two chords of this quartic passing through  $O$ ; the remaining four chords of  $C_{12}$  which lie in the solid must then all coincide with  $OST$ . The plane in which the solid meets  $\Sigma$  will thus have four intersections with the nodal curve at its cusp, and is therefore the osculating plane of the nodal curve there.

\* Cf. Baker, *Principles of Geometry*, 4 (Cambridge, 1925), 38. Also § 350 above.

† Cf. Cremona, *Annali di Matematica* (1), 4 (1861), 92.

‡ The trisecant planes of  $C$  passing through  $O$  cut out a  $g_3^1$  on  $C$ , being one system of planes of a quadric point-cone. This  $g_3^1$  has four double points.

Hence  $C_4$  has four tangents which touch it in points  $B_1, B_2, B_3, B_4$  and meet it again in points  $S_1, S_2, S_3, S_4$ . The nodal curve  $C_6$  has four cusps at the points  $S$ ; the osculating plane of  $C_6$  at a point  $S$  osculating  $C_4$  at the corresponding point  $B$ .

**363.** In the preceding work we have projected from a general point  $O$  of the [4] containing  $C$ . Suppose now that we specialise the position of  $O$  so that it lies on the quadric  $Q$ . Then the polar solid of  $O$  in regard to  $Q$  passes through  $O$ , being the tangent solid of  $Q$  at  $O$ . Hence we obtain in  $\Sigma$  a rational quartic  $C_4$  whose four points of superosculation  $W_1, W_2, W_3, W_4$  are coplanar.

Any line of  $Q$  is met by three tangents of  $C$ , and in fact  $Q$  may be defined as the locus of transversals of triads of tangents of  $C$ . Thus the tangents of  $C_4$  intersect in threes in the points of a conic\*; this conic is the intersection of  $\Sigma$  with the quadric cone in which  $Q$  is met by its tangent solid at  $O$ , and it passes through  $W_1, W_2, W_3, W_4$ .

Hence we have a developable whose nodal curve degenerates into a triple conic; this conic meets the cuspidal edge in the four (coplanar) points at which its osculating plane is stationary.

**364.** Now let us choose  $O$  to lie in an osculating plane  $\alpha$  of  $C$ ;  $\alpha$  contains a tangent  $t$  of  $C$  touching it at a point  $T$ . It is common to the tangent solids of  $Q$  at all the points of  $t$ , and every line in  $\alpha$  touches both  $Q$  and  $F$  at the point where it meets  $t$ .

• A solid through  $O$  meets  $F$  in a rational sextic with four cusps; there is a tangent of this curve passing through  $O$ , viz. the line in which the solid meets  $\alpha$ . Then there must be five chords of the curve passing through  $O$ ; so that the chords of  $F$  through  $O$  give on  $F$  a curve  $C_{10}$  of order 10, and the nodal curve of the developable in  $\Sigma$  is a curve  $C_5$  of the fifth order.

The polar solid of  $O$  in regard to  $Q$  contains  $t$  and therefore meets  $C$  in two other points. Hence when  $C$  is projected on to  $\Sigma$  from  $O$  it becomes a rational quartic  $C_4$  with an inflection  $I$  and two points  $W_1, W_2$  at which the osculating plane is stationary. There are only two planes through  $O$  which contain tangents of  $C$  and meet it again†, so that there are two tangents of  $C_4$  which touch it at points  $B_1, B_2$  and meet it again in points  $S_1, S_2$ .

By repeating previous arguments we see that the nodal curve  $C_5$  passes through  $W_1$  and  $W_2$  and has there the same osculating planes as  $C_4$ . Further,  $C_5$  meets  $C_4$  in  $S_1$  and  $S_2$ ;  $C_5$  has, in fact, cusps at  $S_1$  and  $S_2$ , the osculating planes of  $C_5$  at these points osculating  $C_4$  in  $B_1$  and  $B_2$ .

\* See R. A. Roberts, "Unicursal Twisted Quartics," *Proc. Lond. Math. Soc.* (1), 14 (1883), 22, and Marletta, *Annali di Matematica* (3), 8 (1903), 109.

† The  $g_3^1$  given by the trisecant planes through  $O$  includes a set for which all the three points coincide; this triple point counts for two double points. In general, an  $m$ -ple point of a  $g_n^1$  counts for  $m - 1$  double points.

365. If we consider now any solid belonging to the pencil of solids through  $\alpha$ , it meets  $Q$  in a quadric cone whose vertex lies on  $t$  and which touches  $\alpha$  along its generator  $t$ . The same solid meets  $F$  in a quartic having a cusp (at the one intersection other than  $T$  of the solid with  $C$ ) and touching  $t$  at  $T$ . There are two chords of this quartic passing through  $O$ ; the other three chords of  $C_{10}$  which lie in this solid will all coincide with  $OT$ . Thus the nodal curve  $C_5$  of the developable in  $\Sigma$  passes through  $I$ , and any plane of the pencil determined by the inflectional tangent of  $C_4$  at  $I$  meets  $C_5$  in three points at  $I$ .

Similarly we can shew that a plane through  $I$  which does not contain the inflectional tangent of  $C_4$  meets  $C_5$  in only one point there; a solid containing  $OT$  but not  $t$  meets  $F$  in a rational sextic with four cusps, one of which is at  $T$ , the cuspidal tangent of this curve at  $T$  passing through  $O$ .

Thus we conclude that  $C_5$  has an inflection at  $I$  with the same inflectional tangent as  $C_4$ .

366. We now choose  $O$  to be the intersection of two osculating planes of  $C^*$ . Then on projection we have in  $\Sigma$  a rational quartic  $C_4$  with two inflections  $I$  and  $J$ . The developable of tangents has a nodal curve  $D_4$  which is also a quartic;  $D_4$  has inflections at  $I$  and  $J$  with the same inflectional tangents as  $C_4$ .

367. Finally, let us choose  $O$  to lie on a chord  $XY$  of  $C$ . The chords of  $C$  form a locus  $M_3^3$ , and through any point of  $M_3^3$  there pass two lines lying on it which are not chords of  $C$ . Such lines we shall call axes of  $C$ . The chords of  $C$  which meet an axis form a cubic ruled surface†.

Through  $O$  there pass two axes  $m$  and  $n$  of  $C$ . The cubic ruled surfaces determined by  $m$  and  $n$  have the chord  $XY$  as a common generator, and the solid which contains the tangents of  $C$  at  $X$  and  $Y$  touches both the cubic ruled surfaces along this generator and contains  $m$  and  $n$ . When we project from  $O$  on to  $\Sigma$  we obtain a rational quartic  $C_4$  with a double point; the lines  $m$  and  $n$  meet  $\Sigma$  in points  $A$  and  $B$  which are the vertices of quadric cones containing  $C_4$ ,  $A$  and  $B$  both lying in the plane of the tangents at the double point.

Now take one of the axes through  $O$ , say  $m$ . Then through any point of  $m$  there passes a chord of  $C$  which is a generator of the cubic ruled surface containing  $m$ . The solid containing the two tangents of  $C$  at the points where this chord meets it is the tangent solid of the ruled surface along this generator; it therefore contains  $m$ . Hence there is a transversal from  $O$  to these two tangents.

\* The locus of intersections of pairs of osculating planes of  $C$  is a quartic surface. See § 368.

† See Segre, "Sulle varietà cubiche dello spazio a quattro dimensioni," *Memorie Torino* (2), 39 (1889); in particular p. 35.

Thus either cubic ruled surface gives a cone through  $O$  formed by chords of the developable of tangents of  $C$ ; the nodal curve of the developable in  $\Sigma$  breaks up into two parts.

There are two tangents of  $C$  meeting  $m$ ; these touch  $C$  in two points, the osculating solids at which contain  $m$ . There are generators of the cone through  $O$  which meet  $C$  in these two points.

The line  $OXY$  is a stationary generator of the cone, meeting a pair of tangents of  $C$  at the points where they themselves touch  $C$ . The tangent plane of the cone along this stationary generator will lie in the solid joining  $O$  to the osculating plane of  $C$  at  $X$  and also in the solid joining  $O$  to the osculating plane of  $C$  at  $Y$ .

Now the line  $m$  is not a generator of the cone of chords through  $O$  which it determines; but nevertheless it meets two tangents of  $C$ . Hence it must be a generator of the cone of chords determined by  $n$ . The tangent plane of this latter cone along  $m$  is the intersection of the two solids joining  $O$  to the planes which osculate  $C$  at the points where it is touched by the two tangents meeting  $m$ ; but these are none other than the osculating solids of  $C$  at these two points. Thus we have in  $\Sigma$  an inflection on the nodal curve.

Thus the developable of tangents of  $C_4$  has a nodal curve consisting of two plane cubics  $C_3$  and  $D_3^*$ . Each of these cubics has a cusp at the double point of  $C_4$ , the cuspidal tangent being for each of them the intersection of the two osculating planes of  $C_4$  at this point.  $C_3$  meets  $C_4$  also in two points  $W_1, W_2$ , at each of which the osculating plane of  $C_4$  is stationary. Similarly  $D_3$  meets  $C_4$  also in two points  $W_3, W_4$ , at each of which the osculating plane of  $C_4$  is stationary.  $C_3$  has an inflection at  $B$ , the inflectional tangent being the intersection of the osculating planes of  $C_4$  at  $W_3$  and  $W_4$ , while  $D_3$  has an inflection at  $A$ , the inflectional tangent being the intersection of the osculating planes of  $C_4$  at  $W_1$  and  $W_2$ †.

The tangent solids of a cubic ruled surface in [4] all pass through its directrix; they meet any solid in the tangent planes of a quadric cone. Thus the two quadric cones with vertices  $A$  and  $B$  which contain  $C_4$  belong to the bitangent developable of the surface.

*The sections of the locus formed by the osculating planes  
of a normal rational quartic curve*

**368.** Instead of projecting the points of  $C$  from a point  $O$  (which does not lie on a tangent of  $C$ ) we can take the sections of the osculating solids of  $C$  by a solid  $\Sigma$  (which does not contain an osculating plane

\* Cf. Chasles, *Comptes Rendus*, 54 (1862), 718.

† Cf. Brambilla, *Rendiconti della Reale Acc. di Scienze Fis. e Mat. di Napoli*, 24 (1885), 294.

of  $C$ ). These sections are planes which osculate a rational sextic  $C_6$  with four cusps, and the tangents of this curve, which are the lines in which the osculating planes of  $C$  meet  $\Sigma$ , form a developable of the sixth order. The cusps of  $C_6$  are the four points  $K_1, K_2, K_3, K_4$  in which  $\Sigma$  meets  $C$ .

The nodal curve of this developable in  $\Sigma$  is given by pairs of non-consecutive intersecting osculating planes of  $C$ ; hence its order is equal to the number of points of an arbitrary plane through which pass two osculating planes but not a tangent of  $C$ . The osculating planes meet the arbitrary plane in the points of a rational sextic with six cusps at the points where the plane is met by tangents of  $C$ ; this curve has four double points. Hence there are four points of a given plane at which two distinct osculating planes of  $C$  intersect, so that the developable in  $\Sigma$  has a nodal curve  $C_4$  of the fourth order.

**369.** If a plane meets  $C$  in a point  $K$  then there are four tangents of  $C$  which meet the plane in points other than  $K$ ; for when  $C$  is projected from  $K$  on to a solid it becomes a twisted cubic, of which there are four tangents meeting an arbitrary line.

Also, if a line meets  $C$  in a point  $K$  it is met by three osculating planes of  $C$  in points other than  $K$ ; for when  $C$  is projected from the line on to a plane it becomes a rational cubic which has three inflections.

Thus a plane which meets  $C$  in a point  $K$  is met by the osculating planes of  $C$  in the points of a rational sextic curve with a triple point at  $K$  and four cusps. This curve will have three other double points. We deduce that the nodal curve  $C_4$  of the developable meets the cuspidal edge  $C_6$  in its four cusps  $K_1, K_2, K_3, K_4$ .

But we can prove more than this, for the curves  $C_4$  and  $C_6$  have the same osculating planes at  $K_1, K_2, K_3, K_4$ . The osculating plane of  $C_6$  at  $K_1$  is the intersection of  $\Sigma$  with the osculating solid of  $C$  at  $K_1$ . This solid meets the developable  $F$  formed by the tangents of  $C$  in the tangent at  $K_1$  counted three times and a twisted cubic touching this tangent at  $K_1^*$ ; the osculating plane of  $C$  at  $K_1$  being also the osculating plane of this cubic curve. Then the plane of intersection of the osculating solid with  $\Sigma$  meets this osculating plane in a line  $p$ ; it meets the other osculating planes of  $C$  in the points of a rational quartic with three cusps, one of these cusps is at  $K_1$  and  $p$  is the cuspidal tangent there. The remaining intersection of  $p$  with the quartic curve is the only point of the plane in which two distinct osculating planes of  $C$  intersect. Hence the plane osculates  $C_4$  as well as  $C_6$ .

**370.** Instead of choosing  $\Sigma$  to be a general solid we can choose it to be a tangent solid of the quadric  $Q$ . Then the tangents of  $C$  meet  $\Sigma$  in the

\* Cf. § 361.

points of a rational sextic curve  $C_6$  which has four cusps and lies on a quadric cone. It meets every generator of the cone in three points; the tangent planes of the cone are tritangent planes of the developable.

**371.** Now choose  $\Sigma$  to contain a tangent  $t$  of  $C$ ; it meets  $C$  in two other points  $K_1, K_2$ . Then the tangents of  $C$  other than  $t$  meet  $\Sigma$  in the points of a rational quintic curve  $C_5$  with cusps at  $K_1$  and  $K_2$ . The osculating planes of  $C$  meet  $\Sigma$  in the tangents of  $C_5$ , so that  $t$  will be a stationary tangent of  $C_5$ .  $C_5$  then has an inflection at  $I$ , the point of contact of  $t$  with  $C$ ,  $t$  being the inflectional tangent.

Using similar arguments to those above we find that the tangents of  $C_5$  form a developable surface with a nodal quartic curve  $C_4$ ;  $C_4$  passes through  $K_1$  and  $K_2$  and has at these points the same osculating planes as  $C_5$ .

Any line meeting a tangent of  $C$  will meet four other osculating planes of  $C$ ; for when  $C$  is projected from the line on to a plane we obtain a rational quartic, with one cusp and two double points, which has four inflections. Hence a plane through  $t$  meets the locus formed by the osculating planes of  $C$  in  $t$  counted twice and a rational quartic with cusps at the two other points in which the plane is met by tangents of  $C^*$ . This quartic will have one other double point.

Hence any plane of  $\Sigma$  passing through  $t$  meets  $C_4$  in only one point not on  $t$ . In fact  $C_4$  has an inflection at  $I$  with  $t$  as the inflectional tangent.

**372.** If we take  $\Sigma$  to be the solid containing the tangents at two points  $I$  and  $J$  of  $C$  it meets the developable of tangents further in a rational quartic curve  $C_4$  with inflections at  $I$  and  $J$ . The tangents of  $C_4$  form a developable of the sixth order; this has a nodal curve which is also a quartic, having the same two inflections and inflectional tangents as  $C_4$ . We obtain the same surface as in § 366.

**373.** Finally, choose  $\Sigma$  to contain the plane in which two osculating solids of  $C$  intersect. Then the developable is formed by the tangents of a rational sextic with four cusps and a plane which osculates it in two different points. It is the dual of the developable formed by the tangents of a rational quartic with a double point.

We have already seen (in § 368) that the locus of the intersections of pairs of osculating planes of  $C$  is a quartic surface. If we project the quartic  $C$  on to a plane from a line which lies in one of its osculating planes we obtain a plane quartic with a triple point, the three branches at the triple point having a common tangent which meets the curve in four points there. This tangent is the line of intersection of the plane with the

\* Cf. § 360.

osculating solid of  $C$ . Such a plane quartic has two inflections\*, so that a line lying in an osculating plane of  $C$  meets two other osculating planes of  $C$ , or the osculating planes of  $C$  meet the quartic surface in conics.

Consider now an involution of pairs of points on  $C$ ; the lines joining the pairs form a cubic ruled surface, and the tangent solids of this cubic ruled surface envelop a quadric line-cone. Now, if we reciprocate in regard to the quadric  $Q$  which contains the tangents of  $C$ , a tangent solid of the ruled surface, as containing two tangents of  $C$ , becomes the point of intersection of two osculating planes of  $C$ , and the quadric line-cone becomes a conic. Hence, if we take any involution of pairs of points on  $C$ , the pairs of osculating planes at these points intersect in the points of a conic. Hence, since there are  $\infty^2$  involutions on  $C$ , the quartic surface has  $\infty^2$  conics upon it. The locus of intersections of pairs of osculating planes of  $C$  is therefore a projection of Veronese's surface†.

The conics in which two different osculating planes of  $C$  meet the quartic surface have one intersection, this being the intersection of the two osculating planes. The plane of the two tangents to the conics at their intersection is the tangent plane of the surface; it is in fact the plane of intersection of the two corresponding osculating solids of  $C$ , an osculating solid of  $C$  containing the tangent planes of the surface at all the points of the conic in the osculating plane. Now every solid through a tangent plane of the surface meets it in two conics‡. Hence  $\Sigma$  meets the surface in two conics; these make up the nodal curve of the developable. The conics have one intersection; this lies in the plane which osculates the sextic at two different points. Each conic passes through two of the cusps of the sextic.

If the curve  $C$  is given by

$$x_0 : x_1 : x_2 : x_3 : x_4 = \theta^4 : \theta^3 : \theta^2 : \theta : 1,$$

then the osculating planes at the two points whose parameters are  $\theta$  and  $\phi$  intersect in the point

$$\theta^2\phi^2, \quad \frac{1}{2}\theta\phi(\theta + \phi), \quad \frac{1}{6}(\theta^2 + 4\theta\phi + \phi^2), \quad \frac{1}{2}(\theta + \phi), \quad 1.$$

If either  $\theta$  or  $\phi$  is constant the locus of this point is a conic.

The coordinates of the point may be written, putting  $\theta + \phi = x$  and  $\theta\phi = y$ ,

$$y^2, \quad \frac{1}{2}xy, \quad \frac{1}{6}(x^2 + 2y), \quad \frac{1}{2}x, \quad 1,$$

so that prime sections of the surface are represented by conics in a plane. The surface is of the fourth order.

\* The Hessian of the quartic is made up of the tangent at the triple point, counted four times, and two other lines through the triple point; the remaining intersections of these two lines with the curve are the two inflections.

† A surface, in space of any number of dimensions, which contains  $\infty^2$  conics, is the surface of Veronese or one of its projections; see Bertini, *Geometria proiettiva degli iperspazi* (Messina, 1923), 393.

‡ Cf. Bertini, *ibid.* 412.

If there is a relation  $ax + by + c = 0$  we obtain the points of a conic on the surface, so that the intersections of osculating planes at pairs of points of an involution lie on a conic.

The tangent plane to the surface is the plane of the three points

$$\begin{array}{lllll} \theta^2\phi^2, & \frac{1}{2}\theta\phi(\theta + \phi), & \frac{1}{8}(\theta^2 + 4\theta\phi + \phi^2), & \frac{1}{2}(\theta + \phi), & 1; \\ 2\theta\phi^2, & \frac{1}{2}\phi(2\theta + \phi), & \frac{1}{8}(\theta + 2\phi), & \frac{1}{2}, & 0; \\ 2\theta^2\phi, & \frac{1}{2}\theta(\theta + 2\phi), & \frac{1}{8}(2\theta + \phi), & \frac{1}{2}, & 0; \end{array}$$

and is therefore common to the two solids

$$\begin{aligned} x_0 - 4\theta x_1 + 6\theta^2 x_2 - 4\theta^3 x_3 + \theta^4 x_4 &= 0, \\ x_0 - 4\phi x_1 + 6\phi^2 x_2 - 4\phi^3 x_3 + \phi^4 x_4 &= 0, \end{aligned}$$

which are the osculating solids of  $C$  at the points whose parameters are  $\theta$  and  $\phi$ .

Any solid containing a tangent plane of the surface therefore has an equation

$$\begin{aligned} \lambda(x_0 - 4ax_1 + 6a^2x_2 - 4a^3x_3 + a^4x_4) \\ + \mu(x_0 - 4\beta x_1 + 6\beta^2x_2 - 4\beta^3x_3 + \beta^4x_4) = 0; \end{aligned}$$

and if we substitute in this equation the coordinates of the point on the surface we obtain

$$\lambda(\theta - a)^2(\phi - a)^2 + \mu(\theta - \beta)^2(\phi - \beta)^2 = 0,$$

which splits up into factors

$$(\theta - a)(\phi - a) \pm k(\theta - \beta)(\phi - \beta) = 0.$$

Hence any solid through a tangent plane meets the surface in *two* curves, each curve being given by a relation of the form

$$a\theta\phi + b(\theta + \phi) + c = 0$$

between  $\theta$  and  $\phi$ . The curves are therefore conics.

*The developables of the sixth order considered as curves on  $\Omega$*

374. If we have a rational developable of the sixth order in [3] its generators are represented by the points of a rational sextic curve  $C$  lying on the quadric  $\Omega$  in [5], all the tangents of  $C$  also lying on  $\Omega$ . If, for the moment, we assume that the edge of regression  $\Gamma$  has no inflections,  $C$  will have no cusps.

In [5] the rational sextics  $C$  are  $\infty^{38}$  in aggregate\*, while the quadrics  $\Omega$  are  $\infty^{20}$  in aggregate. Since each sextic lies on  $\infty^7$  quadrics we deduce that each quadric contains  $\infty^{25}$  sextics.

Now if one of the sextic curves lies on a quadric there are eight of its tangents which lie on the quadric†; the tangents form a surface of order ten on which the curve is cuspidal, and the eight tangents, taken with the curve  $C$  counted twice, make up the complete intersection of the surface with the quadric. If a quadric contains  $C$  and nine of its tangents it will contain all the tangents. Given the curve  $C$  there will not, in general, be a quadric containing  $C$  and all its tangents.

But on any given quadric  $\Omega$  there will be  $\infty^{25-9}$  or  $\infty^{16}$  curves  $C$  all of whose tangents lie on  $\Omega$ . There are thus  $\infty^{38}$  curves  $C$  such that quadrics can be taken to contain all their tangents, and a rational sextic curve in

\* § 354.

† § 349.



[5] will have to satisfy two conditions in order that all its tangents should lie on a quadric.

We conclude that in [3] the aggregate of rational sextic developables without stationary generators is  $\infty^{16}$ . This is in accordance with what we should expect, a developable in [3] being completely defined by its cuspidal edge. For rational quartics in [3] are  $\infty^{16}$ ; rational quintics with two cusps are  $\infty^{20-2\cdot 2} = \infty^{16}$  and rational sextics with four cusps are  $\infty^{24-4\cdot 2} = \infty^{16}$ .

**375.** Suppose then that on the quadric  $\Omega$  in [5] we have a curve  $C$  whose tangents also lie on  $\Omega$ ;  $C$  being rational, of the sixth order and without cusps. No plane of [5] can be met by more than six tangents of  $C^*$ , so that there are three possibilities:

(a) Every  $\omega$ -plane of  $\Omega$  meets six tangents of  $C$ , while every  $\rho$ -plane of  $\Omega$  meets four tangents of  $C$ .

(b) Every plane of  $\Omega$  meets five tangents of  $C$ .

(c) Every  $\omega$ -plane of  $\Omega$  meets four tangents of  $C$ , while every  $\rho$ -plane of  $\Omega$  meets six tangents of  $C$ .

In any case there are four osculating planes of  $C$  lying on  $\Omega$ †.

If we project  $C$  from one of its osculating planes on to some other plane we obtain a rational plane cubic; this will have a double point, and, for certain particular osculating planes of  $C$ , it may have a cusp. But no osculating plane of  $C$  can be met by more than one tangent of  $C$ , other than the one which it contains. From this we can conclude that in (a) the four osculating planes of  $C$  which lie on  $\Omega$  are all  $\rho$ -planes, and that in (c) they are all  $\omega$ -planes.

If a rational sextic curve lies on  $\Omega$  there are eight of its trisecant planes lying on  $\Omega$ ‡; thus, besides the four osculating planes of  $C$  which lie on  $\Omega$  there must be four other trisecant planes of  $C$  also lying on  $\Omega$ .

**376.** When  $C$  is in class (a) the cuspidal edge of the developable is a rational quartic  $\Gamma$  with four stationary osculating planes; these are represented on  $\Omega$  by the four  $\rho$ -planes which osculate  $C$ . The four other planes of  $\Omega$  which are trisecant to  $C$  will all be  $\omega$ -planes; each of these in fact contains a tangent of  $C$  and meets  $C$  in a further point: they represent the four points in which  $\Gamma$  is met by its own tangents.

The chords of  $C$  form a locus  $M_3^{10}$  of three dimensions and the tenth order; there being ten chords of  $C$  meeting an arbitrary plane of [5]. The chords of  $C$  which lie on  $\Omega$  form a ruled surface which is the intersection of  $\Omega$  and  $M_3^{10}$ ; this consists in fact of two surfaces, that formed by the tangents of  $C$  and another ruled surface  $R_2^{10}$  of order ten meeting every  $\omega$ -plane of  $\Omega$  in four points and every  $\rho$ -plane of  $\Omega$  in six points.

The tangent prime of  $\Omega$  at any point of  $C$  meets  $C$  four times|| at this

\* For we cannot have a rational plane sextic with more than six cusps. See the reference to Veronese in § 347. † § 350. ‡ § 35. || § 352.

point, and in two other points; the chords to these other points are generators of  $R_2^{10}$ . Thus  $C$  is a double curve on  $R_2^{10}$ . The generators of  $R_2^{10}$  set up a symmetrical (2, 2) correspondence on  $C$ ; there are four of them which touch  $C$ , and these are precisely the tangents of  $C$  at the points where its osculating planes lie on  $\Omega$ ; the tangent prime of  $C$  at one of these points only meets  $C$  in one other point\*.

The pairs of intersections of  $C$  with those of its chords which generate  $R_2^{10}$  represent pairs of intersecting tangents of  $\Gamma$ ; the locus of these points of intersection is a nodal curve on the developable; its points are represented on  $\Omega$  by the  $\varpi$ -planes through the generators of  $R_2^{10}$  and it is thus of the sixth order. It has four cusps; these are represented on  $\Omega$  by those four  $\varpi$ -planes which contain tangents of  $C$  and meet  $C$  again. The four cusps of the nodal curve are at the four points where  $\Gamma$  is met by its own tangents. Also the nodal curve meets  $\Gamma$  in the four points where its osculating plane is stationary; these four points are represented on  $\Omega$  by the  $\varpi$ -planes through the tangents of  $C$  at the four points where its osculating plane lies on  $\Omega$ .

The  $\rho$ -planes through the generators of  $R_2^{10}$  represent planes of pairs of intersecting tangents of  $\Gamma$ ; these envelop a nodal developable of the fourth class to which belong the four stationary osculating planes of  $\Gamma$ .

When  $C$  is in class (c) we have similarly a developable whose cuspidal edge  $\Gamma$  is a sextic curve with four cusps; the cusps are represented on  $\Omega$  by the four  $\varpi$ -planes which osculate  $C$ . The nodal curve is of the fourth order passing through the cusps of  $\Gamma$ .

**377.** When  $C$  is in class (b) the four planes of  $\Omega$  which osculate  $C$  belong two to each system. The developable is thus formed by the tangents of a rational quintic curve  $\Gamma$  which has two cusps and two stationary osculating planes.

There are four other planes trisecant to  $C$  which lie on  $\Omega$ ; these will also be two of each system, and each of them contains a tangent of  $C$  and meets  $C$  in another point. The two  $\varpi$ -planes represent points in which  $\Gamma$  is met by its own tangents, while the two  $\rho$ -planes represent osculating planes of  $\Gamma$  which contain other tangents.

The chords of  $C$  which lie on  $\Omega$  again form a ruled surface  $R_2^{10}$  on which  $C$  is a double curve, but now there are five generators of the surface meeting every plane of  $\Omega$ . The tangents of  $C$ , at those four points at which the osculating planes lie on  $\Omega$ , are generators of  $R_2^{10}$ , as also are those chords of  $C$  which lie in the four other trisecant planes.

The  $\varpi$ -planes through the generators of  $R_2^{10}$  represent the points of a nodal curve; this curve is of the fifth order and passes through the cusps of  $\Gamma$ , and it has cusps at the two points where  $\Gamma$  is met by its own tangents.

\* § 352,

The  $\rho$ -planes through the generators of  $R_2^{10}$  represent planes each of which contains two intersecting tangents of  $\Gamma$ ; these envelop a nodal developable of the fifth class. The stationary osculating planes of  $\Gamma$  belong to this developable, and the two planes which osculate  $\Gamma$ , and also touch it, are stationary planes of the developable.

The tangents of  $\Gamma$  here form a developable which we did not obtain by the methods of projection and section in [4].

**378.** We now suppose that we have a rational sextic curve  $C$  in [5] with one cusp—the projection of a rational normal sextic from a point on one of its tangents. The tangents of  $C$  form a surface of order nine and any quadric containing  $C$  contains six of its tangents\*. Any quadric through  $C$  which contains seven of its tangents must contain them all, and as there are  $\infty^8$  quadrics containing  $C$  there will be  $\infty^1$  quadrics containing  $C$  and all its tangents.

Four conditions are necessary for a curve in [5] to have a cusp; if, for example, the curve is regarded as the projection of a curve in [6] without singularities the point of projection must lie on the two-dimensional locus formed by the tangents. There will thus be  $\infty^{34}$  rational sextic curves in [5] which have cusps. There are  $\infty^{20}$  quadrics in [5];  $\infty^8$  of these contain a given rational sextic with a cusp. Thus on any given quadric  $\Omega$  in [5] there are  $\infty^{22}$  rational sextic curves which have cusps and  $\infty^{15}$  of these curves are such that all their tangents lie on  $\Omega$ . The cuspidal edge  $\Gamma$  of the developable represented by  $C$  has an inflection corresponding to the cusp of  $C$ ; in the next paragraphs we shall see that  $\Gamma$  is either a rational quartic with an inflection or a rational quintic with two cusps and an inflection. In either case the aggregate of curves  $\Gamma$  is  $\infty^{15}$ ; two conditions are necessary for  $\Gamma$  to have a cusp, one condition for it to have an inflection.

**379.** Suppose then that we have on the quadric  $\Omega$  in [5] a rational sextic curve  $C$  with a cusp  $K$ . The tangents of  $C$  form a surface of order nine, and as no plane can be met by more than five tangents of  $C$ † there are two possibilities.

(a) Every  $\omega$ -plane of  $\Omega$  is met by five tangents of  $C$  and every  $\rho$ -plane by four tangents of  $C$ .

(b) Every  $\omega$ -plane of  $\Omega$  is met by four tangents of  $C$  and every  $\rho$ -plane by five tangents of  $C$ .

In either case there are two osculating planes of  $C$  lying on  $\Omega$ ‡.

If  $C$  is projected from one of its osculating planes (other than that at the cusp) on to another plane we obtain a plane cubic with a cusp. Hence no osculating plane of  $C$  can be met by any tangent of  $C$  other than the one which it contains. This is sufficient to shew that the two osculating planes of  $C$  which lie on  $\Omega$  are both  $\rho$ -planes in (a) and both  $\omega$ -planes in (b).

\* § 349.

† Just as in § 375; we cannot have a rational plane sextic with more than six cusps.

‡ § 350.

**380.** Suppose, to fix ideas, that  $C$  is of the type (a). The chords of  $C$  form a locus  $M_3^9$  of three dimensions and of order nine; there are nine chords of  $C$  meeting an arbitrary plane of [5]. The osculating plane of  $C$  at  $K$  belongs to  $M_3^9$ ; for if we regard  $C$  as the projection of a rational normal sextic from a point on one of its tangents the osculating solid of  $C$  at the point of contact of this tangent touches the  $M_3^{10}$  formed by the chords all along this tangent.

The points of  $C$  represent the generators of a developable in [3] whose cuspidal edge  $\Gamma$  is a rational quartic with an inflection.  $\Gamma$  has two stationary osculating planes; these are represented on  $\Omega$  by the two  $\rho$ -planes which osculate  $C$ .

The intersection of  $\Omega$  and  $M_3^9$  consists of the surface formed by the tangents of  $C$  and of a ruled surface  $R_2^9$  consisting of chords of  $C$  which lie on  $\Omega$ . This latter surface meets every  $\omega$ -plane of  $\Omega$  in four points and every  $\rho$ -plane of  $\Omega$  in five points. It has two generators passing through  $K$ ; these are the cuspidal tangent of  $C$  and the other line in which the osculating plane of  $C$  at  $K$  meets  $\Omega$ . The generators of  $R_2^9$  set up a symmetrical (2, 2) correspondence on  $C$ ; the four coincidences include  $K$  counted twice, so that we have two other tangents of  $C$  which are generators of  $R_2^9$ , these being the two tangents which touch  $C$  at the points where its osculating planes lie on  $\Omega$ .

The  $\omega$ -planes through the generators of  $R_2^9$  represent the points of the nodal curve of the developable in [3]. This curve is then of the fifth order and meets  $\Gamma$  in the two points at which its osculating plane is stationary, and also in its inflection.

There are two  $\omega$ -planes of  $\Omega$  which pass through tangents of  $C$  and meet  $C$  again; these represent the two points in which  $\Gamma$  is met by its own tangents. These points are cusps on the nodal curve.

The  $\rho$ -planes through the generators of  $R_2^9$  represent planes of [3] which envelop a nodal developable. This developable is of the fourth class and the two stationary osculating planes of  $\Gamma$  belong to it, as also does the osculating plane of  $\Gamma$  at its inflection.

**381.** When  $C$  is of the type (b) we have a surface dual to the one just discussed. The cuspidal edge  $\Gamma$  is now of the fifth order having two cusps and an inflection; the two cusps are represented on  $\Omega$  by the  $\omega$ -planes which osculate  $C$ , the inflection by the  $\omega$ -plane through the cuspidal tangent of  $C$ .

The nodal curve is of the fourth order passing through the cusps and the inflection of  $\Gamma$ . The nodal developable is of the fifth class; the osculating planes of  $\Gamma$  at its cusps and inflection belong to this developable. There are two planes of  $\Gamma$  which both osculate and touch it; these are stationary planes of the nodal developable.

**382.** Suppose now that we have in [4] a rational sextic with two cusps—the projection of a rational normal sextic from a line meeting two of its tangents. There are  $\infty^3$  quadric threefolds containing the curve, and each of these contains four of its tangents. There will not, in general, be a quadric containing all the tangents of the curve.

The rational sextics in [4] are  $\infty^{31}$  in aggregate\*, and as three conditions are necessary for a curve in [4] to have a cusp the rational sextics  $C$  which have two cusps are  $\infty^{25}$  in aggregate. The aggregate of quadrics in [4] is  $\infty^{14}$ , while there are  $\infty^3$  quadrics through each curve  $C$ ; hence each quadric contains  $\infty^{14}$  curves  $C$ . If a quadric should contain five tangents of  $C$  as well as  $C$  itself it necessarily contains all the tangents; thus on any given quadric there will be  $\infty^9$  curves  $C$  all of whose tangents also lie on the quadric.

Thus in [4] we have  $\infty^{23}$  curves  $C$  such that there are quadrics which contain all their tangents; hence a general rational sextic in [4] with two cusps must be specialised twice in order that a quadric should contain all its tangents.

**383.** Suppose now that we have a quadric in [4] and a rational sextic curve  $C$  with two cusps  $K_1, K_2$  whose tangents all lie on the quadric. The tangents form a ruled surface of order eight; since no line on the quadric can meet more than four tangents of  $C$  every line on the quadric must meet precisely four tangents of  $C$ .

Now regard the quadric as a prime section of  $\Omega$ . Then the points of  $C$  represent the generators of a developable surface in [3]; the edge of regression  $\Gamma$  being of the fourth order and having two inflections. The tangents of  $\Gamma$  belong to a linear complex.

The chords of  $C$  form a locus  $M_3^8$ , there being eight of them meeting an arbitrary line of [4]. The osculating planes of  $C$  at  $K_1$  and  $K_2$  belong to this locus. The quadric meets this locus in the surface formed by the tangents of  $C$  and in a ruled surface  $R_2^8$  also of order eight; this surface also meets every line on  $\Omega$  in four points and there are two of its generators passing through any point of  $C$ . The cuspidal tangents belong also to this surface; there is one other generator of  $R_2^8$  through each cusp, the other line in which the osculating plane there meets the quadric.

The  $\omega$ -planes through the generators of  $R_2^8$  represent points of the nodal curve of the developable; this passes through the inflections of  $\Gamma$  and is also of the fourth order. We also have a nodal developable of the fourth class.

**384.** One condition is necessary for a curve in [3] to have an inflection; if the curve is regarded as the projection of a curve in [4] the point of projection is constrained to lie on the three-dimensional locus formed by

\* § 354.

the osculating planes. Thus there are in [3]  $\infty^{16-2} = \infty^{14}$  curves of the fourth order with two inflections.

On the other hand, given a quadric  $\Omega$  in [5] it has  $\infty^5$  prime sections, and on each of these there lie  $\infty^9$  rational sextic curves  $C$  which have two cusps, and are such that all their tangents lie on the quadric. Hence we have  $\infty^{14}$  curves  $C$  on  $\Omega$  in agreement with the former result.

*The classification of the developable surfaces*

385. We now give a table of the different developables of the sixth order in [3]. In [4] there is only one developable of the sixth order—that formed by the tangents of a normal rational quartic curve. We have obtained ten different types of surfaces in [3]; in the first column of the table we give the double curve, in the second the bitangent developable and in the third a description of the cuspidal edge.  $C_n$  denotes a curve of order  $n$ ,  $E_n$  a developable of class  $n$  and  $T$  or  $T'$  a stationary generator.  $D_n$  is also used for a curve,  $F_n$  for a developable when necessary. All curves and developables which occur are rational.

When a bar is placed over a part of the double curve it means that the points in which a plane meets this curve are cusps and not ordinary double points on the plane section. Similarly, when a bar is placed over a part of the bitangent developable it means that the planes of this part which pass through a point are stationary planes and not ordinary double planes of the enveloping cone.

There are no developables of the sixth order in spaces of higher dimension than 4.

	Double curve	Bitangent developable	Cuspidal edge
1	$\bar{C}_4 + C_6$	$\bar{E}_6 + E_4$	Rational quartic
2	$\bar{C}_5 + C_5$	$\bar{E}_5 + E_5$	Rational quintic with two cusps
3	$\bar{C}_6 + C_4$	$\bar{E}_4 + E_6$	Rational sextic with four cusps
4	$\bar{C}_4 + 3C_2$	$\bar{E}_6 + E_4$	Rational quartic whose four stationary osculating planes have their points of contact coplanar
5	$\bar{C}_6 + C_4$	$E_4 + 3E_2$	Rational sextic with four cusps, the osculating planes at the cusps being concurrent
6	$\bar{C}_4 + C_3 + D_3$	$\bar{E}_6 + E_2 + F_2$	Rational quartic with a double point
7	$\bar{C}_6 + C_2 + D_2$	$\bar{E}_4 + E_2 + F_2$	Rational sextic with four cusps and a doubly osculating plane
8	$\bar{C}_4 + \bar{T} + C_6$	$\bar{E}_5 + \bar{T} + E_4$	Rational quartic with an inflection
9	$\bar{C}_6 + \bar{T} + C_4$	$\bar{E}_4 + \bar{T} + E_6$	Rational quintic with two cusps and an inflection
10	$\bar{C}_4 + \bar{T} + \bar{T}' + C_4$	$\bar{E}_4 + \bar{T} + \bar{T}' + E_4$	Rational quartic with two inflections

386. We can give another table shewing the singularities of the cuspidal edges of these developables. We shall denote the order of a curve by  $n$ , the rank by  $r$ , the class by  $n'$ , the number of cusps by  $\kappa$ , the number of inflections by  $i$  and the number of stationary osculating planes by  $\kappa'$ . Also let  $h$  denote the number of apparent double points and  $\delta$  the number of actual double points;  $\delta'$  the number of doubly osculating planes and  $h'$  the number of lines in any given plane which are intersections of two osculating planes of the curve. Also let  $b$  be the number of planes through any point which are bitangent to the curve and  $b'$  the order of the nodal curve of the developable.

Then we can give the following table; we omit the developables (4) and (5) because the table would not distinguish them from (1) and (2).

	$n$	$r$	$n'$	$\kappa$	$i$	$\kappa'$	$b$	$b'$	$h$	$\delta$	$h'$	$\delta'$
1	4	6	6	0	0	4	4	6	3	0	6	0
2	5	6	5	2	0	2	5	5	4	0	4	0
3	6	6	4	4	0	0	6	4	6	0	3	0
6	4	6	6	0	0	4	4	6	2	1	6	0
7	6	6	4	4	0	0	6	4	6	0	2	1
8	4	6	5	0	1	2	4	5	3	0	4	0
9	5	6	4	2	1	0	5	4	4	0	3	0
10	4	6	4	0	2	0	4	4	3	0	3	0

## CHAPTER VI

### SEXTIC RULED SURFACES (CONTINUED)

#### *Further types of rational sextic ruled surfaces*

**387.** The developable surface of the fourth order has for its cuspidal edge a twisted cubic; it has no nodal curve. It is a particular case of the general rational quartic ruled surface which has a twisted cubic for its double curve.

The developable surface of the fifth order has for its cuspidal edge a rational quartic with a cusp; its nodal curve is a conic passing through the cusp of the quartic and meeting it in two other points. It is a particular case of the rational quintic ruled surface whose double curve breaks up into a rational quartic and a conic; the quartic has a double point through which the conic passes and the two curves have two other intersections (cf. § 92 and § 130).

When we turn to the developable surfaces of the sixth order we at once notice the fact that, except for the two developables that we have numbered (6) and (7), we have not obtained any rational sextic ruled surfaces of which they are special cases. The developable (6) has two nodal plane cubics and, for its cuspidal edge, a rational quartic with a double point; it is a particular case of the rational sextic ruled surface whose double curve consists of two twisted cubics and a rational quartic (cf. § 180). The developable (7) has two nodal conics and, for its cuspidal edge, a rational sextic with four cusps; it is a particular case of the rational sextic ruled surface whose double curve consists of two conics and a rational sextic with four double points (cf. § 179).

This leads us to suspect that there still remain types of rational sextic ruled surfaces other than the sixty-seven types we have already obtained. For example: the existence of the first developable, whose cuspidal edge is a rational quartic and nodal curve a rational sextic, leads us to expect a type of rational sextic ruled surface whose double curve consists of a rational quartic and a rational sextic, and we have not yet obtained such a surface. Since the generators of the developable are tangents of the cuspidal edge and chords of the double curve we expect the generators of the ruled surface to meet each part of the double curve in two points.

This last remark gives the clue to the discovery of these further types of surfaces. For if we consider the double curve of any type of rational sextic ruled surface that we have obtained, and omit any double generators that may occur, the remainder of the double curve consists of a number of



components of which all, with at most one exception, are directrices, meeting each generator in one point. The solid  $S$  from which we project the normal surface meets an infinity of chords of one or more directrix curves of the normal surface.

**388.** Let us then take a curve meeting each generator of the normal surface  $F$  in two points, the surface having  $\infty^1$  directrix cubic curves. We take first an elliptic curve  $C_8^1$  of order eight; such a curve is given by a quadric primal containing four generators of  $F$ . The chords joining pairs of points of a rational involution on  $C_8^1$  form a rational sextic ruled surface with, in general,  $\infty^1$  directrix cubic curves each meeting  $C_8^1$  twice; but there are 16 involutions for which the sextic ruled surface has a directrix conic\*, this conic not meeting  $C_8^1$ . Let us choose  $S$  to contain the plane of such a conic  $\vartheta_2$ ; it meets the locus  $M_5^6$  formed by the chords of  $F$  in  $\vartheta_2$  and a rational quartic  $\vartheta_4^0$  having four intersections with  $\vartheta_2$ . The chords of  $F$  which meet  $\vartheta_4^0$  include four tangents and meet  $F$  in the points of an elliptic curve  $C_{12}^1$ ;  $C_{12}^1$  meets each generator twice and has four double points. The two chords of  $F$  which pass through any one of these double points and meet  $\vartheta_4^0$  meet  $F$  again in intersections of  $C_{12}^1$  and  $C_8^1$ ; this accounts for eight of the sixteen intersections of the two curves. The remaining eight lie in pairs on the chords of  $F$  which pass through the four intersections of  $\vartheta_2$  and  $\vartheta_4^0$ .

Hence, projecting from  $S$ , we obtain a surface  $f$  in [3] whose double curve is  $C_4^0 + C_6^0$ .  $C_6^0$  has four double points through all of which  $C_4^0$  passes, and the two curves have four other intersections.

If we project  $F$  from the plane of  $\vartheta_2$  on to a [4] we obtain a rational sextic ruled surface with a rational normal quartic double curve. The generators of the surface are chords of the curve, through each point of the curve there pass two of them. The surface is the intersection of the cubic primal formed by the chords of the curve with a quadric containing the curve.

**389.** The involution on  $C_8^1$  sets up a symmetrical (2, 2) correspondence between the generators of  $F$ . There are four generators  $g$  of  $F$  for which the two corresponding generators coincide; these are the four generators  $g$  which touch  $C_8^1$ . There are four generators of  $f$  touching  $C_4^0$  and four touching  $C_6^0$ .

The (2, 2) correspondence between the generators of  $F$  gives a (2, 2) correspondence on any directrix cubic of  $F$ ; the chords of the cubic which join pairs of corresponding points form a rational quartic ruled surface. Hence the planes which are determined by those pairs of generators of  $f$  which meet in the points of  $C_4^0$  are the projections, from  $S$ , of the solids of an  $M_4^8$ . Each solid of  $M_4^8$  is determined by two corresponding lines in a (1, 1) correspondence, without united elements, between the generators of two rational quartic ruled surfaces. The projection of  $M_4^8$  from the solid

\* Segre, *Math. Ann.* 27 (1886), 296.

$S$ , which contains a conic  $\vartheta_2$  meeting each solid of  $M_4^8$ , is a rational developable  $E_6^0$  of the sixth class. This is part of the bitangent developable  $E_6^0 + E_4^0$  of  $f$ .

The solids which contain the directrix cubics of  $F$  form a locus  $M_4^4$ ; there are four of them which meet  $S$ . The [6] containing  $S$  and any one of these four solids meets the space on to which we are projecting in a tri-tangent plane of  $f$ . It meets  $S$  in a directrix cubic  $\Delta$  and in three generators; it contains eight points of  $C_8^1$ , two on each generator and two on  $\Delta$ . Now the pairs of points in which  $C_8^1$  is met by the directrix cubics of  $F$  are the pairs of a  $g_2^1$ , and two  $g_2^1$ 's on an elliptic curve cannot have a common pair of points. Hence the chord of  $C_8^1$  which joins its intersections with  $\Delta$  does not meet  $\vartheta_2$ . The eight points of  $C_8^1$  which lie in the [6] are joined in pairs by four lines which meet  $\vartheta_2$ ; the [6] meets the space on to which we are projecting in a plane which is a plane of  $E_4^0$  and a double plane of  $E_6^0$ .

**390.** We now choose the solid  $S$  to contain  $\vartheta_2$  and also to meet the solid containing a pair  $g, g'$  of generators of  $F$  in a line  $l$ . It can be shewn that  $l$  must meet the plane of  $\vartheta_2$  in a point  $O$  of  $\vartheta_2$  itself, and that the chord of  $C_8^1$  through  $O$  meets  $F$  on the generators  $g$  and  $g'$ .  $S$  meets  $M_5^6$  in  $\vartheta_2$ ,  $l$  and a twisted cubic  $\vartheta_3^0$  which meets  $l$  once and  $\vartheta_2$  three times.

The chords of  $F$  which meet  $S$  meet  $F$  in  $C_8^1, g, g'$  and an elliptic curve  $C_{10}^1$ ; there are four tangents of  $C_{10}^1$  meeting  $\vartheta_3^0$ , and it meets each generator of  $F$  in two points.  $C_{10}^1$  has two double points.

Let  $OAA'$  be the chord of  $F$  through  $O$ , meeting  $g$  in  $A$  and  $g'$  in  $A'$ .  $C_8^2$  passes through  $A$  and  $A'$ ; let it meet  $g$  and  $g'$  again in  $B$  and  $B'$  respectively. We have chords  $BC, B'C'$  of  $C_8^1$  meeting  $\vartheta_2$  and chords  $BD', B'D$  meeting  $l$ ;  $D'$  is on  $g'$  and  $D$  on  $g$ . The lines  $CD'$  and  $C'D$  are chords of  $C_{10}^1$  and meet  $\vartheta_3^0$ ; the other intersections of  $C_{10}^1$  with  $g$  and  $g'$  lie on the chord of  $F$  which passes through the intersection of  $l$  and  $\vartheta_3^0$ . There are three pairs of intersections of  $C_8^1$  and  $C_{10}^1$  collinear with the three intersections of  $\vartheta_2$  and  $\vartheta_3^0$ ; there are four other intersections of  $C_8^1$  and  $C_{10}^1$ , these are associated in pairs with the two double points of  $C_{10}^1$ .

The projected surface  $f$  has a double curve  $C_4^0 + G + C_5^0$ .  $G$  is a trisecant of  $C_4^0, C_5^0$  passing through two of their intersections.  $C_5^0$  has two double points;  $C_4^0$  passes through both these double points, as well as meeting  $C_5^0$  in three further points which do not lie on  $G$ .

The locus  $M_4^8$  now meets  $S$  in the conic  $\vartheta_2$  and the line  $l$ ; the planes of the pairs of generators of  $f$  which intersect in the points of  $C_4^0$  form a developable  $E_5^0$  of the fifth class. The bitangent developable of  $f$  is  $E_5^0 + G + E_4^0$ . The solid  $S$  meets four solids which contain directrix cubics of  $F$ ; two of these four meet  $S$  in points on  $l$ . Thus  $f$  has two tri-tangent planes which do not pass through  $G$ ; these are planes of  $E_4^0$  and double planes of  $E_5^0$ .

391. We can further choose  $S$  to contain  $\mathfrak{D}_2$ , a line  $l$  lying in the solid containing a pair of generators  $g, g'$  of  $F$  and a line  $m$  lying in the solid containing a pair of generators  $h, h'$  of  $F$ .  $l$  and  $m$  meet  $\mathfrak{D}_2$ .  $S$  meets  $M_5^6$  in  $\mathfrak{D}_2, l, m$  and a conic  $\phi_2$ ;  $\phi_2$  meets  $l$  and  $m$  and meets  $\mathfrak{D}_2$  twice.

The chords of  $F$  which meet  $S$  meet  $F$  in  $C_8^1, g, g', h, h'$  and an elliptic curve  $D_8^1$ ;  $D_8^1$  meets each generator of  $F$  twice and has four of its tangents meeting  $\phi_2$ .  $C_8^1$  and  $D_8^1$  have eight intersections.

The projected surface  $f$  has two double generators, its double curve being  $C_4^0 + G + H + D_4^0$ .  $C_4^0$  and  $D_4^0$  have two intersections on  $G$ , two on  $H$ , and two others.  $G$  and  $H$  are trisecants both of  $C_4^0$  and  $D_4^0$ . The bitangent developable is  $E_4^0 + G + H + F_4^0$ .

392. A curve of order ten which meets each generator of  $F$  twice is elliptic provided that it has two double points. Let us then consider such a curve  $C_{10}^1$ ; it can be given by a quadric containing two generators of  $F$  and touching  $F$  twice.

The joins of the pairs of points of a rational involution on  $C_{10}^1$  form an octavic ruled surface; in general this surface has  $\infty^1$  directrix quartic curves, but there are 25 involutions for which the resulting surface has a directrix cubic. The curve  $C_{10}^1$  is the projection of a normal curve in [9] from a line meeting two of its chords. An involution on an elliptic curve is determined when *one* of its pairs of points has been assigned; the normal curve is projected from a line which meets two chords joining pairs of points of the *same* involution. This involution gives an octavic ruled surface which is projected into the sextic surface  $F$ .

We choose the solid  $S$ , from which  $F$  is to be projected, to contain the directrix cubic  $\mathfrak{D}_3^0$  of the ruled surface determined by one of the 25 special involutions.  $S$  meets  $M_5^6$  in  $\mathfrak{D}_3^0$  and a second twisted cubic  $\phi_3^0$ , the two curves having four intersections. There are four tangents of  $C_{10}^1$  meeting  $\mathfrak{D}_3^0$ ; the chords of  $F$  which meet  $S$  meet  $F$  in  $C_{10}^1$  and a second elliptic curve  $D_{10}^1$ . There are four tangents of  $D_{10}^1$  meeting  $\phi_3^0$ ; it meets each generator of  $F$  in two points and has two double points.  $C_{10}^1$  and  $D_{10}^1$  have sixteen intersections; eight of these are collinear in pairs with the four intersections of  $\mathfrak{D}_3^0$  and  $\phi_3^0$ , the remaining eight fall into four pairs associated with the two double points of  $C_{10}^1$  and the two double points of  $D_{10}^1$ .

The double curve of the projected surface  $f$  is  $C_5^0 + D_5^0$ . Each of  $C_5^0$  and  $D_5^0$  has two double points through which the other passes, and the two curves have four further intersections.

The bitangent developable is  $E_5^0 + F_5^0$ . Either part of this developable is the projection of a locus  $M_4^8$  from a solid  $S$  which contains a cubic curve meeting all the solids of  $M_4^8$ .

We can also obtain surfaces in [3], with double curves and bitangent developables as above, by projection from the normal surface with a directrix conic. A surface whose double curve is  $C_5^0 + D_5^0$  can be obtained

by projection from the normal surface with a directrix line; we have a bitangent developable  $10R$  and a double curve  $C_5^0 + D_5^0$ .

**393.** The sextic ruled surface whose double curve is  $C_4^0 + C_6^0$  is generated by a symmetrical (2, 2) correspondence\* on a rational quartic  $C_4^0$ . If we specialise this correspondence so that the two points corresponding to some point  $P$  of the curve are the points  $Q$  and  $R$  where the curve is met by its trisecant through  $P$ , the surface has a double generator and its double curve is  $C_4^0 + G + C_6^0$ ;  $C_6^0$  passes through  $Q$  and  $R$ . We can specialise the correspondence further and obtain two double generators. We cannot specialise the correspondence still further in this way† without it degenerating into the  $g_3^1$  cut out by the trisecants of the curve; the sextic ruled surface would then degenerate into a quadric counted three times.

If we have a rational quintic  $C_5^0$  in [3], with two double points, and set up on the curve a symmetrical (2, 2) correspondence in which the two points on the different branches at either double point of  $C_5^0$  correspond to one another, the chords joining pairs of corresponding points generate a sextic ruled surface whose double curve is  $C_5^0 + D_5^0$ . This is the projection of the ruled surface of order eight, generated by a symmetrical (2, 2) correspondence on a rational normal quintic curve in [5], from a line meeting two of its generators.

**394.** We now give the representation of these ruled surfaces as curves on  $\Omega$ . Through each point of a rational sextic curve  $C$  on  $\Omega$  there pass four of its chords which lie on  $\Omega$ ; the chords of  $C$  which lie on  $\Omega$  set up a symmetrical (4, 4) correspondence on  $C$ . It is characteristic of the types of ruled surfaces that we have just obtained that this correspondence degenerates into the sum of two symmetrical (2, 2) correspondences.

Suppose that  $C$  has no double points. The joins of corresponding points in a symmetrical (2, 2) correspondence on  $C$  form a ruled surface of order ten; the ruled surface  $R_2^{20}$  formed by the chords of  $C$  which lie on  $\Omega$  (cf. § 173) here breaks up into two ruled surfaces  $R_2^{10}$  and  $S_2^{10}$ , both of order ten and both rational.

Suppose that  $PQR$  is a trisecant plane of  $C$  which lies on  $\Omega$ . Then the chords  $QR$ ,  $RP$ ,  $PQ$  of  $C$  cannot all belong to the same ruled surface, for then the symmetrical (2, 2) correspondence would be an involution of sets of three points‡, and we should have  $\infty^1$  trisecant planes of  $C$  lying on  $\Omega$ . Hence two of the three chords must belong to one of the two ruled surfaces, the other chord belonging to the other ruled surface. The double curve of the ruled surface in [3] breaks up into two parts; one of these parts has its points in (1, 1) correspondence with the generators of  $R_2^{10}$ ,

\* See footnote to § 19.

† The symmetrical (2, 2) correspondence and the  $g_3^1$  given by the trisecants have four common pairs of points and no more. See § 15.

‡ Cf. Baker. *Principles of Geometry*, 2 (Cambridge 1922), 135-6.

the other has its points in (1, 1) correspondence with the generators of  $S_2^{10}$ . At each of the four triple points of the sextic ruled surface there is one part of the double curve which has a double point. Similarly each tri-tangent plane of the surface is a double plane of one of the two parts of the bitangent developable.

The surfaces  $R_2^{10}$  and  $S_2^{10}$  have four common generators; the two parts of the double curve have four intersections other than at triple points of the ruled surface while the two parts of the bitangent developable have in common four planes which are not tri-tangent planes of the ruled surface.

Neither of the ruled surfaces  $R_2^{10}$  and  $S_2^{10}$  can meet a plane of  $\Omega$  in more than six points. Suppose, first, that  $R_2^{10}$  meets every  $\omega$ -plane of  $\Omega$  in four points and every  $\rho$ -plane of  $\Omega$  in six points; then, since each plane of  $\Omega$  is met by ten chords of  $C$ ,  $S_2^{10}$  meets every  $\omega$ -plane of  $\Omega$  in six points and every  $\rho$ -plane of  $\Omega$  in four points. The surface in [3] has a double curve  $C_4^0 + C_6^0$  and a bitangent developable  $E_6^0 + E_4^0$ . The four  $\omega$ -planes of  $\Omega$  which are trisecant to  $C$  contain each two generators of  $S_2^{10}$  and one generator of  $R_2^{10}$ ;  $C_6^0$  has four double points through each of which  $C_4^0$  passes. Also  $E_6^0$  has four double planes, these being also planes of  $E_4^0$ .

It may also happen that both  $R_2^{10}$  and  $S_2^{10}$  meet every plane of  $\Omega$  in five points. Then the surface in [3] has a double curve  $C_5^0 + D_5^0$  and a bitangent developable  $E_5^0 + F_5^0$ .

**395.** Suppose now that  $C$  has a double point  $P$ ; and that the points of  $C$  on the two different branches at  $P$  are corresponding points in both the symmetrical (2, 2) correspondences. These correspondences give rise to two rational ruled surfaces  $R_2^9$  and  $S_2^9$ , both of the ninth order. Each of these surfaces must meet the planes of one system each in four points and the planes of the opposite system each in five points; each plane of  $\Omega$  is met by four generators of one surface and five of the other. The corresponding ruled surface in [3] has a double curve  $C_5^0 + G + C_4^0$  and a bitangent developable  $E_4^0 + G + E_5^0$ .

The plane of the two tangents of  $C$  at  $P$  meets  $\Omega$  in two lines which are common generators of  $R_2^9$  and  $S_2^9$ ; the two surfaces have two other common generators and each surface has a third generator passing through  $P$ . There are four trisecant planes of  $C$  lying on  $\Omega$ , two being  $\omega$ -planes and two  $\rho$ -planes. Thus  $C_5^0$  has two double points through each of which  $C_4^0$  passes; the curves have two intersections on  $G$  in addition to two further intersections, and  $G$  is a trisecant of both curves. There are similar statements for the bitangent developable.

We can suppose also that  $C$  lies in [4] and has two double points  $P$  and  $Q$ ; and that the points of  $C$  on the two different branches at either double point are corresponding points in both symmetrical (2, 2) correspondences. These correspondences give rise to two rational ruled surfaces

$R_2^8$  and  $S_2^8$ , both of the eighth order. Each of these surfaces meets every plane of  $\Omega$  in four points on the line in which it meets the [4] containing  $C$ , and the corresponding ruled surface in [3] has a double curve  $C_4^0 + G + H + D_4^0$  and a bitangent developable  $E_4^0 + G + H + F_4^0$ .

There are two common generators of  $R_2^8$  and  $S_2^8$  in the plane containing the tangents of  $C$  at  $P$ , and there is a third generator of each surface through  $P$ ; similarly for  $Q$ . There are no other common generators of  $R_2^8$  and  $S_2^8$  other than the two at  $P$  and the two at  $Q$ , and there are no trisecants of  $C$ . Hence  $C_4^0$  and  $D_4^0$  have two intersections on  $G$  and two intersections on  $H$ ;  $G$  and  $H$  are trisecants both of  $C_4^0$  and of  $D_4^0$ .

396. There are two types of rational sextic ruled surfaces which were overlooked in the first section of Chapter IV; we will obtain them now.

Take the rational normal sextic ruled surface  $F$  in [7] which has  $\infty^1$  directrix cubic curves, and on this surface two directrix quartics  $E$  and  $E'$ . These quartic curves have two intersections, and we consider two involutions of pairs of generators of  $F$ , the generators through the two intersections of  $E$  and  $E'$  forming a pair of both involutions. The first involution determines a cubic ruled surface containing  $E$  and the second determines similarly a cubic ruled surface containing  $E'$ ; these two cubic ruled surfaces have directrices  $\lambda$  and  $\lambda'$  respectively, and we choose the two involutions so that  $\lambda$  and  $\lambda'$  intersect on the common chord of  $E$  and  $E'$ . We take, further, a pair of generators  $g, g'$  which belong to the first involution and a line  $l$  meeting  $\lambda$  and lying in the solid  $gg'$ . We then project  $F$  on to a [3]  $\Sigma$  from the solid  $S$  determined by  $\lambda, \lambda'$  and  $l$ .

$S$  meets  $M_5^0$  in  $\lambda, \lambda', l$  and a twisted cubic  $\mathfrak{d}_3^0$  which meets  $l$  and  $\lambda$  and has  $\lambda'$  as a chord. The chords of  $F$  which meet  $S$  meet  $F$  in  $E, E', g, g'$  and an elliptic curve  $C_{10}^1$  of which four tangents meet  $\mathfrak{d}_3^0$ ;  $C_{10}^1$  meets each generator in two points and has two double points, and meets each of  $E$  and  $E'$  in six points.

Two of the intersections of  $C_{10}^1$  with  $E$  are on the chord of  $F$  which passes through the intersection of  $\lambda$  and  $\mathfrak{d}_3^0$ ; the remaining four are associated in pairs with the two double points of  $C_{10}^1$ . The chord of  $F$  which passes through the intersection of  $\mathfrak{d}_3^0$  and  $l$  accounts for an intersection of  $C_{10}^1$  with each of  $g$  and  $g'$ ; the remaining intersections of  $C_{10}^1$  with  $g$  and  $g'$  are associated with the intersections of  $E'$  with  $g'$  and  $g$  (respectively) and with two of the intersections of  $E'$  and  $C_{10}^1$ . The remaining intersections of  $E'$  and  $C_{10}^1$  are collinear in pairs with the two intersections of  $\lambda'$  and  $\mathfrak{d}_3^0$ .

We obtain, on projection, a surface  $f$  in  $\Sigma$  whose double curve is  $C_2 + D_2 + G + C_5^0$ .  $G$  is a chord of  $D_2$  and  $C_5^0$  passes through their intersections, meeting  $D_2$  in two other points and  $G$  in one other point.  $C_5^0$  has two double points through each of which  $C_2$  passes;  $C_2$  meets  $C_5^0$  in one other point and meets  $G$  and  $D_2$  each in one point.

The surface is generated by a (2, 2) correspondence between two conics  $C_2$  and  $D_2$  with a common point, this being a doubly united point. The correspondence is to be specialised to give the double generator; to the second intersection of  $C_2$  with the plane of  $D_2$  there corresponds a pair of points of  $D_2$  collinear with it. The planes of the pairs of generators which intersect in the points of  $C_2$  envelop a quadric cone  $E_2$ ; the planes of the pairs of generators which intersect in the points of  $D_2$  envelop a cubic developable  $E_3^0$ . There are two planes of  $E_3^0$  and one plane of  $E_2$  passing through  $G$ . The bitangent developable of the surface is  $E_2 + E_3^0 + G + E_4^0$ .

397. There is also a surface dual to the one just mentioned; the double curve of this surface will be  $C_2 + C_3^0 + G + C_4^0$ , the bitangent developable  $E_2 + F_2 + G + E_5^0$ .

To obtain this surface by projection we consider, on the normal surface  $F$  with  $\infty^1$  directrix cubic curves, a directrix quartic  $E$  and a prime section  $C_6^0$ ; these have four intersections. We then consider two involutions on  $F$ ; the first of these determines, by means of  $E$ , a cubic ruled surface with a directrix line  $\lambda$ , and the second determines, by means of  $C_6^0$ , a quintic ruled surface with a directrix conic  $\Gamma$ . We choose the involutions so that  $\lambda$  and  $\Gamma$  meet on a chord of  $F$  which joins two of the intersections of  $E$  and  $C_6^0$ . We can determine a solid  $S$  which not only contains  $\lambda$  and  $\Gamma$  but also a line  $l$ , meeting  $\lambda$ , which lies in the solid determined by a pair of generators  $g, g'$  belonging to the first involution. We project  $F$  from  $S$  on to  $\Sigma$ .

$S$  meets  $M_5^6$  in  $\lambda, l, \Gamma$  and a conic  $\Delta$  meeting  $\Gamma$  twice and  $l$  and  $\lambda$  each once. The chords of  $F$  which meet  $S$  meet  $F$  in  $E, g, g', C_6^0$  and an elliptic curve  $C_8^1$  of which four tangents meet  $\Delta$ ;  $C_8^1$  meets each generator in two points, has no double points, and meets  $E$  in four points and  $C_6^0$  in eight points.

Two of the intersections of  $C_8^1$  with  $E$  are on the chord of  $F$  which passes through the intersection of  $\lambda$  and  $\Delta$ ; the remaining two are associated with those two intersections of  $E$  and  $C_6^0$  which do not lie on the chord of  $F$  passing through the intersection of  $\lambda$  and  $\Gamma$  and with two of the intersections of  $C_8^1$  and  $C_6^0$ . There are two other intersections of  $C_8^1$  and  $C_6^0$  associated with the intersections of  $C_6^0$  with  $g$  and  $g'$  and with intersections of  $C_8^1$  with  $g'$  and  $g$  (respectively). The other intersections of  $C_8^1$  with  $g'$  and  $g$  lie on the chord of  $F$  through the intersection of  $\Delta$  and  $l$ , while the four remaining intersections of  $C_8^1$  and  $C_6^0$  are collinear in pairs with the two intersections of  $\Gamma$  and  $\Delta$ .

We obtain, on projection, a surface  $f$  in  $\Sigma$  whose double curve is  $C_2 + C_3^0 + G + C_4^0$ .  $C_2$  and  $C_3^0$  have three intersections through two of which  $C_4^0$  passes;  $C_4^0$  meets  $C_2$  in one other point and  $C_3^0$  in four other points.  $G$  joins two of these four points; it meets  $C_4^0$  again and also meets  $C_2$ .

The surface is generated by a (2, 2) correspondence between a conic  $C_2$

and a twisted cubic  $C_3^0$ ; the curves have three intersections one of which is a doubly united point, the other two being ordinary united points. The correspondence is specialised to give the double generator; there is a point of  $C_2$  to which correspond the two points in which  $C_3^0$  is met by the chord passing through the point. The planes joining the points of  $C_2$  to the pairs of points of  $C_3^0$  which correspond to them touch a quadric cone  $E_2$ , while the planes joining the points of  $C_3$  to the pairs of points of  $C_2$  which correspond to them touch a quadric cone  $F_2$ . The bitangent developable is  $E_2 + F_2 + G + E_5^0$ . There are two planes of  $F_2$  and one plane of  $E_2$  passing through  $G$ .

398. We now shew how the two surfaces which we have just obtained are represented as rational sextic curves  $C$  on the quadric  $\Omega$  in [5]. Since the surfaces are not self-dual their generators cannot belong to a linear complex; hence  $C$  actually belongs to [5] and is not contained in a space of lower dimension.  $C$  has a double point  $P$  representing the double generator  $G$ .

$C$  is the projection of a rational normal sextic curve from a point on one of its chords. The chords which join the pairs of points of an involution on a rational normal sextic curve form a rational normal quintic ruled surface with a directrix conic; we take two such involutions and project from a point on a chord belonging to one of them. We obtain a curve  $C$ , with a double point  $P$ , in [5]; we have, containing  $C$ , a quartic ruled surface and a quintic ruled surface with a common generator. Of the  $\infty^{20}$  quadrics in [5] there are  $\infty^8$  which contain  $C$ ; of these  $\infty^8$  there are  $\infty^3$  containing the quintic ruled surface,  $\infty^5$  containing the quartic ruled surface, and  $\infty^1$  containing both surfaces. We take one of these last quadrics to be  $\Omega$ .

There is, on the quintic surface, a plane cubic curve with a double point at  $P$  (cf. § 191); the plane of this curve lies on  $\Omega$ . Every plane of  $\Omega$  of the same system as this meets two generators of the quintic surface; every plane of the opposite system meets three generators of the quintic surface. The quartic ruled surface meets every plane of  $\Omega$  in two points. The chords of  $C$  which lie on  $\Omega$  form the quartic ruled surface  $R_2^4$ , the quintic ruled surface  $R_2^5$ , and a rational ruled surface  $R_2^9$  of order nine on which  $C$  is a double curve. The plane of the two tangents of  $C$  at  $P$  meets  $\Omega$  in two lines one of which is a generator of  $R_2^4$  and the other of  $R_2^9$ . There are two generators of  $R_2^5$  passing through  $P$ ; these are also generators of  $R_2^9$ ;  $R_2^5$  and  $R_2^9$  have two other common generators while  $R_2^4$  and  $R_2^9$  have one common generator.  $R_2^4$  and  $R_2^5$  have one common generator. A plane of  $\Omega$  which belongs to the same system as the plane of the cubic curve on  $R_2^5$  is met by five generators of  $R_2^9$ ; a plane of the opposite system is met by four generators of  $R_2^9$ .

Suppose that the plane cubic on  $R_2^5$  lies in a  $\rho$ -plane of  $\Omega$ . Then the



surface in [3] has a double curve  $C_2 + D_2 + G + C_5^0$  and a bitangent developable  $E_2 + E_3^0 + G + E_4^0$ . If, on the other hand, the plane cubic on  $R_2^5$  lies in a  $\omega$ -plane of  $\Omega$  the surface has a double curve  $C_2 + C_3^0 + G + C_4^0$  and a bitangent developable  $E_2 + F_2 + G + E_5^0$ .

*Sextic ruled surfaces with a triple curve*

399. One of the developables of the sixth order which we have obtained has a triple conic; in order to obtain a sextic ruled surface with a triple curve by projection we must choose, for the centre of projection, a space which is met in lines by  $\infty^1$  trisecant planes of the normal surface.

Consider a prime section  $C_6^0$  of the rational normal sextic ruled surface  $F$  in [7] which has  $\infty^1$  directrix cubics. Let us consider the planes which contain the triads of points which form sets of a  $g_3^1$  on  $C_6^0$ ; they form a three-dimensional locus whose order is the number of planes which meet an arbitrary solid,  $\Pi$  say, in the prime which contains  $C_6^0$ . Now the trisecant planes of  $C_6^0$  are the intersections of corresponding primes of four projectively related triply infinite systems\*; to obtain the planes which contain the sets of a  $g_3^1$  we consider four corresponding pencils of primes belonging to the four systems. The solid  $\Pi$  meets these pencils in four projectively related pencils of planes; and it is a well-known result that there are *four* points which are intersections of corresponding planes of the four pencils†. Hence the planes which contain the sets of a  $g_3^1$  on  $C_6^0$  form a three-dimensional quartic locus  $V_3^4$ .

This locus  $V_3^4$  is normal in [6]; it can be generated by the planes which join triads of corresponding points of two lines and a conic, projectively related to each other‡. It therefore contains the quadric surface  $Q$  determined by the projective relation between the lines; there is a generator of  $Q$  in each plane of  $V_3^4$ . We select the solid  $S$  which contains  $Q$  as the centre of projection, and project  $F$  from  $S$  on to a [3]  $\Sigma$ .

400. The curve  $C_6^0$  does not meet  $Q$ ; the chords of  $C_6^0$  which lie in the planes of  $V_3^4$  form a ruled surface of order ten which meets  $Q$  in a rational quartic curve  $\vartheta_4^0$ , the generators of  $Q$  which lie in the planes of  $V_3^4$  being the trisecants of  $\vartheta_4^0$ . Since the  $g_3^1$  on  $C_6^0$  has four double points there are four tangents of  $C_6^0$  which meet  $\vartheta_4^0$ . The solid  $S$  meets the locus  $M_5^6$  formed by the chords of  $F$  in  $\vartheta_4^0$  and a conic  $\vartheta_2, \vartheta_2$  meeting  $\vartheta_4^0$  in four

\* The bases of the four systems may be taken to be any four fixed trisecant planes of  $C_6^0$ .

† If  $l_1, l_2, l_3, l_4$  denote the axes of the four pencils, the two pencils whose axes are  $l_2$  and  $l_3$  generate, by means of the lines of intersection of corresponding planes, a regulus. This regulus contains the cubic curve generated by the pencils  $l_1, l_2, l_3$  and also that generated by the pencils  $l_2, l_3, l_4$ ; these two cubic curves have four intersections.

‡ Segre, *Atti Torino*, 21 (1885), 95.

points. The chords of  $F$  which meet  $S$  meet  $F$  in the curve  $C_8^0$  counted twice (since two of them pass through each point of  $C_8^0$ ) and an elliptic curve  $C_8^1$ , four of whose tangents meet  $\mathfrak{D}_2$ , meeting each generator twice. The eight intersections of  $C_8^1$  and  $C_8^0$  are collinear in pairs with the four intersections of  $\mathfrak{D}_2$  and  $\mathfrak{D}_4^0$ .

On projection we obtain a surface  $f$  in  $\Sigma$  whose double curve is  $3C_2 + C_4^0$ ,  $C_2$  and  $C_4^0$  having four intersections. The surface is generated by a conic  $C_2$  and a twisted cubic in (1, 3) correspondence with three united points. The planes of pairs of generators which intersect in the points of  $C_2$  envelop a developable  $E_4^0$ ; the bitangent developable of the surface is  $E_4^0 + E_6^0$ .

The surface is also generated by the chords of  $C_4^0$  which meet  $C_2$ .

There is also a dual surface whose double curve is  $C_4^0 + C_6^0$  and bitangent developable  $3E_2 + E_4^0$ .

401. We can choose the  $g_3^1$  on  $C_6^0$  so that  $S$  meets  $M_5^6$  in  $\mathfrak{D}_4^0$  and two lines  $l$  and  $m$ ; these lines are chords of  $\mathfrak{D}_4^0$  and intersect. They are axes of two directrix quartics  $E$  and  $E'$ , the common chord of  $E$  and  $E'$  passing through the intersection of  $l$  and  $m$ . On projection from  $S$  we obtain a surface  $f$  whose double curve  $3B_2 + C_2 + D_2$  consists of a triple conic and two double conics;  $C_2$  and  $D_2$  have one intersection and each of them meets  $B_2$  in two points. The surface is generated by two conics,  $B_2$  and  $C_2$ , in (2, 3) correspondence with two doubly united points. The planes of the pairs of generators which intersect in the points of  $C_2$  or  $D_2$  envelop a developable of the third class. The bitangent developable of the surface is  $E_4^0 + E_3^0 + F_3^0$ .

The surface is generated by the lines which meet the three conics.

There is also a dual surface whose double curve is  $C_4^0 + C_3^0 + D_3^0$  and bitangent developable  $3E_2 + F_2 + G_2$ .

402. The planes which contain the triads of points of a  $g_3^1$  on a rational sextic curve  $C$  in [5] form a locus  $v_3^4$  which is the projection of the normal locus  $V_3^4$  formed by the planes containing the triads of points of a  $g_3^1$  on the normal curve. We obtain different types of loci  $v_3^4$  according as to the position we select for the point of projection; the only locus  $v_3^4$  which lies on a non-degenerate quadric  $\Omega$  of [5] is obtained by projecting  $V_3^4$  from a point which lies in the solid containing the quadric  $Q$ . We thus obtain a locus  $v_3^4$  with a double plane, the  $\infty^1$  generating planes of  $v_3^4$  meeting this double plane in the tangents of a conic.

The generating planes of  $v_3^4$  are all of the same system on the quadric  $\Omega$ , the double plane being of the opposite system. If the points of  $\Omega$  represent the lines of [3] then the points of  $C$  represent the generators of a rational sextic ruled surface  $f$ . If the double plane of  $v_3^4$  is a  $\rho$ -plane  $f$  has a triple

conic  $C_2$  whose plane is represented on  $\Omega$  by this  $\rho$ -plane. If the double plane of  $v_3^4$  is a  $\varpi$ -plane  $f$  has as tritangent planes the tangent planes of a quadric cone  $E_2$  whose vertex is represented on  $\Omega$  by the  $\varpi$ -plane.

**403.** The four types of sextic ruled surfaces that we have just obtained can be derived by projection either from the normal surface with  $\infty^1$  directrix cubics or from the normal surface with a directrix conic  $\Gamma$ . On the latter surface  $C_6^0$  and  $\Gamma$  have two intersections; these belong, in general, to different sets of the  $g_3^1$  on  $C_6^0$ .

We now proceed to consider the particular case when the two intersections,  $a_1$  and  $a_2$  say, of  $C_6^0$  and  $\Gamma$  belong to the same set of the  $g_3^1$ ; the third point of this set is some other point  $b$  on  $C_6^0$ . The line  $a_1a_2$  meets the quadric  $Q$  on  $V_3^4$ , so that we are projecting from a solid  $S$  which meets the plane of  $\Gamma$  in a point  $O$ . Moreover, the  $[5]S\Gamma$  contains the plane  $a_1a_2b$ , and therefore also the generator  $g_0$  of  $F$  which passes through  $b$ . Hence the projected surface  $f$  has a directrix line  $R$  which is itself a generator; through any point of  $R$  there pass two other generators while any plane through  $R$  contains three other generators.

The generators which meet  $C_6^0$  in the sets of the  $g_3^1$  give also sets of a  $g_3^1$  on  $\Gamma$ ; since a  $g_3^1$  and a  $g_2^1$  on  $\Gamma$  have two common pairs of points there is one pair of points, other than  $a_1$  and  $a_2$ , which belong to a set of the  $g_3^1$  on  $\Gamma$  and have their join passing through  $O$ . The two generators  $g$  and  $g'$  of  $F$  which pass through this pair of points lie in a solid which meets  $S$  in a line  $l$  passing through  $O$ .  $S$  meets  $M_5^6$  in  $l$ ,  $m$  and  $\vartheta_4^0$ , where  $m$  is the line of intersection of  $S$  with the solid  $\Gamma g_0$  and  $\vartheta_4^0$  is the quartic curve on  $Q$ .  $l$ ,  $m$  and  $\vartheta_4^0$  all pass through  $O$ .

The chords of  $F$  which meet  $S$  meet  $F$  in  $C_6^0$ ,  $\Gamma$  and  $g_0$  each counted twice, and in  $g$  and  $g'$ . On projection from  $S$  we obtain a surface  $f$  whose double curve is  $3R + G + 3C_2$ ;  $R$  and  $C_2$  have one intersection and  $G$  meets both  $R$  and  $C_2$ .

The surface is generated by a line  $R$  and a conic  $C_2$  in (3, 2) correspondence with a doubly united point; the correspondence must be specialised to give the double generator. The joins of the pairs of points of  $C_2$  which correspond to the points of  $R$  envelop a rational curve of class three, one point of  $R$  lying on the tangent of the curve to which it gives rise. Hence the planes of the pairs of generators which intersect in the points of  $R$  touch a developable  $E_3^0$  which has two planes passing through  $R$  and one through  $G$ . The bitangent developable of  $f$  is  $6R + G + E_3^0$ .

There is also a dual surface whose double curve is  $6R + G + C_3^0$  and bitangent developable  $3R + G + 3E_2$ .

**404.** In order to obtain a rational sextic ruled surface  $f$  in [3] which has an infinity of tritangent planes we must project the normal surface  $F$ ,

in [7], which contains  $\infty^1$  directrix cubics from a solid  $S$  which meets every solid  $K$  that contains one of the directrix cubics.

The solids  $K$  form\* a locus  $M_4^4$ . This locus has  $\infty^3$  linear directrices, one of these passing through any given point of the locus; it can in fact be generated by a projectivity between any two of its solids  $K$ , the lines which join corresponding points of the two solids being the directrices. The generators of  $F$  are themselves directrices of  $M_4^4$ ; the cubic curves in which the two solids  $K$  meet  $F$  correspond to each other in the projectivity.

Suppose now that we project  $F$  on to a [3]  $\Sigma$  from a solid  $S$  which contains a directrix  $d$  of  $M_4^4$ . Each point of  $d$  lies in a solid  $K$ , and through it there passes a chord of the corresponding cubic curve on  $F$ . These chords all meet  $F$  on the same pair of generators†  $g$  and  $g'$ , and  $d$  lies in the solid  $gg'$ . Hence on projection we obtain a surface  $F$ , with a double generator  $G$ , whose bitangent developable is  $G + 3E_3^0$ ; the solids  $K$  when projected from  $S$  become the osculating planes of a twisted cubic. The double curve of the surface is  $G + C_9^2$ .

There is also the dual surface whose double curve is  $G + 3C_3^0$  and bitangent developable  $G + E_9^2$ . We take a curve of order nine lying on the normal surface and meeting each generator in two points; this curve is of genus 2, and the planes which contain the sets of a  $g_3^1$  on the curve form a locus  $V_3^5$ . There is a quadric‡  $Q$  which has a generator in each plane of  $V_3^5$ , and we project from the solid  $S$  which contains  $Q$ . The chords of the curve which lie in the planes of  $V_3^5$  form a ruled surface of order fourteen§, meeting  $Q$  in a curve of order five which is trisecant to the generators of  $Q$  which lie in the planes of  $V_3^5$ . This quintic curve forms part of the intersection of  $S$  with the locus  $M_5^6$  formed by the chords of the ruled surface; the residual intersection is a line which lies in a solid containing a pair of generators of the ruled surface.

In order to obtain by projection the surfaces which have as tritangent planes the tangent planes of a quadric cone we project from a solid  $S$  which contains a directrix conic of  $M_4^4$ .

**405.** We have seen that the surface which is generated by means of a (3, 2) correspondence, with a doubly united point, between a line  $R$  and a conic  $C_2$  is a rational sextic ruled surface, provided the correspondence

\* p. 143 *supra*.

† Each directrix of  $M_4^4$  lies in a solid containing two generators of  $F$ . Each solid which contains two generators of  $F$  meets  $M_4^4$  in a quadric surface, on which lie  $\infty^1$  directrices of  $M_4^4$ .

‡ There are two types of locus  $V_3^5$ ; a locus of the first type is generated by means of two lines and a cubic curve, a locus of the second type by a line and two conics. We are concerned with a locus of the first type here.  $V_3^5$  is normal in [7].

§ The ruled surface is generated by the joins of pairs of corresponding points in a symmetrical (2, 2) correspondence of valency 1 on a curve of genus 2. See the footnote to § 19.

is specialised to give a double generator. If the correspondence is not specialised to give a double generator the resulting ruled surface is elliptic. We therefore proceed to enquire whether we can obtain an elliptic sextic ruled surface with a triple conic by projection; the surface must be the projection of a normal surface in [5] from a line  $l$  which lies in  $\infty^1$  of its trisecant planes. The triple conic must be the projection from  $l$  of the section of the surface by a prime through  $l$ , there being  $\infty^1$  trisecant planes of this prime section passing through  $l$  and forming a quadric line-cone.

Suppose we have an elliptic normal sextic curve in [5]; through a point of general position there pass two of its trisecant planes. There are four Veronese surfaces containing the curve\*, and through a point of any of these surfaces there pass not two but an infinity of trisecant planes of the curve. The surfaces may be defined by this property. The tangent planes of any one of the Veronese surfaces form a cubic primal, and the four cubic primals so obtained belong to a pencil whose base is the  $M_3^9$  formed by the chords of the curve. The four cubic primals together form the locus of points which are such that the two trisecant planes of the curve which pass through them coincide †.

Take now a point  $O$  on one of the Veronese surfaces. The  $\infty^1$  trisecant planes of the sextic which pass through  $O$  meet the Veronese surface in conics; if the sextic is projected from  $O$  on to a [4] it becomes a curve lying on a cubic ruled surface, meeting every generator in three points. The tangent plane  $\pi$  of the Veronese surface at  $O$  meets all the trisecant planes through  $O$  in lines; the projection from any point of  $\pi$  on to a [4] gives an elliptic sextic curve such that  $\infty^1$  of its trisecant planes pass through a line. The trisecant planes through  $O$  form a cubic cone; so that, after projection from a point of  $\pi$ , we obtain  $\infty^1$  trisecant planes of an elliptic sextic curve in [4] which generate a quadric line-cone.

The chords of the normal curve which lie in the trisecant planes through  $O$  meet  $\pi$  in the points of a cubic curve; we can either project from a general point of  $\pi$  or from a point of this cubic curve.

406. Suppose first that we project from a general point of  $\pi$ . We obtain, in [4], an elliptic sextic curve with only one trisecant; there are  $\infty^1$  trisecant planes of this curve all passing through a line, this line being met also by the trisecant. We therefore consider a prime section  $C_6^1$  of the normal surface  $F$  which has only one directrix cubic  $\Gamma$ . We have on  $C_6^1$  a  $g_3^1$  cut out by trisecant planes through a line  $l$ ; the three points of intersection of  $C_6^1$  and  $\Gamma$  lie on a line meeting  $l$  in a point  $O$ . The chords of  $F$

\* Rosati, *Rend. Ist. Lomb.* (2), 35 (1902), 407.

† There are four quadric cones which contain an elliptic quartic curve in [3]; the two chords of the curve which pass through a point of any of the cones coincide. There is an analogous result for a normal elliptic curve of any even order; that for the sextic is as above.

which meet  $l$  meet  $F$  in  $C_6^1$  and  $\Gamma$  each counted twice; on projection we obtain a surface with a triple directrix line  $R$  and a triple conic  $C_2$ . The double curve is  $3R + 3C_2$ ;  $R$  and  $C_2$  intersect. The surface is generated by a (3, 3) correspondence between  $R$  and  $C_2$  with a trebly united point; to the point of intersection regarded as a point of either curve there correspond three points of the other curve which all coincide with it. The triads of points of  $C_2$  which correspond to the points of  $R$  are such that their joins envelop an elliptic curve of the sixth class; the tangent of  $C_2$  at the united point is a stationary tangent of this envelope, which has three other tangents passing through the point. The bitangent developable of the surface is  $3R + E_6^1$ , there being three planes of  $E_6^1$  passing through  $R$ .

407. If we project from a point of  $\pi$  which lies on a chord of the normal curve we obtain an elliptic sextic curve with a double point in [4]. There is on this curve a  $g_3^1$  cut out by the planes of a quadric line-cone; the two points on the different branches at the double point belong to the same set of the  $g_3^1$  and the line joining them to the remaining point of the set meets the vertex of the line-cone. We therefore consider a prime section  $C_6^1$  of the normal surface  $F$  with a double line  $\lambda$ .  $C_6^1$  has a double point; the plane joining this to the vertex  $l$  of the line-cone meets  $C_6^1$  again, and the generator  $g$  through this other intersection lies in the solid  $l\lambda$ . We obtain, on projection from  $l$ , a surface in [3] with a directrix line  $R$  which is itself a generator; through each point of  $R$  there pass two other generators while each plane through  $R$  contains three other generators.  $R$  is the projection of  $g$  and  $\lambda$ ; there is also a triple conic  $C_2$ , the projection of  $C_6^1$ .  $R$  and  $C_2$  have one intersection; the surface is generated by a (3, 2) correspondence, with a doubly united point, between  $R$  and  $C_2$ . The joins of the pairs of points of  $C_2$  which correspond to the points of  $R$  form a rational envelope of the third class, one of whose tangents passes through the point of  $R$  which gives rise to it. The planes which contain the pairs of generators which intersect in the points of  $R$  therefore form a developable  $E_3^0$  two of whose planes pass through  $R$ . The bitangent developable of the surface is  $6R + E_3^0$ .

408. If we project the normal elliptic surface  $F$  which has  $\infty^1$  directrix cubic curves from a line  $l$  we obtain a surface  $f$  in [3] which has  $\infty^1$  tri-tangent planes. The planes of the cubic curves on the normal surface form a locus  $V_3^3$ , and are projected into the planes of a developable of class three, there being three of them which pass through an arbitrary point of [3]. Hence we obtain a surface whose bitangent developable is  $3E_3^0$ . The double curve is  $C_9^4$ .

The dual surface whose double curve is  $3C_3^0$  is obtained by projection from the normal surface which has two directrix cubic curves; the bi-

tangent developable is  $E_9^4$ . This surface is generated by a symmetrical (3, 3) correspondence between the points of a twisted cubic  $C_3^0$ .

When we represent these surfaces on  $\Omega$  we obtain an elliptic sextic curve with  $\infty^1$  trisecant planes on  $\Omega$ ; these planes are all of the same system, two of them passing through any point of the curve. They form a three-dimensional sextic locus with a double Veronese surface.

When we project the surface  $F$  which contains  $\infty^1$  directrix cubics from a line which meets one of the planes which contain the cubic curves we obtain a surface  $f$  which has as tritangent planes the tangent planes of a quadric cone. These planes are the projections of the planes of a locus  $V_3^3$  from a line  $l$  which meets one of them.

In general the line  $l$ , meeting the plane of a directrix cubic  $\Gamma$ , is not such that the solid which joins it to the plane of  $\Gamma$  contains also a generator of  $F$ . The chords of  $F$  which meet  $l$  meet  $F$  in  $\Gamma$  counted twice and a curve  $C_{12}^2$ , six of whose tangents meet  $l$ , meeting each generator in two points. The projected surface has a double curve  $3R + C_6^1$  and a bitangent developable  $3R + 3E_2$ ,  $R$  being the projection of  $\Gamma$ .

If we choose  $l$  to lie in the solid which contains  $\Gamma$  and a generator  $g$  then the chords of  $F$  which meet  $l$  meet  $F$  in  $\Gamma$  and  $g$  each counted three times and a prime section  $C_6^1$ . We obtain a surface  $f$  with a directrix  $R$  which is also a generator. The double curve is  $6R + C_3^0$  and the bitangent developable  $3R + 3E_2$ .

409. The following three surfaces were overlooked in the third section of Chapter IV.

The chords of a normal sextic curve  $C$  of genus 2 in [4] which lie on a quadric containing  $C$  set up on  $C$  a symmetrical (4, 4) correspondence. If this breaks up into the sum of two (2, 2) correspondences we have in [3] a ruled surface, normal in [3], whose double curve is  $C_4^1 + D_4^1$  and bitangent developable  $E_4^1 + F_4^1$ .  $C_4^1$  and  $D_4^1$  have four common points;  $E_4^1$  and  $F_4^1$  have four common planes.

Suppose that we take a prime section  $C_6^2$  of the normal surface  $F$  (§ 335) which lies on a cubic cone\* and project  $F$  on to a [3] from the vertex  $O$  of this cone. There are two tangents of  $C_6^2$  through  $O$ ; the remaining chords of  $F$  through  $O$  meet  $F$  in a curve  $C_8^5$  having two intersections with each generator, eight tangents of  $C_8^5$  passing through  $O$ . We obtain in [3] a surface whose double curve is  $R + C_3^1 + C_4^1$  and bitangent developable  $6R + E_2$ ; it is generated by a (2, 2) correspondence between  $R$  and  $C_3^1$  with a doubly united point.  $C_4^1$  meets  $C_3^1$  four times but does not meet  $R$ .

There is also the dual surface, normal in [3], whose double curve is  $6R + C_2$  and bitangent developable  $R + E_3^1 + E_4^1$ .

\* The intersection of an elliptic cubic cone with a cubic surface containing three of its generators and touching it is a  $C_6^2$  with a double point.

## TABLES SHEWING THE DIFFERENT TYPES OF RULED SURFACES IN [3] UP TO AND IN- CLUDING THOSE OF THE SIXTH ORDER

We give here a set of tables shewing the different kinds of ruled surfaces in [3] that we have studied. In the first two columns of a table we give the double curve and bitangent developable of the surface, following the model of the table at the end of Cremona's paper on quartic ruled surfaces.

The symbols  $B_n^p$ ,  $C_n^p$ ,  $D_n^p$  denote curves of order  $n$  and genus  $p$ , while  $E_n^p$ ,  $F_n^p$ ,  $G_n^p$  denote developables of class  $n$  and genus  $p$ .  $R$  is used for a directrix,  $G$  and  $H$  for generators.

We shall not give here the developable surfaces, since a table of these is on p. 284.

### *Cubic Ruled Surfaces*

The cubic ruled surfaces are of only two kinds, both are rational and are projections of a normal surface  $F$  in [4] with a directrix line  $\lambda$ . They can be exhibited as follows:

Double curve	Bitangent developable
$2R$	$2R'$
$2R$	$2R$

### *Quartic Ruled Surfaces*

When we consider quartic ruled surfaces there are ten rational surfaces and two elliptic surfaces. Of the ten rational surfaces six are projections of the normal surface in [5] with directrix conics, while the remaining four are projections of the normal surface in [5] with a directrix line.

We give also in the table the type of rational quartic  $C$  on  $\Omega$  from which any rational ruled quartic surface arises (the list of these curves  $C$  is found in § 58) and also the numbers attached to the surfaces by Cremona and Cayley. The last two surfaces in the table are elliptic.



Double curve	Bitangent developable	Type of curve $C$ on $\Omega$	Order of minimum directrix on normal surface $F$	Cremona's type	Cayley's type
$C_3^0$	$E_3^0$	I	2	1	10
$3R$	$E_3^0$	II (A)	2	8	9
$C_3^0$	$3R$	II (B)	1	7	8
$R + C_2$	$R + E_2$	II (C)	2	2	7
$3R$	$R + E_2$	III (A)	2	3	—
$R + C_2$	$3R$	III (B)	1	4	—
$3R$	$3R'$	IV (A)	1	9	3
$R + G + R'$	$R + G + R'$	IV (B)	2	5	2
$2R + G$	$2R + G$	V (A)	2	6	5
$3R$	$3R$	V (B)	1	10	6
<hr/>					
$R + R'$	$R + R'$	—	—	11	1
$2R$	$2R$	—	—	12	4

*Quintic Ruled Surfaces*

Of quintic ruled surfaces in [3] there are twenty-four kinds of rational surfaces, six kinds of elliptic surfaces, and two surfaces whose plane sections are of genus 2.

Of the rational surfaces seventeen are projections of the normal surface in [6] with a directrix conic and seven are projections of the normal surface in [6] with a directrix line.

For the purpose of classifying the rational quintic surfaces we distinguished nineteen different types of rational quintic curves  $C$  on  $\Omega$ . We give the type of curve corresponding to each surface (a list of the curves is given in § 85) and also the number associated with the surface by Schwarz\*.

The table is divided into four sections by horizontal lines drawn across it. The surfaces in the first section have not a directrix line; those in the second section have a directrix line which is not a generator; those in the third section have a directrix line which is also a generator, while those in the fourth section are surfaces whose generators belong to a linear congruence.

\* *Journal für Math.* 67 (1867), 57.

Double curve	Bitangent developable	Type of curve $C$ on $\Omega$	Order of minimum directrix on normal surface $F$	Schwarz's type
$C_6^1$ $C_2 + C_4^0$ $B_2 + C_2 + D_2$ $G + C_5^0$	$E_6^1$ $E_2 + E_4^0$ $E_2 + F_2 + G_2$ $G + E_5^0$	I, II (A) I, II (A) I, II (A) II (B)	2 2 2 2	II VIII IX X
$6R$ $6R$ $6R$ $C_6^1$ $C_2 + C_4^0$ $B_2 + C_2 + D_2$ $3R + G + C_2$ $R + G + E_4^0$ $3R + C_3^0$ $R + C_5^0$	$E_6^1$ $E_2 + E_4^0$ $E_2 + F_2 + G_2$ $6R$ $6R$ $6R$ $R + G + E_4^0$ $3R + G + E_2$ $R + E_5^0$ $3R + E_3^0$	III (A) III (A) III (A) III (B) III (B) III (B) III (B) III (C) III (D) III (E) III (F)	2 2 2 1 1 1 1 2 2 2 2	I I I II VIII IX IV VII III VI
$6R$ $R + C_5^0$ $3R + G + C_2$ $3R + C_3^0$ $6R$ $3R + C_3^0$	$R + E_5^0$ $6R$ $3R + G + E_2$ $3R + E_3^0$ $3R + E_3^0$ $6R$	IV (A) IV (B) IV (C) IV (D) V (A) V (B)	2 1 2 2 2 1	I VI IV III I III
$6R$ $3R + G + H + R'$ $4R + G + H$ $6R$	$6R'$ $R + G + H + 3R'$ $4R + G + H$ $6R$	VI (A) VI (B) VII (A) VII (B)	1 2 2 1	I V V I

To classify the elliptic quintic ruled surfaces in [3] we divided the elliptic quintic curves  $C$  on  $\Omega$  into six classes (§ 151). Of the six types of surface two are projections of the normal surface with directrix cubic curves, while four are projections of the normal surface with a double line.

We have the following table:

Double curve	Bitangent developable	Type of curve $C$ on $\Omega$	Order of minimum directrix on normal surface $F$	Schwarz's type
$C_5^1$	$E_5^1$	VIII	3	I
$3R + C_2$	$R + E_4^1$	IX (A)	3	II
$R + C_4^1$	$3R + E_2$	IX (B)	2	IV
$3R + C_2$	$3R + E_2$	X	2	II
$3R + G + R'$	$R + G + 3R'$	XI	2	II
$4R + G$	$4R + G$	XII	2	III

There are only two types of quintic ruled surface in [3] whose plane sections are of genus 2; they are not projections of quintic surfaces in higher space. They are given by

Double curve	Bitangent developable	Type of curve $C$ on $\Omega$
$3R + R'$	$R + 3R'$	XIII
$4R$	$4R$	XIV

*Sextic Ruled Surfaces*

Proceeding now to the sextic ruled surfaces we have eighty-three kinds of rational surfaces, thirty-four kinds of elliptic surfaces, thirteen kinds of surfaces whose plane sections are of genus 2, four kinds of surfaces whose plane sections are of genus 3 and two kinds of surfaces whose plane sections are of genus 4.

As a basis for the classification of the rational surfaces we have thirty-eight types of rational curves  $C$  on  $\Omega$  (§ 171).

Corresponding to curves  $C$  of types I and II we have twenty-seven different kinds of ruled surfaces; twelve of these are without multiple generators. We have the following table:

Double curve	Bitangent developable	Type of curve $C$ on $\Omega$
$C_{10}^3$ $C_5^0 + D_5^0$ $C_4^0 + C_6^0$ $C_2 + C_8^1$ $C_3^0 + C_7^1$ $C_2 + D_2 + C_6^0$ $C_3^0 + D_3^0 + C_4^0$ $C_2 + C_3^0 + C_5^0$ $3C_2 + C_4^0$ $3B_2 + C_2 + D_2$ $C_4^0 + C_6^0$ $C_4^0 + C_3^0 + D_3^0$	$E_{10}^3$ $E_5^0 + F_5^0$ $E_6^0 + E_4^0$ $E_3^0 + E_7^1$ $E_2 + E_8^1$ $E_3^0 + F_3^0 + E_4^0$ $E_2 + F_2 + E_3^0$ $E_3^0 + E_2 + E_5^0$ $E_4^0 + E_6^0$ $E_4^0 + E_3^0 + F_3^0$ $3E_2 + E_4^0$ $3E_2 + F_2 + G_2$	I (A)    II (A) I (A) I (A) I (A) I (A) I (A) I (A) I (A) I (A) I (A) I (A) I (A)
$G + C_9^2$ $C_2 + G + C_7^1$ $C_2 + D_2 + G + C_5^0$ $C_4^0 + G + C_5^0$ $C_2 + G + C_7^0$ $C_3^0 + G + C_6^0$ $C_2 + D_2 + G + C_5^0$ $C_2 + C_3^0 + G + C_4^0$ $G + 3C_3^0$ $G + C_9^2$ $G + H + C_8^1$ $C_4^0 + G + H + D_4^0$ $C_2 + G + H + C_6^0$ $C_2 + D_2 + G + H + C_4^0$ $3G + C_7^0$	$G + E_9^2$ $E_2 + G + E_7^1$ $E_2 + F_2 + G + E_5^0$ $E_5^0 + G + E_4^0$ $E_3^0 + G + E_6^0$ $E_2 + G + E_7^0$ $E_2 + E_3^0 + G + E_4^0$ $E_2 + F_2 + G + E_5^0$ $G + E_9^2$ $G + 3E_3^0$ $G + H + E_8^1$ $E_4^0 + G + H + F_4^0$ $E_2 + G + H + E_6^0$ $E_2 + F_2 + G + H + E_4^0$ $3G + E_7^0$	I (B)    II (B) I (B)    II (B) I (B)    II (B) I (B) I (B) I (B) I (B) I (B) I (B) I (B) I (B) I (B) I (B) I (B) II (C) II (C) II (C) II (C) II (C) II (D)

The remaining types of curves  $C$  on  $\Omega$  give rise to fifty-six kinds of surfaces; each of these surfaces is the projection of a unique surface in [7], twenty-seven are projections of the normal surface with directrix cubic curves, twenty are projections of the normal surface with a directrix conic and nine are projections of the normal surface with a directrix line. They are exhibited in the following table. The table is divided into three sections; the surfaces in the first section have a directrix line which is not itself a generator, those in the second section have a directrix line which is also a generator and those in the third section are such that their generators belong to a linear congruence.

Double curve	Bitangent developable	Type of curve $C$ on $\Omega$	Order of minimum directrix on normal surface $F$
$10R$	$E_{10}^3$	III (A)	3
$10R$	$E_5^0 + F_5^0$	III (A)	3
$C_{10}^3$	$10R$	III (B)	1
$C_5^0 + D_5^0$	$10R$	III (B)	1
$6R + C_4^0$	$R + E_9^1$	III (C)	3
$6R + C_4^0$	$R + E_3^0 + E_6^0$	III (C)	3
$6R + C_4^0$	$R + E_3^0 + F_3^0 + G_3^0$	III (C)	3
$R + C_9^1$	$6R + E_4^0$	III (D)	2
$R + C_3^0 + C_6^0$	$6R + E_4^0$	III (D)	2
$R + B_3^0 + C_3^0 + D_3^0$	$6R + E_4^0$	III (D)	2
$6R + G + C_3^0$	$R + G + E_8^1$	III (E)	3
$6R + G + C_3^0$	$R + G + E_2 + E_6^0$	III (E)	3
$6R + G + C_3^0$	$R + G + E_3^0 + E_5^0$	III (E)	3
$6R + G + C_3^0$	$R + G + E_2 + E_3^0 + F_3^0$	III (E)	3
$R + G + C_8^1$	$6R + G + E_3^0$	III (F)	2
$R + G + C_2 + C_6^0$	$6R + G + E_3^0$	III (F)	2
$R + G + C_3^0 + C_5^0$	$6R + G + E_3^0$	III (F)	2
$R + G + C_2 + C_3^0 + D_3^0$	$6R + G + E_3^0$	III (F)	2
$6R + G + H + C_2$	$R + G + H + E_7^1$	III (G)	3
$6R + G + H + C_2$	$R + G + H + E_2 + E_5^0$	III (G)	3
$6R + G + H + C_2$	$R + G + H + E_3^0 + E_4^0$	III (G)	3
$6R + G + H + C_2$	$R + G + H + E_2 + F_2 + E_3^0$	III (G)	3
$R + G + H + C_7^1$	$6R + G + H + E_2$	III (H)	2
$R + G + H + C_2 + C_5^0$	$6R + G + H + E_2$	III (H)	2
$R + G + H + C_3^0 + C_4^0$	$6R + G + H + E_2$	III (H)	2
$R + G + H + C_2 + D_2 + C_3^0$	$6R + G + H + E_2$	III (H)	2
$3R + C_7^0$	$3R + E_7^0$	III (I)	3
$3R + G + C_6^0$	$3R + G + E_6^0$	III (J)	3
$3R + G + H + C_5^0$	$3R + G + H + E_5^0$	III (K)	3
$3R + 3G + C_4^0$	$3R + 3G + E_4^0$	III (L)	3

Double curve	Bitangent developable	Type of curve $C$ on $\Omega$	Order of minimum directrix on normal surface $F$
$10R$	$R + E_9^1$	IV (A)	3
$10R$	$R + E_3^0 + E_6^0$	IV (A)	3
$10R$	$R + E_3^0 + F_3^0 + G_3^0$	IV (A)	3
$R + C_9^1$	$10R$	IV (B)	1
$R + C_3^0 + C_6^0$	$10R$	IV (B)	1
$R + B_3^0 + C_3^0 + D_3^0$	$10R$	IV (B)	1
$6R + C_4^0$	$3R + E_7^0$	IV (C)	3
$3R + C_7^0$	$6R + E_4^0$	IV (D)	2
$6R + G + C_3^0$	$3R + G + E_5^0$	IV (E)	3
$6R + G + C_3^0$	$3R + G + 3E_2$	IV (E)	3
$3R + G + C_6^0$	$6R + G + E_3^0$	IV (F)	2
$3R + G + 3C_2$	$6R + G + E_3^0$	IV (F)	2
$6R + G + H + C_2$	$3R + G + H + E_5^0$	IV (G)	3
$3R + G + H + C_5^0$	$6R + G + H + E_2$	IV (H)	2
$10R$	$3R + E_7^0$	V (A)	3
$3R + C_7^0$	$10R$	V (B)	1
$6R + C_4^0$	$6R + E_4^0$	V (C)	2
$6R + G + C_3^0$	$6R + G + E_3^0$	V (D)	2
$10R$	$6R + E_4^0$	VI (A)	2
$6R + C_4^0$	$10R$	VI (B)	1
$10R$	$10R'$	VII (A)	1
$6R + G + H + K + R'$	$R + G + H + K + 6R'$	VII (B)	2
$3R + G + H + J + K + 3R'$	$3R + G + H + J + K + 3R'$	VII (C)	3
$10R$	$10R$	VIII (A)	1
$7R + G + H + K$	$7R + G + H + K$	VIII (B)	2
$6R + G + H + J + K$	$6R + G + H + J + K$	VIII (C)	3

Of the thirty-four types of elliptic sextic ruled surfaces in [3] sixteen are projections of the general normal surface which has two directrix cubic curves and eleven are projections of the normal surface with a double line. Five types are projections of the normal surface with  $\infty^1$  plane cubics on it (denoted in the table by  $3_\infty$ ) and the two remaining types are the projections of the normal surface with only one cubic curve on it (denoted in the table by  $3_1$ ).

There are twenty different types of elliptic sextic curves  $C$  on  $\Omega$  (§ 255).

The table of the elliptic sextic ruled surfaces is divided into four sections by horizontal lines drawn across it. The surfaces in the first section have not a directrix line; those in the second section have a directrix

line which is not itself a generator; those in the third section have a directrix line which is also a generator; those in the fourth section belong to a linear congruence.

Double curve	Bitangent developable	Type of curve $C$ on $\Omega$	Order of minimum directrix on normal surface $F$
$C_9^4$ $C_2 + C_7^3$ $C_2 + D_2 + C_5^2$ $3C_3^0$ $C_9^4$ $G + C_8^3$ $C_2 + G + C_6^2$	$E_9^4$ $E_2 + E_7^3$ $E_2 + F_2 + E_5^2$ $E_9^4$ $3E_3^0$ $G + E_8^3$ $E_2 + G + E_6^2$	I, II (A) I, II (A) I, II (A) I I II (B) II (B)	3 3 3 3 $3_\infty$ 3 3
$R + C_8^3$ $R + C_2 + C_6^2$ $R + C_2 + D_2 + C_4^1$ $6R + C_3^0$ $6R + C_3^0$ $6R + C_3^0$ $R + G + C_7^3$ $R + G + C_2 + C_5^2$ $6R + G + C_2$ $6R + G + C_2$ $3R + C_4^1$ $3R + 3C_2$ $3R + C_6^1$ $3R + G + C_5^1$	$6R + E_3^0$ $6R + E_3^0$ $6R + E_3^0$ $R + E_8^3$ $R + E_2 + E_6^2$ $R + E_2 + F_2 + E_4^1$ $6R + G + E_2$ $6R + G + E_2$ $R + G + E_7^3$ $R + G + E_2 + E_5^2$ $3R + E_6^1$ $3R + E_6^1$ $3R + 3E_2$ $3R + G + E_5^1$	III (A) III (A) III (A) III (B) III (B) III (B) III (C) III (C) III (D) III (D) III (E) III (E) III (E) III (F)	2 2 2 3 3 3 2 2 3 3 3 3 3 3 3
$3R + C_6^1$ $3R + 3C_2$ $6R + C_3^0$ $6R + C_3^0$ $3R + G + C_5^1$ $6R + G + C_2$ $6R + C_3^0$	$6R + E_3^0$ $6R + E_3^0$ $3R + E_6^1$ $3R + 3E_2$ $6R + G + E_2$ $3R + G + E_5^1$ $6R + E_3^0$	IV (A) IV (A) IV (B) IV (B) IV (C) IV (D) V	2 2 3 $3_\infty$ 2 3 2
$6R + G + H + R'$ $3R + G + H + J + 3R'$ $3R + 3G + 3R'$ $7R + G + H$ $6R + G + H + J$ $6R + 3G$	$R + G + H + 6R'$ $3R + G + H + J + 3R'$ $3R + 3G + 3R'$ $7R + G + H$ $6R + G + H + J$ $6R + 3G$	VI (A) VI (B) VI (C) VII (A) VII (B) VII (C)	2 3 $3_\infty$ 2 3 $3_\infty$

We give now the list of the sextic ruled surfaces in [3] whose plane sections are of genus 2, classified according to their double curves and bitangent developables.

Double curve	Bitangent developable	Type of curve $C$ on $\Omega$
$C_8^5$	$E_8^5$	I
$C_4^1 + D_4^1$	$E_4^1 + F_4^1$	I
$R + C_7^5$	$6R + E_2$	II (A)*
$R + C_3^1 + C_4^1$	$6R + E_2$	II (A)*
$6R + C_2$	$R + E_7^5$	II (B)
$6R + C_2$	$R + E_3^1 + E_4^1$	II (B)
$3R + C_5^2$	$3R + E_5^2$	II (C)
$3R + C_5^2$	$6R + E_2$	III (A)*
$6R + C_2$	$3R + E_5^2$	III (B)
$6R + G + R'$	$R + G + 6R'$	IV (A)*
$3R + G + H + 3R'$	$3R + G + H + 3R'$	IV (B)
$7R + G$	$7R + G$	V (A)*
$6R + G + H$	$6R + G + H$	V (B)

The list of curves  $C$  on  $\Omega$  of order 6 and genus 2 is found in § 327.

The five surfaces marked with an asterisk are the projections of a normal surface in [4], but the others are not the projections of a sextic ruled surface in higher space, and are themselves normal.

For surfaces whose plane sections are of genus 3 we have

Double curve	Bitangent developable
$6R + R'$	$R + 6R'$
$3R + G + 3R'$	$3R + G + 3R'$
$7R$	$7R$
$6R + G$	$6R + G$

and for surfaces whose plane sections are of genus 4

Double curve	Bitangent developable
$3R + 3R'$	$3R + 3R'$
$6R$	$6R$



## NOTE

### THE INTERSECTIONS OF TWO CURVES ON A RULED SURFACE

1. Suppose that we have in space  $[r]$  a ruled surface of order  $n$  and on this surface two curves of orders  $m$  and  $m'$ . A curve will meet all the generators of the surface in the same number of points; suppose then that the curve of order  $m$  meets each generator in  $k$  points and that the curve of order  $m'$  meets each generator in  $k'$  points. We wish to find an expression for the number of intersections of the two curves.

In the first instance we shall assume that each curve is a simple curve on the ruled surface; this means that through each point of a curve, with possibly a finite number of exceptions, there passes only one generator of the surface.

Take then an arbitrary  $[r - 2]$ ; this will meet the surface in  $n$  points. We establish a correspondence between primes  $P$  and  $P'$  passing through this  $[r - 2]$ ; two primes  $P$  and  $P'$  correspond when  $P$  joins  $[r - 2]$  to a point of the first curve and  $P'$  joins  $[r - 2]$  to a point of the second curve, these two points being on the same generator of the surface. Then any prime  $P$  meets the curve of order  $m$  in  $m$  points through each of which there passes a generator of the surface, so that we have  $mk'$  corresponding primes  $P'$ . Similarly, given a prime  $P'$ , we have  $m'k$  corresponding primes  $P$ . Hence, by Chasles' principle of correspondence, there will be  $mk' + m'k$  coincidences of pairs of corresponding primes.

The prime joining  $[r - 2]$  to an intersection of the two curves clearly counts among these coincidences. But there are also other coincidences; for  $[r - 2]$  meets the surface in  $n$  points, and the prime joining  $[r - 2]$  to the generator of the surface through any one of these points contains  $k$  points of the first curve and  $k'$  points of the second curve, all on this same generator. This prime in fact counts  $kk'$  times among the coincidences. There are no other coincidences than those already mentioned.

Hence we conclude that, if  $i$  is the number of intersections of the two curves\*,

$$mk' + m'k = i + nkk',$$

or

$$i = mk' + m'k - nkk'.$$

For example, if we consider two prime sections we have  $m = m' = n$  and  $k = k' = 1$ , so that they have  $n$  intersections. These are, of course, the  $n$  points in which the  $[r - 2]$  common to the primes meets the surface.

\* Segre, *Rom. Acc. Lincei Rend.* (4), 3<sup>a</sup> (1887), 3.

We have assumed implicitly in the above proof that none of the intersections of the two curves is a multiple point on the ruled surface; for if one of the  $i$  points is at an intersection of two or more generators of the ruled surface we should have to count it more than once among the coincidences, although both the curves might only have simple points there. In such a case the number  $i$  given by the formula would have to be modified.

2. Let us consider now the intersections of two multiple curves on a ruled surface. We take as before a ruled surface of order  $n$  in  $[r]$  and on it two curves of orders  $m$  and  $m'$ , meeting each generator of the surface in  $k$  and  $k'$  points respectively. But now the curves are of multiplicities  $s$  and  $s'$  upon the surface; through a general point of the curve of order  $m$  there pass  $s$  generators, while through a general point of the curve of order  $m'$  there pass  $s'$  generators. The preceding is the particular case of this when  $s = s' = 1$ .

We set up just as before a correspondence between the primes  $P$  and  $P'$  of a pencil; but now to any given prime  $P$  there will correspond  $msk'$  primes  $P'$ , while to any given prime  $P'$  there will correspond  $m's'k$  primes  $P$ . We thus have

$$msk' + m's'k - nkk'$$

coincidences of pairs of corresponding primes which are to be accounted for by intersections of the two curves.

The question then remains to be answered, "How many times does each intersection count among this number of coincidences?"—and at first sight it seems as though the answer cannot be given without some difficulty. There may be points of the first curve through which there pass more than  $s$  generators of the surface just as there may be points of the second curve through which there pass more than  $s'$  generators of the surface. It seems as though we ought to know the number of generators which meet at any given intersection of the two curves before we can tell how many times this intersection should count among the coincidences, and we may have to know more than this.

3. It will help to explain some of the difficulties if we illustrate them by one or two examples.

(A) Suppose that we take a line and a conic with one intersection and place them in (2, 2) correspondence with a doubly united point. The joining lines will then generate a ruled surface of the fourth order on which the line and the conic are double curves. For the line  $m = 1$ ,  $s = 2$ ,  $k = 1$ , while for the conic  $m' = 2$ ,  $s' = 2$ ,  $k' = 1$ , and  $n = 4$ . Hence

$$msk' + m's'k - nkk' = 2,$$

and the single intersection is counted twice.

(B) If we consider the number of intersections of a prime section and a double curve the formula for the number of coincidences gives twice the

order of the double curve, whereas the number of intersections is actually equal to the order of the double curve. Each intersection then must be counted twice.

(C) Let us take a rational quartic with a double point and a conic which passes through the double point and meets the quartic in two other points. Then the chords of the quartic which meet the conic generate a quintic ruled surface on which the two curves are double curves. The double point of the quartic curve is, in fact, a triple point of the ruled surface, there being three generators passing through this point.

Then for the quartic

$$m = 4, \quad s = 2, \quad k = 2,$$

and for the conic

$$m' = 2, \quad s' = 2, \quad k' = 1,$$

while  $n = 5$ . Hence

$$msk' + m's'k - nkk' = 6.$$

But the number of actual intersections of the two curves is four, if we count the double point as two intersections, and the relation between the number of intersections and the number of coincidences does not seem clear without further investigation.

Now it is known, from the general theory of ruled surfaces, that on this quintic ruled surface there is a simple conic; through each point of this conic not on the double curve there passes one generator of the surface, while each generator (with three exceptions) meets the conic in one point. Then for this conic  $m'' = 2$ ,  $s'' = 1$ ,  $k'' = 1$ , so that, considering the two conics,

$$m's'k'' + m''s''k' - nk'k'' = 1.$$

It certainly cannot be maintained that any coincidence has been counted twice in this result, and there is in fact precisely one intersection of the two conics. Hence in this particular case the formula gives the correct result without modification.

Also

$$m''s''k + msk'' - nk''k = 2,$$

which is also correct; the simple conic does, in fact, meet the double quartic in two points.

But if we were to calculate the number of intersections of either the double conic or the double quartic with a plane section we should get twice the correct result, as in (B).

Thus it appears that when we have two curves of given multiplicities on a ruled surface we cannot, without further examination, state the number of times that each intersection counts among the coincidences. We certainly cannot for two double curves, or for a double curve and a simple curve.

4. It may be that the ruled surface which we are considering is the projection of a normal ruled surface in higher space. When this is so it

may happen further that the curves of orders  $m$  and  $m'$  and multiplicities  $s$  and  $s'$  are the projections of simple curves of orders  $ms$  and  $m's'$  on the normal surface. Then, since the order of the surface and the number of points in which a curve meets a generator are unaltered by projection, the formula

$$i = msk' + m's'k - nkk'$$

gives the number of intersections of these two simple curves on the normal surface\*. Thus if an intersection of the multiple curves is the projection of only one of the intersections of the simple curves it will count only once among the coincidences; if, however, it is the projection of two intersections of the simple curves it will count twice, and so on.

We can thus clearly explain the fact that the formula, when applied to a prime section and a double curve, gives twice the correct number of intersections. We assume that the surface in  $[r]$  is the projection of a normal surface of the same order in  $[R]$  from a space  $[R - r - 1]$  not meeting this normal surface. Then we suppose that the double curve of the surface in  $[r]$  is the projection from  $[R - r - 1]$  of a simple curve  $C$  on the normal surface; there will be  $\infty^1$  chords of  $C$  meeting  $[R - r - 1]$ , each of these chords being joined to  $[R - r - 1]$  by an  $[R - r]$  meeting  $[r]$  in a point of the double curve. A prime section of the surface in  $[r]$  is joined to  $[R - r - 1]$  by a space  $[R - 1]$  which gives the corresponding prime section  $S$  of the normal surface; then through any intersection of  $S$  and  $C$  there must pass a chord meeting  $[R - r - 1]$ ; the other intersection of this chord with  $C$  then necessarily lies also on  $S$ . Hence each intersection of the double curve and the prime section on the surface in  $[r]$  is the projection of two intersections of  $S$  and  $C$ .

5. It will be advisable now to elaborate at some length the account of the curves on the quintic surface in our third example; the whole theory of the quintic ruled surfaces has been fully investigated, and this process of elaboration will throw much light on more general examples†.

We had a rational quartic  $C_4$  with a double point and a conic  $C_2$  passing through this double point and meeting  $C_4$  in two other points. The quintic ruled surface formed by the chords of  $C_4$  which meet  $C_2$  has both  $C_4$  and  $C_2$  for double curves; it is rational since its plane sections have six double points. It can therefore be regarded as the projection of a normal quintic ruled surface  $F$  in  $[6]$ .

The centre of projection will be a plane  $\omega$ . The surface  $F$  will have a directrix conic  $\Gamma$ , and the projection of  $\Gamma$  from  $\omega$  gives the simple conic  $c_2$  on the projected surface. The surface  $F$  has  $\infty^4$  directrix quartic curves lying on it, and we choose  $\omega$  to contain an axis  $p$  of one of these quartics  $E$  (§ 130).

\* By § 1.

† See § 130.

It is known that the chords of  $F$  form a locus  $M_5^3$  of five dimensions and the third order; thus  $\omega$  will meet this locus in the line  $p$  and a conic  $\mathfrak{D}$ . Through every point of  $p$  there passes a chord of  $E$ , while through every point of  $\mathfrak{D}$  there passes a chord of  $F$ ; for different points of  $\mathfrak{D}$  the different chords of  $F$  trace out on  $F$  a curve  $C_8$  of the eighth order meeting each generator in two points. It can be proved that  $C_8$  has one double point  $A$ . Through  $A$  there will then pass two chords  $AB$  and  $AC$  of  $C_8$  which both meet  $\mathfrak{D}$ ; the line  $BC$  is then also a chord of  $F$  meeting  $\omega$ , its intersection with  $\omega$  lies on  $p$ , and  $B$  and  $C$  lie on  $E$  as well as on  $C_8$ . Also through the intersections of  $p$  and  $\mathfrak{D}$  there pass lines which are common chords of  $E$  and  $C_8$ .

When we project from  $\omega$  on to a solid the curve  $E$  becomes a conic  $C_2$  and the curve  $C_8$  becomes a quartic  $C_4$ ; both  $C_2$  and  $C_4$  are double curves on the projected surface. The solid through  $\omega$  containing  $A, B, C$  meets the solid on to which we are projecting in a point which lies on  $C_2$  and is a double point on  $C_4$ , this being the projection of two intersections, viz.  $B$  and  $C$ , of  $E$  and  $C_8$ .  $C_2$  and  $C_4$  have two other intersections each of which is the projection of two intersections of  $E$  and  $C_8$ .

We now see clearly how to reckon the intersections of  $C_2$  and  $C_4$ ; the double point of  $C_4$  lies on  $C_2$  and counts twice among the coincidences, while each other intersection of  $C_2$  and  $C_4$  counts twice among the coincidences. The curves  $C_8$  and  $E$  on the normal surface have six intersections and no more.

The curve  $E$ , meeting each generator of  $F$  in one point, will meet the conic  $\Gamma$  in one point, while the curve  $C_8$ , meeting each generator of  $F$  in two points, will meet  $\Gamma$  in two points. Through the intersection of  $\Gamma$  and  $E$  there passes a chord of  $E$  meeting  $p$ , but the other intersection of this chord with  $E$  is not on  $\Gamma$  nor does the solid through  $\omega$  which contains it meet  $\Gamma$  again. Hence, on the quintic surface in [3], the double conic  $C_2$  and the simple conic  $c_2$  have one intersection, and this is the projection of the single intersection of  $\Gamma$  and  $E$ . Similarly the double quartic  $C_4$  meets the simple conic  $c_2$  in two points each of which is the projection of one intersection of  $\Gamma$  and  $C_8$ .

If, however, we were to consider intersections of either  $C_4$  or  $C_2$  with a plane section each would be the projection of two intersections of  $C_8$  or  $E$  with a prime section of  $F$ . The prime would contain  $\omega$ , two chords of  $E$  meeting  $p$  and four chords of  $C_8$  meeting  $\mathfrak{D}$ .

6. It is usual to assume that a general ruled surface has no singularities unless it lies in three or four-dimensional space; a ruled surface in [4] will have a finite number of double points, while a ruled surface in [3] has a double curve with a finite number of triple points which are also triple points of the ruled surface. This is, of course, a statement which is only

true in very general instances; we can have many ruled surfaces with multiple curves in any space. For example, if a curve of order  $n$  lies on a quadric then the chords of the curve which lie on the quadric form a ruled surface on which the curve is a multiple curve of multiplicity  $n - 2$ . Also any ruled surface generated by any correspondence other than a (1, 1) correspondence between two curves has necessarily one or both of the curves as multiple curves.

Let us, however, for the present confine our attention to surfaces in [3] with only double curves. It may happen, as in the case of the quintic ruled surface which we have been discussing, that the double curve is composite, consisting of two or more parts. We shall examine in detail how the intersections of two of these parts are given in the expression

$$msk' + m's'k - nkk'.$$

7. Suppose that the surface  $f$  in [3] is the projection of a normal surface  $F$  in  $[r]$  from a space  $[r - 4]$  not meeting  $F$ . Then a point of the double curve of  $f$  is the intersection with [3] of an  $[r - 3]$  containing  $[r - 4]$  and meeting  $F$  in two points; the chord of  $F$  joining these two points will meet  $[r - 4]$ . There are, in fact,  $\infty^1$  chords of  $F$  meeting  $[r - 4]$ ; of the  $\infty^{2(r-1)}$  lines in  $[r]$  there are  $\infty^{2r-5}$  meeting  $[r - 4]$  and  $r - 3$  conditions are necessary for one of these to meet  $F$ . The curve in which these chords of  $F$  meet  $[r - 4]$  (if  $r > 4$ ) is in birational correspondence with the double curve of  $f$ .

If the double curve of  $f$  has a triple point we shall have an  $[r - 3]$  containing  $[r - 4]$  and meeting  $F$  in three points; thus we have a trisecant plane of  $F$  meeting  $[r - 4]$  in a line. Of the  $\infty^{3(r-2)}$  planes of  $[r]$  there are  $\infty^{3(r-4)}$  meeting  $[r - 4]$ , and  $r - 4$  conditions are necessary for one of these to meet  $F$ . We thus expect  $f$  to have a finite number of triple points. In general,  $F$  will not have any trisecant lines, and, even if it had, none of them would meet a general  $[r - 4]$ .

The chords of  $F$  meeting  $[r - 4]$  will then meet  $F$  in the points of a curve. If a trisecant plane meets  $F$  in points  $A, B, C$  and also meets  $[r - 4]$  in a line the chords  $BC, CA, AB$  will all meet  $[r - 4]$  so that the curve on  $F$  has double points at each of  $A, B, C$ .

If now the double curve of  $f$  is composite this curve on  $F$  will also be composite. It is then at once clear that *if the two parts of the double curve on  $f$  have an intersection which is a simple point on both of them and not a triple point of  $f$  then this intersection is the projection of two distinct intersections of the corresponding curves on  $F$ , the common chord of the two curves meeting  $[r - 4]$ .*

It remains to consider the intersection of the two parts at a triple point of  $f$ . Suppose that the first component of the double curve has a double point here, then the second component will only have a simple

point. Then, the triple point being the intersection of [3] with an  $[r - 3]$  meeting  $F$  in three points  $A, B, C$ , the first component of the curve on  $F$  will have a double point at one of these points, say  $A$ , and pass through the other two, while the second component passes through  $B$  and  $C$ . Thus *the intersection of the two curves at the triple point of  $f$  is the projection of two intersections  $B$  and  $C$  of the corresponding curves on  $F$* . The quintic ruled surface in § 5 is an example of this.

On the other hand, it may happen that both components of the double curve pass simply through the triple point, there being a third component also passing through the triple point. When this happens the three components must be projections of three curves on  $F$  one of which passes through  $B$  and  $C$ , another through  $C$  and  $A$  and another through  $A$  and  $B$ . Then *the intersection of the two components is the projection of only one intersection of the two corresponding curves on  $F$* .

We can then state the following rule.

*Suppose that we have on a ruled surface  $f$ , in [3], two double curves of orders  $m$  and  $m'$  meeting each generator in  $k$  and  $k'$  points respectively. Then the mutual intersections of the two curves are of three possible kinds:*

*I. A simple intersection at a point which is not a triple point of the ruled surface.*

*II. An intersection at a triple point of the ruled surface, one curve having a double point and the other a simple point.*

*III. An intersection at a triple point of the ruled surface, both curves having only simple points.*

*Then, if we count each intersection under I or II as two intersections and each intersection under III as one intersection, the total number of intersections so obtained will be*

$$2 (mk' + m'k) - nkk'$$

*where  $n$  is the order of the surface.*

Both the quartic and quintic ruled surfaces in the examples in § 3 illustrate this rule.

8. An abundance of other examples could be given to illustrate the rule; we give only a few:

(A) Take three conics circumscribing the triangles of three faces of a tetrahedron. Then the lines which meet all three conics form a quintic ruled surface on which the conics are double curves; the point common to the three conics is a triple point of the ruled surface. Then, for any pair of the conics,  $m = m' = 2$ ,  $s = s' = 2$ ,  $k = k' = 1$ , while  $n = 5$ , so that

$$msk' + m's'k - nkk' = 3,$$

the intersection of the two conics at the triple point only counting once in this result, while the other intersection counts twice.

(B) Let us take an elliptic quartic curve and a line meeting it once.

Then the chords of the curve which meet the line form a ruled surface of the fifth order on which the line and the curve are both double. For the curve  $m = 4$  and  $k = 2$ , while for the line  $m' = 1$  and  $k' = 1$ . Thus, since  $n = 5$ ,  $\frac{1}{2}(mk' + m'k) - nkk' = 2$ ; the single intersection counting twice.

(C) If we take a conic  $C_2$  and a twisted cubic and place them in (1, 2) correspondence with a united point we obtain a ruled surface of the sixth order on which  $C_2$  is a double curve. The surface has also a double curve  $C_3$  of the eighth order meeting each of its generators in three points;  $C_3$  has two triple points. It has also two double points through which  $C_2$  passes and has three simple intersections with  $C_2$  (cf. §§ 175, 176).  $n = 6$ .

$$\text{For } C_3 \quad m = 8, \quad s = 2, \quad k = 3.$$

$$\text{For } C_2 \quad m' = 2, \quad s' = 2, \quad k' = 1.$$

$$msk' + m's'k - nkk' = 10 = 2 \cdot 2 + 3 \cdot 2.$$

(D) Take two twisted cubics and place them in (1, 2) correspondence with three united points. Then the joins of corresponding points give a ruled surface of the sixth order on which one of the cubics  $C_3$  is a double curve. On this surface there is also a double curve  $C_7$  meeting each generator in three points.  $C_7$  has four double points through all of which  $C_3$  passes; it has also three simple intersections with  $C_3$  (cf. §§ 175, 177).  $n = 6$ .

$$\text{For } C_7 \quad m = 7, \quad s = 2, \quad k = 3.$$

$$\text{For } C_3 \quad m' = 3, \quad s' = 2, \quad k' = 1.$$

$$msk' + m's'k - nkk' = 14 = 4 \cdot 2 + 3 \cdot 2.$$

(E) Take two conics  $C_2$  and  $D_2$  with one intersection and place them in (2, 2) correspondence with a doubly united point. Then the joins of pairs of corresponding points form a ruled surface of the sixth order on which  $C_2$  and  $D_2$  are both double curves. There is also on this surface a double curve  $C_6$  of the sixth order meeting each generator in two points. This curve has four double points, two of which are on  $C_2$  and two on  $D_2$ , while it has two simple intersections with each of  $C_2$  and  $D_2$  (cf. §§ 178, 179).  $n = 6$ .

$$\text{For } C_6 \quad m = 6, \quad s = 2, \quad k = 2.$$

$$\text{For } C_2 \quad m' = 2, \quad s' = 2, \quad k' = 1.$$

$$\text{For } D_2 \quad m'' = 2, \quad s'' = 2, \quad k'' = 1.$$

$$m's'k'' + m''s''k' - nk'k'' = 2 = 2 \cdot 1,$$

$$m''s''k + msk'' - nk''k = 8 = 2 \cdot 2 + 2 \cdot 2,$$

$$msk' + m's'k - nkk' = 8 = 2 \cdot 2 + 2 \cdot 2.$$

(F) Take two twisted cubics  $C_3$  and  $D_3$  with five intersections and place them in (2, 2) correspondence with four ordinary united points and one doubly united point. Then the joins of pairs of corresponding points generate a ruled surface of the sixth order on which  $C_3$  and  $D_3$  are double



curves. There is further on this surface a quartic double curve  $C_4$  meeting each generator in two points; this passes through the four ordinary united points of the correspondence and meets each cubic in two other points (cf. §§ 178, 180).  $n = 6$ .

For  $C_4$   $m = 4, \quad s = 2, \quad k = 2.$

For  $C_3$   $m' = 3, \quad s' = 2, \quad k' = 1.$

For  $D_3$   $m'' = 3, \quad s'' = 2, \quad k'' = 1.$

$$m's'k'' + m''s''k' - nk'k'' = 6 = 2 \cdot 1 + 4,$$

$$m''s''k + msk'' - nk''k = 8 = 2 \cdot 2 + 4,$$

$$msk' + m's'k - nkk' = 8 = 2 \cdot 2 + 4.$$

(G) Take a conic  $C_2$  and a twisted cubic  $C_3$  which has three intersections with the conic. Then place these in (2, 2) correspondence with two ordinary united points and one doubly united point. Then the joins of pairs of corresponding points form a ruled surface of the sixth order on which  $C_2$  and  $C_3$  are double curves. On this surface there is also a double curve  $C_5$  of the fifth order which meets each generator in two points and passes through the two ordinary united points of the correspondence.  $C_5$  has two double points through which  $C_3$  passes; it has two intersections with  $C_3$  other than these double points and the united points already mentioned. Also it has two intersections with  $C_2$  other than these united points (cf. § 181).  $n = 6$ .

For  $C_5$   $m = 5, \quad s = 2, \quad k = 2.$

For  $C_3$   $m' = 3, \quad s' = 2, \quad k' = 1.$

For  $C_2$   $m'' = 2, \quad s'' = 2, \quad k'' = 1.$

$$m's'k'' + m''s''k' - nk'k'' = 4 = 2 \cdot 1 + 2,$$

$$m''s''k + msk'' - nk''k = 6 = 2 \cdot 2 + 2,$$

$$msk' + m's'k - nkk' = 10 = 2 \cdot 2 + 2 \cdot 2 + 2.$$

9. At a triple point of the ruled surface there are three generators meeting; we may have part of the double curve with a triple point, or one part with a double point and another part with a simple point, or three parts all with simple points. In any case we have three distinct branches of the double curve; let  $P_1, P_2, P_3$  be points of the different branches at the triple point. Then we can regard the three generators of the surface which pass through the triple point as the lines  $P_2P_3, P_3P_1,$  and  $P_1P_2$ ; each generator joining the points on two of the branches of the curve. This statement is obscure without further elucidation; but is made much clearer when the triple point is regarded as the projection of three different points of the normal surface, at each of which the curve which projects into the

double curve has a double point. It can also be elucidated in another way—by regarding the generators as points of a curve on a quadric in [5]\*.

Suppose then that at the triple point we have two parts of the double curve, one having a double point  $P_1P_2$  and the other a simple point  $P_3$ . Then given an arbitrary line the plane joining it to  $P$  counts for two among the coincidences; for it joins the point  $P_3$  of the second curve to the point  $P_1$  of the first on the same generator  $P_1P_3$  and also joins the point  $P_3$  of the second curve to the point  $P_2$  of the first on the generator  $P_2P_3$ .

If, however, there are three parts of the double curve each passing simply through  $P$  they have there simple points  $P_1, P_2, P_3$ . If we consider intersections of the first two curves, the plane joining an arbitrary line to  $P$  only counts once among the coincidences, as joining the point  $P_1$  of the first curve to the point  $P_2$  of the second on the same generator.

But if two parts of the double curve intersect at a point which is not a triple point of the ruled surface the plane joining an arbitrary line to this intersection counts for two among the coincidences as joining the line to an intersection of the curves which lies on two different generators.

10. We know that we can represent the generators of a ruled surface  $f$  of order  $n$  in [3] by the points of a curve  $C$  of order  $n$  on a quadric fourfold  $\Omega$  in [5]. Thus we can enquire how the intersections of different parts of the double curve of  $f$  are represented on  $\Omega$ . The points of the double curve of  $f$  are in (1, 1) correspondence with the chords of  $C$  which lie on  $\Omega$ ; these chords form a ruled surface  $R_2$ .

If there is a triple point on  $f$  we shall have a plane of  $\Omega$ , representing this point, which is trisecant to  $C$ ; the points of the plane representing all the lines of [3] which pass through the triple point. Then the three points  $X, Y, Z$  in which this plane meets  $C$  represent the three generators of  $f$  which pass through the triple point; while the three lines  $YZ, ZX, XY$  which are all generators of the ruled surface formed by the chords of  $C$  lying on  $\Omega$  are all, in the (1, 1) correspondence between the generators of  $R_2$  and the points of the double curve, in correspondence with the triple point of the double curve.

If the double curve of  $f$  breaks up into component parts then  $R_2$  will have to break up into corresponding component ruled surfaces. If two parts of the double curve both pass through a triple point then the corresponding ruled surfaces will have to have generators in the same plane of  $\Omega$  which is trisecant to  $C$ ; if, however, two parts of the double curve both pass through a point which is not a triple point of the surface the corresponding ruled surfaces on  $\Omega$  must have a common generator. For the points of the double curve are in correspondence with the generators of  $R_2$ , and the same point cannot be in correspondence with two different

\* See § 10 below.

generators unless these lie in a plane which lies on  $\Omega$  and is trisecant to  $C^*$ . Thus we see that there is a sharp distinction to be drawn between the intersections of two double curves on the surface according as the intersections do or do not lie at triple points of the surface.

If, at a triple point of the ruled surface, one part of the double curve has a double point and another a simple point then, in the corresponding trisecant plane  $XYZ$  of  $C$ , two of the three chords are generators of one ruled surface, while the remaining chord is a generator of another. If, however, three different parts of the double curve of  $f$  pass through the triple point then the three chords  $YZ, ZX, XY$  of  $C$  belong to three different ruled surfaces.

Also we now obtain a new aspect of the statement made above that at a triple point  $P$  of the ruled surface where we have three points  $P_1, P_2, P_3$  on three different branches of the double curve, we regard the generators as the lines  $P_2P_3, P_3P_1, P_1P_2$ . The points  $P_1, P_2, P_3$  of the double curve are represented by the generators  $YZ, ZX, XY$  respectively of  $R_2$ , and to say that one generator is the line  $P_2P_3$  and does not (strictly) pass through  $P_1$  is the same as saying that  $X$  lies on  $ZX$  and  $XY$  but not on  $YZ$ .

11. Suppose now that we have on a ruled surface  $f$  in a space  $[r]$  a double curve and a triple curve. We shall assume that this surface is the projection of a normal ruled surface  $F$  of the same order in a space  $[R]$ ; the centre of projection being a space  $[R - r - 1]$  which does not meet  $F$ . Then there will be  $\infty^1$  trisecant planes of  $F$  which meet  $[R - r - 1]$  in lines; the spaces  $[R - r]$  containing  $[R - r - 1]$  and these planes meeting  $[r]$  in the points of the triple curve on  $f$ . Also there will be  $\infty^1$  chords of  $F$  which meet  $[R - r - 1]$ ; the spaces  $[R - r]$  containing  $[R - r - 1]$  and these chords meeting  $[r]$  in the points of the double curve on  $f$ .

If  $Q_1, Q_2, Q_3$  denote the three intersections of  $F$  with one of its trisecant planes which meets  $[R - r - 1]$  in a line, then the points  $Q$  describe a curve on  $F$  whose order is three times that of the triple curve on  $f$ , and which meets each generator of  $F$  in the same number of points as the triple curve on  $f$  meets the generators of  $f$ . Also, if  $P_1, P_2$  denote the two intersections of  $F$  with one of its chords which meets  $[R - r - 1]$  in a point then the points  $P$  describe a curve on  $F$  whose order is twice that of the double curve on  $f$  and which meets each generator of  $F$  in the same number of points as the double curve on  $f$  meets the generators of  $f$ .

Now an intersection of the double curve and the triple curve on  $f$  is joined to  $[R - r - 1]$  by an  $[R - r]$  which contains two points  $P$  and three points  $Q$  on  $F$ . Unless, then, we are to have a quadrisecant solid of  $F$  meeting  $[R - r - 1]$  in a plane, two of the points  $Q$  must coincide with the points  $P$ , and the intersection on  $f$  will be the projection of two inter-

\* We may here refer to § 66, § 92 and § 100.

sections of the corresponding curves on  $F$ . There are two possibilities in which a quadrisecant solid of  $F$  could meet  $[R - r - 1]$  in a plane; either the intersection of the two curves on  $f$  could be a multiple point on one or both of them or a third multiple curve on  $f$  might pass through this same point.

We can therefore state the following:

*If on a ruled surface  $f$  we have a double curve of order  $m$  meeting each generator in  $k$  points and a triple curve of order  $m'$  meeting each generator in  $k'$  points and if all their intersections are simple points on both curves, and if, further, none of their intersections lies on a third multiple curve of  $f$ , then the number of these intersections is*

$$\frac{1}{2} (2mk' + 3m'k - nkk'),$$

where  $n$  is the order of the surface. In obtaining this we have assumed that the multiple curves are projections of simple curves on a normal surface.

And similarly we have the following:

*If on a ruled surface  $f$  we have a curve of order  $m$  and multiplicity  $s$  meeting each generator in  $k$  points and also a curve of order  $m'$  and multiplicity  $s'$  meeting each generator in  $k'$  points, and if all their intersections are simple points on both curves, and if, further, none of their intersections lies on a third multiple curve of  $f$ , then the number of these intersections is*

$$\frac{1}{s_0} (msk' + m's'k - nkk'),$$

where  $n$  is the order of the surface and  $s_0$  is the smaller of  $s$  and  $s'$ . The result holds if  $s = s' = s_0$ .

12. We can illustrate these last statements by examples:

(A) If we take a line  $R$  and a conic and place them in (1, 3) correspondence the joins of pairs of corresponding points will generate a ruled surface of the fifth order on which  $R$  is a triple line. This surface has also a double curve; this is a twisted cubic  $C_3$  which has  $R$  for a chord. A plane through  $R$  meets the conic in two points and contains two generators of the surface; their intersection is a point of  $C_3$  (cf. §§ 139, 140).  $n = 5$ .

$$\text{For } C_3 \quad m = 3, \quad s = 2, \quad k = 1.$$

$$\text{For } R \quad m' = 1, \quad s' = 3, \quad k' = 1.$$

$$msk' + m's'k - nkk' = 4 = 2 \cdot 2.$$

The ruled surface is rational, the plane sections being quintic curves each with a triple point and three double points. It can therefore be obtained by projection from a rational normal quintic ruled surface  $F$  in [6]. We project from a plane  $\omega$ , and for this particular surface we choose  $\omega$  to meet a solid  $K$  containing a directrix cubic  $\Delta$  of the normal surface

in a line  $p^*$ . Then the pencil of planes through  $p$  in  $K$  gives a system of trisecant planes of  $F$  which all meet the centre of projection in lines (in this case the same line). The plane  $\omega$  meets  $M_5^3$ , the locus of the chords of  $F$ , in  $p$  and a conic  $\mathfrak{D}$ . Through each point of  $\mathfrak{D}$  passes a chord of  $F$ , and these chords will meet  $F$  in the points of a rational sextic curve  $C_6$  meeting each generator of  $F$  once. Then  $C_6$  and  $\Delta$  have four intersections, these being the points of  $F$  on the two chords which pass through the intersections of  $p$  and  $\mathfrak{D}$ .

Each intersection of  $R$  and  $C_3$  is the projection of two intersections of  $\Delta$  and  $C_6$ .

The conic which we put in correspondence with  $R$  is a simple conic on the surface; it does not meet  $R$  but it meets  $C_3$  three times.

For the conic  $m'' = 2, \quad s'' = 1, \quad k'' = 1,$   
 so that  $m''s''k'' + m's'k'' - nk'k'' = 0,$   
 $m''s''k + msk'' - nkk'' = 3.$

The three intersections of the conic with  $C_3$  are the projections of three distinct intersections of  $C_6$  and the directrix conic of the normal surface.

(B) If we take a line  $R$  and a twisted cubic in (1, 3) correspondence the joins of pairs of corresponding points form a ruled surface of the sixth order on which  $R$  is a triple line. This surface has also a double curve  $C_7$  of the seventh order which meets every generator in two points and  $R$  in four points (cf. § 222).  $n = 6$ .

For  $C_7$   $m = 7, \quad s = 2, \quad k = 2.$   
 For  $R$   $m' = 1, \quad s' = 3, \quad k' = 1.$   
 $msk' + m's'k - nkk' = 8 = 2 \cdot 4.$

The ruled surface is rational and is obtained by projection from a normal surface in [7]. The normal surface has  $\infty^1$  directrix cubic curves on it; we project from a solid  $S$  meeting the solid  $K$  containing one of these cubic curves in a line  $p$ . The chords of  $F$  meeting  $S$  do so in  $p$  and a rational quintic  $\mathfrak{D}_5^0$  of which  $p$  is the quadrisecant. The chords of  $F$  through the intersections of  $p$  and  $\mathfrak{D}_5^0$  give four pairs of points on  $F$ ; each of these pairs of points projects into one intersection of  $R$  and  $C_7$ .

(C) Finally, we will give an example in which the conditions required do not hold, there being three different multiple curves on a ruled surface with points common to all of them.

Let us take a rational quintic curve  $C$  in [4]; this has a trisecant chord  $t$  meeting it in three points  $P, Q, R$ . Suppose that we have a quadric  $\Omega$  containing  $C$ ; this quadric will also contain  $t$ .

\* There are  $\infty^2$  curves  $\Delta$  on  $F$  with corresponding solids  $K$ . These solids  $K$  generate the  $M_5^3$  formed by the chords of  $F$ , and a general plane of [6] will not meet any of them in a line.

Then the chords of  $C$  which lie on  $\Omega$  form a ruled surface of order 12 on which  $C$  and  $t$  are both triple curves. There is also on this surface a double curve of the sixth order meeting each generator in one point and passing through  $P$ ,  $Q$  and  $R$ \*.  $n = 12$ .

For  $C$   $m = 5, \quad s = 3, \quad k = 2.$

For the double curve

$$m' = 6, \quad s' = 2, \quad k' = 1.$$

$$msk' + m's'k - nkk' = 15.$$

The number of intersections of  $C$  with the sextic curve other than  $P$ ,  $Q$ ,  $R$  is known to be six, and each of these will be counted twice according to our rule. Hence *the three intersections which are on  $t$  are only counted once.*

We have here an example in which a triple curve and a double curve intersect, three of their intersections being on another triple curve of the ruled surface. It appears that these last three intersections of the triple curve and the double curve only give rise to one coincidence each.

The ruled surface can be shown to be elliptic; it can therefore be obtained by projection from an elliptic ruled surface of order 12 in [11]; the centre of projection is a [6] not meeting the surface.

\* See § 96.









