

A GEOMETRIC ANALYSIS OF THE LAGERSTROM MODEL PROBLEM

NIKOLA POPOVIĆ AND PETER SZMOLYAN

ABSTRACT. Lagerstrom’s model problem is a classical singular perturbation problem which was introduced to illustrate the ideas and subtleties involved in the analysis of viscous flow past a solid at low Reynolds number by the method of matched asymptotic expansions. In this paper, the corresponding boundary value problem is analyzed geometrically by using methods from the theory of dynamical systems, in particular invariant manifold theory. As an essential part of the dynamics takes place near a line of non-hyperbolic equilibria, a blow-up transformation is introduced to resolve these singularities. This approach leads to a constructive proof of existence and local uniqueness of solutions and to a better understanding of the singular perturbation nature of the problem. In particular, the source of the logarithmic switchback phenomenon is identified.

1. INTRODUCTION

Viscous flow past a solid at low Reynolds number is a classical singular perturbation problem from fluid dynamics. Steady low Reynolds number flow of an incompressible fluid past a circular cylinder was studied by [Sto51]: as a first approximation, he took the Reynolds number to be zero in the governing equations and found that the resulting boundary value problem has no solution (*Stokes paradox*). For flow past a sphere, Stokes did in fact find an approximation which has been widely used. In an attempt to derive a higher-order approximation for the spherical case, however, [Whi89] found that the next term has a singularity at infinity (*Whitehead paradox*). More than half a century later, [Ose10] observed that these seeming paradoxes were due to an incorrect treatment of the flow far from the cylinder respectively the sphere and that they could be avoided by linearizing about the flow at infinity. Oseen solved the resulting equation for spherical flow and obtained a solution which improves Stokes’ solution; however, he failed to give a systematic expansion procedure. The conceptual structure of the problem was clarified much later by Kaplun and Lagerstrom [Kap57, KL57] and Proudman and Pearson [PP57], who showed that it could be solved by the systematic use of the method of *matched asymptotic expansions*.

Later still, Lagerstrom proposed his model problem to illustrate the mathematical ideas and techniques used by Kaplun in the asymptotic treatment of low Reynolds number flow, see [Kap57, Lag66, KL57]. In its simplest formulation, the model is given by the nonlinear, non-autonomous second-order boundary value problem

$$(1) \quad \ddot{u} + \frac{n-1}{x} \dot{u} + u\dot{u} = 0,$$

1991 *Mathematics Subject Classification.* 34E15, 37N10, 76D05.

Key words and phrases. Singular perturbations; Invariant manifolds; Blow-up.

with boundary conditions

$$(2) \quad u(\varepsilon) = 0, \quad u(\infty) = 1.$$

Here, $n \in \mathbb{N}$, $0 < \varepsilon \leq x \leq \infty$, and the overdot denotes differentiation with respect to x . Heuristically speaking, (1),(2) models slow incompressible viscous flow in n dimensions, scaled in such a way that the dependence of u on ε (the analogue of the Reynolds number) occurs through the inner boundary condition. Here, $u(\varepsilon) = 0$ corresponds to the no-slip condition at the surface of an “ n -sphere” of diameter ε , whereas $u(\infty) = 1$ requires the “flow” to be uniform far away from the “ n -sphere”. We focus on the physically relevant cases $n = 2$ and $n = 3$ corresponding to flow around a cylinder and a sphere, respectively. This problem, which is not a model in the physical sense, but a mathematical model equation, was analyzed by Lagerstrom himself and many others, see e.g. [Bus71, CFL78, HTB90, KW96, KWK95, Lag88, RS75, Ski81], and the references therein. While it displays similar qualitative properties as the original fluid dynamical problem, Lagerstrom’s model is analytically much simpler, owing largely to the fact that it is an ordinary rather than a partial differential equation. Both the original problem of viscous flow past a solid at low Reynolds number and Lagerstrom’s model example have been quite influential for the development of singular perturbation theory in general and of the method of matched asymptotic expansions in particular.

More recently, an alternative approach to singularly perturbed problems known as *geometric singular perturbation theory* has been developed. This approach is based on methods from the theory of dynamical systems, in particular on invariant manifold theory. In this context, outer solutions and their expansions find a geometric explanation in terms of slow center-like manifolds which depend smoothly on the singular perturbation parameter. Standard exponential layer (inner) solutions are explained geometrically as invariant foliations of stable or unstable manifolds of slow center-like manifolds, which again depend smoothly on the singular perturbation parameter, see [Fen79] or [Jon95].

However, this well-developed geometric theory does not apply at points where normal hyperbolicity is lost, i.e., at points where the slow manifold ceases to be exponentially attractive respectively repelling.

Recently, the geometric approach has been extended to the case where normal hyperbolicity fails due to a single zero eigenvalue, a situation which arises frequently in applications, e.g. in relaxation oscillations. This advance has been possible due to the use of the blow-up method [DR91, Dum93, DR96]. Blow-up can be described as a sophisticated rescaling which allows one to identify the dominant scales in various regions near a singularity. In particular, the blow-up method has been applied in a detailed analysis of the simple fold problem, see [KS01, vGKS]. In these works, slow manifolds are continued beyond the fold point. Additionally, the complicated structure of the corresponding asymptotic expansions is explained and an algorithm to compute them is given.

The aim of the present work and its sequel [PS] is to analyze Lagerstrom’s model problem in a similar spirit.

The paper is organized as follows. In Section 2, a brief introduction to Lagerstrom’s model equation and its analysis by means of matched asymptotic expansions is given. Section 3 contains a dynamical systems reformulation of the Lagerstrom

model. The governing equations are rewritten as an autonomous system of ordinary differential equations. A solution of the original boundary value problem is seen to correspond to an orbit connecting an (unknown) point in a one-dimensional manifold \mathcal{V}_ε representing the boundary condition at $x = \varepsilon$ to a degenerate equilibrium Q which corresponds to the boundary condition at $x = \infty$. In the limit of $\varepsilon = 0$, Q becomes even more degenerate, which – at least partially – explains the singular perturbation nature of the problem. A singular orbit Γ is identified which connects the manifold \mathcal{V}_0 to the equilibrium Q for $\varepsilon = 0$. To resolve this singular behavior, a blow-up transformation is introduced. In Section 4, the dynamics of the blown-up problem is analyzed in detail. In the blown-up system, existence and uniqueness of solutions for Lagerstrom’s model is proven by carefully tracking the manifold \mathcal{V}_ε of boundary values along the singular orbit Γ to show that it intersects transversely the stable manifold of Q . In most parts of the analysis, one has to distinguish between the cases $n = 3$ and $n = 2$, the latter being difficult due to its more degenerate nature.

2. LAGERSTROM’S MODEL EQUATION

By introducing the inner (stretched) variable

$$(3) \quad \xi = \frac{x}{\varepsilon}$$

in (1),(2), we obtain the equivalent formulation

$$(4) \quad u'' + \frac{n-1}{\xi}u' + \varepsilon uu' = 0$$

of Lagerstrom’s model equation, with $1 \leq \xi \leq \infty$ and boundary conditions

$$(5) \quad u(1) = 0, \quad u(\infty) = 1;$$

here, the prime denotes differentiation with respect to ξ .

Lagerstrom’s model equation is a classical example of a singularly perturbed problem, as the solution obtained by setting $\varepsilon = 0$ in (4) is not a uniformly valid approximation to the solution of (4) for $1 \leq \xi \leq \infty$. Moreover, it shows that having the small parameter ε multiply the highest derivative in a differential equation is not a necessary condition for a problem to be singular. In what is to come, we will restrict ourselves to the cases $n = 2$ and $n = 3$, which correspond to the physically relevant settings of flow in two and three dimensions, respectively.

We briefly discuss Lagerstrom’s analysis of (1),(2) based on the method of matched asymptotic expansions. Let us first consider $n = 3$: by assuming a (regular) perturbation expansion for u of the form

$$(6) \quad u(\xi, \varepsilon) = u_0(\xi) + \varepsilon u_1(\xi) + \dots,$$

one obtains

$$(7a) \quad u_0'' + \frac{2}{\xi}u_0' = 0,$$

$$(7b) \quad u_1'' + \frac{2}{\xi}u_1' = -u_0 u_0'$$

for ξ fixed as $\varepsilon \rightarrow 0$; in analogy with the fluid dynamical problem, (7a) is called the *Stokes equation*.

With $u_0 = 0$ for $\xi = 1$ and $u_0 \rightarrow 1$ for $\xi \rightarrow \infty$, the leading term of the inner approximation (6) is

$$(8) \quad u_0 = 1 - \frac{1}{\xi};$$

the solution of (7b) which satisfies $u_1 = 0$ at $\xi = 1$ is given by

$$(9) \quad u_1 = - \left(1 + \frac{1}{\xi}\right) \ln \xi + \alpha \left(1 - \frac{1}{\xi}\right).$$

However, no choice of the constant α can prevent u_1 from being logarithmically infinite for $\xi \rightarrow \infty$. This is the analogue of the fluid dynamical *Whitehead paradox*. Thus, the naive expansion (6) is not uniformly valid for ξ large; this forces one to apply the rescaling in (3).

For $\xi = \mathcal{O}(\varepsilon^{-1})$, (6) and (9) imply $u = 1 + \mathcal{O}(\varepsilon \ln \varepsilon)$, which suggests to replace (6) by an expansion of the form

$$(10) \quad u(\xi, \varepsilon) = 1 - \frac{1}{\xi} + \varepsilon \ln \varepsilon \tilde{u}_1(\xi) + \varepsilon u_1(\xi) + \dots$$

for ξ fixed as $\varepsilon \rightarrow 0$.

To approximate solutions of (1) for $x = \varepsilon \xi$ fixed as $\varepsilon \rightarrow 0$, one uses the outer expansion

$$(11) \quad u(x, \varepsilon) = U_0(x) + \varepsilon \ln \varepsilon \tilde{U}_1(x) + \varepsilon U_1(x) + \dots$$

which is akin to the Oseen expansion in the fluid dynamical problem. The leading order term satisfying (1) is found to be $U_0 = 1$, while \tilde{U}_1 has to satisfy the homogeneous Oseen equation

$$(12) \quad \frac{d^2 \tilde{U}_1}{dx^2} + \left(\frac{2}{x} + 1\right) \frac{d\tilde{U}_1}{dx} = 0,$$

with $\tilde{U}_1(\infty) = 0$; the same is true of U_1 . Equation (12) is linear; its solution can be given in terms of certain exponential integrals, with the constants left to be determined by matching.

For $n = 2$, the situation is even more involved: the same intuitive reasoning as before yields

$$(13) \quad u_0 = \alpha \ln \xi$$

for the leading term of the inner approximation. Obviously, the condition at infinity cannot be satisfied with any choice of α (*Stokes paradox*). Nevertheless, the rescaling in (3) is applicable again, which implies that the troublesome condition at infinity is in the region of x fixed as $\varepsilon \rightarrow 0$. As u must be $\mathcal{O}(1)$ there, (6) and (13) imply that $\alpha = \mathcal{O}((\ln \varepsilon)^{-1})$, which suggests asymptotic expansions

$$(14) \quad u(\xi, \varepsilon) = -\frac{1}{\ln \varepsilon} u_0(\xi) + \frac{1}{(\ln \varepsilon)^2} u_1(\xi) + \dots$$

and

$$(15) \quad u(x, \varepsilon) = 1 - \frac{1}{\ln \varepsilon} U_1(x) + \frac{1}{(\ln \varepsilon)^2} U_2(x) + \dots,$$

respectively. As for $n = 3$, matching these expansions is still possible, although the overlap domain is now very small, see [LC72]. This is in essence Kaplun's resolution

of the Stokes paradox: the Stokes solution is an inner solution which must satisfy a matching condition, but not necessarily the boundary condition at infinity.

Rigorous results for Lagerstrom's model equation have been obtained by numerous researchers using a variety of methods. Existence and uniqueness of solutions was shown in [RS75] by transforming (4) into a pair of integral equations and by applying a contraction mapping theorem. Hsiao [Hsi73] gave a rigorous discussion of existence for $n = 2$ and $\varepsilon \rightarrow 0$, whereas Cole [Col68] utilized an invariance group of (1) to obtain a similar result. In [CFL78], a related initial value problem was transformed into an integral equation, which was then shown to have a unique solution by constructing suitable super- and subsolutions. Hunter et al. [HTB90] proved that the so-called Oseen iteration, an iterative scheme based on using the outer approximation throughout, converges to a unique solution for all ε .

Remark 1 (Logarithmic switchback). The introduction of an $\varepsilon \ln \varepsilon$ -term in (6) is unexpected, as it is not directly forced by the equation, but by the matching. Perturbation problems in which the small parameter ε (but not $\ln \varepsilon$) occurs in the formulation of the problem, whereas $\ln \varepsilon$ occurs in the asymptotic expansion, have been encountered conspicuously often in the resolution of paradoxes in problems of fluid dynamics. The phenomenon is known as *logarithmic switchback*, see [Lag88] for details. \square

Remark 2. A generalization of (1) to arbitrary integral (and even real) dimensions is feasible and has indeed been undertaken by several researchers, see e.g. [LR84]. Our approach applies for any $n \in \mathbb{R}$, $n \geq 2$, as well, with only a few minor changes required.

Notably, the form of the simpler inner expansion (6) depends even more critically on the value of n than the outer expansion. The larger n is, the further the occurrence of switchback terms is postponed; Stokes' paradox is thus only delayed, as it will always occur sooner or later. In particular, there is no switchback for n irrational. \square

3. A DYNAMICAL SYSTEMS APPROACH

3.1. Our strategy. We will employ a shooting argument to prove existence and (local) uniqueness of solutions to the boundary value problem (1),(2). To that end, we rewrite Lagerstrom's model problem as an equivalent autonomous first-order dynamical system. As is usual in geometric singular perturbation theory, the starting point of the analysis are the equations on the fast (inner) scale, i.e., (4). We replace $\xi \in [1, \infty)$ by $\eta := \xi^{-1} \in (0, 1]$; $\xi' = 1$ implies the equation $\eta' = -\eta^2$. By setting $u' = v$, we obtain the system

$$(16) \quad \begin{aligned} u' &= v, \\ v' &= -(n-1)\eta v - \varepsilon uv, \\ \eta' &= -\eta^2, \end{aligned}$$

with boundary conditions

$$(17) \quad u(1) = 0, \quad \eta(1) = 1, \quad u(\infty) = 1;$$

note that (17) in fact entails $\eta(\infty) = 0$ and $v(\infty) = 0$ for the solution to (16), whereas $v(1)$ remains yet to be determined.




FIGURE 2. Geometry of (16) for $\varepsilon = 0$ and (a) $n = 3$, respectively, (b) $n = 2$.

Proposition 3.2. *Let $k \in \mathbb{N}$ be arbitrary, and let $\varepsilon > 0$.*

- (1) *There exists an attracting two-dimensional center manifold $\mathcal{W}_\varepsilon^c$ for (16) which is given by $\{v = 0\}$.*
- (2) *For $|u - 1|$, v , and η sufficiently small, there is a stable invariant \mathcal{C}^k -smooth foliation $\mathcal{F}_\varepsilon^s$, with base $\mathcal{W}_\varepsilon^c$ and one-dimensional \mathcal{C}^k -smooth fibers.*

Proof. The first assertion is obvious from Lemma 3.1 and the fact that $\{v = 0\}$ is an invariant subspace for (16); the second assertion follows from standard invariant manifold theory (see e.g. [Fen79] or [CLW94]). \square

Given $\varepsilon > 0$ fixed, one can thus define the stable manifold $\mathcal{W}_\varepsilon^s$ of Q as

$$(22) \quad \mathcal{W}_\varepsilon^s := \bigcup_{P \in \Upsilon} F_\varepsilon^s(P),$$

where $\Upsilon := \{(1, 0, \eta) \mid 0 \leq \eta \ll 1\}$, i.e., as a union of fibers $F_\varepsilon^s \in \mathcal{F}_\varepsilon^s$ with base points in the weakly stable orbit Υ , see Figure 1. Of particular importance is the fiber $F_\varepsilon^s(Q)$ with base point Q ; note that by Lemma 3.1, $F_\varepsilon^s(Q)$ is tangent to $(1, -\varepsilon, 0)^T$ at Q .

Remark 3. In fact, due to the simple structure of (16) for $\eta = 0$, $F_\varepsilon^s(Q)$ can be found explicitly by writing e.g.

$$(23) \quad \frac{dv}{du} = -\varepsilon u$$

and solving for v to obtain

$$(24) \quad v(u) = \frac{\varepsilon}{2} (1 - u^2);$$

here, we have used $v(1) = 0$. \square

We will in the following write $\mathcal{W}_\varepsilon^{ss}$ instead of $F_\varepsilon^s(Q)$ to stress that $F_\varepsilon^s(Q)$ is the one-dimensional strongly stable manifold of Q .

A solution of the boundary value problem (16),(17) corresponds to a forward orbit starting in \mathcal{V}_ε and converging to Q as $\xi \rightarrow \infty$. Hence, existence and uniqueness of solutions will follow by showing that the saturation of \mathcal{V}_ε under the flow defined by (16), which we call $\mathcal{M}_\varepsilon := \mathcal{V}_\varepsilon \cdot [1, \infty)$, intersects $\mathcal{W}_\varepsilon^s$ in a unique orbit; here, the dot denotes the application of the flow induced by (16).

For $\varepsilon = 0$, it is straightforward to obtain *singular orbits* connecting \mathcal{V}_0 to Q . It is these orbits we will use as templates for orbits of the full problem ($\varepsilon > 0$). The case $n = 3$ is the simpler one, as the forward orbit

$$(25) \quad \gamma := \{(1 - \eta, \eta^2, \eta) \mid \eta \in (0, 1]\}$$

through $P := (0, 1, 1)$ obtained by solving (16) for $\varepsilon = 0$ is asymptotic to Q . We thus define the singular orbit Γ by

$$(26) \quad \Gamma := \gamma \cup \{Q\},$$

see Figure 2. For $n = 2$, the situation is more involved: recall that there is no solution to (16),(17) for $\varepsilon = 0$ in that case. However, a singular orbit Γ can still be defined: let $P := (0, 0, 1)$, and let γ denote the orbit

$$(27) \quad \gamma := \{(0, 0, \eta) \mid \eta \in (0, 1]\}$$

through P , which is forward asymptotic to the origin O . Then,


$$(28) \quad \Gamma := \gamma \cup \{O\} \cup \{(u, 0, 0) \mid u \in (0, 1)\} \cup \{Q\}.$$

For $n = 2$, Γ thus contains a segment of the line of equilibria ℓ_0 , which accounts for the complicated nature of the problem.

We now proceed as follows to prove existence and uniqueness of solutions to (16): we track \mathcal{M}_ε through phase space and show that it intersects transversely the stable manifold $\mathcal{W}_\varepsilon^s$ of Q (see again Figure 1). Due to the fact that we are only interested in ε small, we are going to take a perturbational approach, i.e., we intend to set $\varepsilon = 0$ in (16), and to track \mathcal{M}_0 along Γ under the resulting flow. For $\varepsilon = 0$, however, the equations in (16) are even more degenerate than they are for $\varepsilon > 0$, see Lemma 3.1. Due to the non-hyperbolic character of the problem for $\varepsilon = 0$, there is no stable foliation \mathcal{F}_0^s ; hence, a stable manifold \mathcal{W}_0^s does not exist, either. We therefore have to modify our approach. To that end, we extend (16) by appending the (trivial) equation $\varepsilon' = 0$, obtaining

$$(29) \quad \begin{aligned} u' &= v, \\ v' &= -(n-1)\eta v - \varepsilon uv, \\ \eta' &= -\eta^2, \\ \varepsilon' &= 0 \end{aligned}$$

in extended phase space, where the boundary conditions are still given by (17). Contrary to the above, the parameter ε is not fixed now, but is allowed to vary in an interval $[0, \varepsilon_0]$, with $\varepsilon_0 > 0$ small. Correspondingly, for the extended system (29), we define the manifolds \mathcal{V} and \mathcal{M} by $\mathcal{V} := \bigcup_{\varepsilon \in [0, \varepsilon_0]} \mathcal{V}_\varepsilon \times \{\varepsilon\}$ and $\mathcal{M} := \bigcup_{\varepsilon \in [0, \varepsilon_0]} \mathcal{M}_\varepsilon \times \{\varepsilon\}$, respectively. We will see that by using blow-up, we will be able to define stable manifolds \mathcal{W}^{ss} and \mathcal{W}^s in a smooth way down to $\varepsilon = 0$.

FIGURE 3. The blow-up transformation Φ .

Remark 4. Though Lagerstrom's model equation is a singular perturbation problem, it is not strictly so in the sense of [Fen79]. Indeed, the dynamics of (29) is to be characterized as center-like rather than slow-fast. \square

3.2. The blow-up transformation. To analyze the dynamics of (29) near the line $\ell := \{(u, 0, 0, 0) \mid u \in \mathbb{R}^+\}$ of equilibria² of (29), we introduce a (*polar*) *blow-up transformation*

$$(30) \quad \Phi : \begin{cases} \mathbb{R} \times B \rightarrow \mathbb{R}^4, \\ (\bar{u}, \bar{v}, \bar{\eta}, \bar{\varepsilon}, \bar{r}) \mapsto (\bar{u}, \bar{r}\bar{v}, \bar{r}\bar{\eta}, \bar{r}\bar{\varepsilon}), \end{cases}$$

with $B := \mathbb{S}^2 \times \mathbb{R}$. Here, \mathbb{S}^2 denotes the two-sphere in \mathbb{R}^3 , i.e., $\mathbb{S}^2 = \{(\bar{v}, \bar{\eta}, \bar{\varepsilon}) \mid \bar{v}^2 + \bar{\eta}^2 + \bar{\varepsilon}^2 = 1\}$. Note that obviously $\Phi^{-1}(\ell) = \mathbb{R} \times \mathbb{S}^2 \times \{0\}$, which is the blown-up locus obtained by setting $\bar{r} = 0$. Moreover, for $\bar{r} \neq 0$, i.e., away from $\Phi^{-1}(\ell)$, Φ is a C^∞ -diffeomorphism. We will only be interested in $\bar{r} \in [0, r_0]$, with $r_0 > 0$ small.

The reason for introducing (30) is that degenerate equilibria, such as those in ℓ , can in many cases be neatly analyzed by means of blow-up techniques, see [Dum93]. The blow-up is a (singular) coordinate transformation whereby the degenerate equilibrium is blown up to some n -sphere. Transverse to the sphere and even on the sphere itself one often gains enough hyperbolicity to allow for a complete analysis by standard techniques. For planar vector fields, the method is widely known, see e.g. [GH83]; not unexpectedly, however, difficulties mount with rising dimension. A general discussion of blow-up can be found in [DR91]. The analysis of non-hyperbolic points in singular perturbation problems was initiated by Dumortier and Roussarie, see [Dum93, DR96], and was further developed in [KS01, vGKS]. We refer to these works for an introduction and more background material.

²We will restrict ourselves to $u \geq 0$ in the following, due to the boundary conditions imposed.

The vector field on $\mathbb{R} \times B$, which is induced by the vector field corresponding to (29), is best studied by introducing different charts for the manifold $\mathbb{R} \times B$. In what is to come, it suffices to consider two charts K_1 and K_2 corresponding to $\bar{\eta} > 0$ and $\bar{\varepsilon} > 0$ in (30), respectively, see Figure 3. The coordinates $(u_1, v_1, r_1, \varepsilon_1)$ in K_1 are given by

$$(31) \quad u_1 = \bar{u}, \quad v_1 = \bar{v}\bar{\eta}^{-1}, \quad r_1 = \bar{r}\bar{\eta}, \quad \varepsilon_1 = \bar{\varepsilon}\bar{\eta}^{-1}$$

for $\bar{\eta} > 0$. Similarly, for (u_2, v_2, η_2, r_2) in K_2 , one obtains

$$(32) \quad u_2 = \bar{u}, \quad v_2 = \bar{v}\bar{\varepsilon}^{-1}, \quad \eta_2 = \bar{\eta}\bar{\varepsilon}^{-1}, \quad r_2 = \bar{r}\bar{\varepsilon},$$

where $\bar{\varepsilon} > 0$. We will see that these two charts correspond precisely to the inner and outer regions in the method of matched asymptotic expansions.

Remark 5 (Notation). Let us introduce the following notation: for any object \square in the original setting, let $\bar{\square}$ denote the corresponding object in the blow-up; in charts K_i , $i = 1, 2$, the same object will appear as \square_i when necessary. \square

In K_1 , the blow-up transformation (30) reads

$$(33) \quad \Phi_1 : \begin{cases} \mathbb{R}^4 \rightarrow \mathbb{R}^4, \\ (u_1, v_1, r_1, \varepsilon_1) \mapsto (u_1, r_1 v_1, r_1, r_1 \varepsilon_1), \end{cases}$$

which is a *directional blow-up* in the direction of positive η . With

$$(34) \quad u = u_1, \quad v = r_1 v_1, \quad \eta = r_1, \quad \varepsilon = r_1 \varepsilon_1,$$

the blown-up vector field in K_1 is then given by

$$(35) \quad \begin{aligned} u'_1 &= r_1 v_1, \\ v'_1 &= (2-n)r_1 v_1 - r_1 \varepsilon_1 u_1 v_1, \\ r'_1 &= -r_1^2, \\ \varepsilon'_1 &= r_1 \varepsilon_1, \end{aligned}$$

which can be desingularized by setting $\frac{d}{d\bar{\varepsilon}} = r_1 \frac{d}{d\varepsilon_1}$ in (35) and by dividing out the common factor r_1 on both sides of the equations:

$$(36) \quad \begin{aligned} u'_1 &= v_1, \\ v'_1 &= (2-n)v_1 - \varepsilon_1 u_1 v_1, \\ r'_1 &= -r_1, \\ \varepsilon'_1 &= \varepsilon_1. \end{aligned}$$

This desingularization is necessary to obtain a non-trivial flow for $r_1 = 0$; it corresponds to a rescaling of time, leaving the phase portrait unchanged.

The equilibria of (36) are easily seen to lie in $\ell_1 := \{(u_1, 0, 0, 0) \mid u_1 \in \mathbb{R}^+\}$; the linearization there is

$$(37) \quad \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 2-n & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Depending on n , two cases have to be considered:

Lemma 3.3. *For $n = 3$, -1 is an eigenvalue of multiplicity two, whereas 0 and 1 are simple eigenvalues of (37). The corresponding eigenspaces are*

$$(38) \quad \text{span} \{(0, 0, 1, 0)^T, (1, -1, 0, 0)^T\}, \quad \text{span} \{(1, 0, 0, 0)^T\}, \quad \text{span} \{(0, 0, 0, 1)^T\}.$$

For $n = 2$, the multiplicity of 0 is two, with -1 and 1 simple and the eigenspaces given by

$$(39) \quad \text{span} \{(1, 0, 0, 0)^T, (0, 1, 0, 0)^T\}, \quad \text{span} \{(0, 0, 1, 0)^T\}, \quad \text{span} \{(0, 0, 0, 1)^T\};$$

here, $(0, 1, 0, 0)^T$ is a generalized eigenvector of the eigenvalue 0 .

Proof. Computation. □

In chart K_2 , the blow-up transformation (30) is given by

$$(40) \quad \Phi_2 : \begin{cases} \mathbb{R}^4 \rightarrow \mathbb{R}^4, \\ (u_2, v_2, \eta_2, r_2) \mapsto (u_2, r_2 v_2, r_2 \eta_2, r_2); \end{cases}$$

it follows that

$$(41) \quad u = u_2, \quad v = r_2 v_2, \quad \eta = r_2 \eta_2, \quad \varepsilon = r_2,$$

which is simply an ε -dependent rescaling of the original variables, since $r_2 = \varepsilon$. Given (41), we obtain

$$(42) \quad \begin{aligned} u'_2 &= r_2 v_2, \\ v'_2 &= (1 - n)r_2 \eta_2 v_2 - r_2 u_2 v_2, \\ \eta'_2 &= -r_2 \eta_2^2, \\ r'_2 &= 0 \end{aligned}$$

for the blown-up vector field in K_2 . Desingularizing (dividing by r_2) once again yields

$$(43) \quad \begin{aligned} u'_2 &= v_2, \\ v'_2 &= (1 - n)\eta_2 v_2 - u_2 v_2, \\ \eta'_2 &= -\eta_2^2, \\ r'_2 &= 0; \end{aligned}$$

these equations are simple insofar as r_2 occurs only as a parameter. For $r_2 \in [0, r_0]$, the equilibria of (43) are given by $\ell_2 := \{(u_2, 0, 0, r_2) \mid u_2 \in \mathbb{R}^+\}$, with corresponding linearization

$$(44) \quad \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & -u_2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Lemma 3.4. *The eigenvalues of (44) are 0 and $-u_2$, where the multiplicity of 0 is three. The corresponding eigenspaces are*

$$(45) \quad \text{span} \{(1, 0, 0, 0)^T, (0, 0, 1, 0)^T, (0, 0, 0, 1)^T\}, \quad \text{span} \{(1, -u_2, 0, 0)^T\}.$$

Proof. Computation. □

Note that ℓ_2 corresponds exactly to the original line ℓ_ε , i.e., the point Q we are ultimately interested in is retrieved in chart K_2 after the blow-up. We have the following result for the change of coordinates between charts K_1 and K_2 on their overlap domain:

Lemma 3.5. *Let κ_{12} denote the change of coordinates from K_1 to K_2 , and let $\kappa_{21} = \kappa_{12}^{-1}$ be its inverse. Then, κ_{12} is given by*

$$(46) \quad u_2 = u_1, \quad v_2 = v_1 \varepsilon_1^{-1}, \quad \eta_2 = \varepsilon_1^{-1}, \quad r_2 = r_1 \varepsilon_1,$$

and κ_{21} is given by

$$(47) \quad u_1 = u_2, \quad v_1 = v_2 \eta_2^{-1}, \quad r_1 = r_2 \eta_2, \quad \varepsilon_1 = \eta_2^{-1}.$$

Proof. Computation. □

For the following analysis, it is convenient to define *sections* Σ_1^{in} , Σ_1^{out} , and Σ_2^{in} in K_1 and K_2 , respectively, where

$$(48a) \quad \Sigma_1^{in} := \{(u_1, v_1, r_1, \varepsilon_1) \mid u_1 \geq 0, v_1 \geq 0, \varepsilon_1 \geq 0, r_1 = \rho\},$$

$$(48b) \quad \Sigma_1^{out} := \{(u_1, v_1, r_1, \varepsilon_1) \mid u_1 \geq 0, v_1 \geq 0, r_1 \geq 0, \varepsilon_1 = \delta\},$$

$$(48c) \quad \Sigma_2^{in} := \{(u_2, v_2, \eta_2, r_2) \mid u_2 \geq 0, v_2 \geq 0, r_2 \geq 0, \eta_2 = \delta^{-1}\},$$

with $0 < \rho, \delta \ll 1$ arbitrary, but fixed; see Figure 4. Note that $\kappa_{12}(\Sigma_1^{out}) = \Sigma_2^{in}$.

The shooting argument outlined in Section 3.1 is now carried out in the blown-up system, or, to be precise, in charts K_1 and K_2 . The sole reason for introducing (30) and considering (36) and (43) instead of (29), however, is that we have gained enough hyperbolicity to extend the argument down to and including $\varepsilon = 0$, i.e., to define the stable manifold $\overline{\mathcal{W}}^s$ of $\overline{Q} \in \overline{\ell}$ for $\overline{r} = 0$. This follows from chart K_2 , as the linearization of (43) at Q_2 has a negative eigenvalue irrespective of the value of r_2 , in contrast to the linearization of the original system (29) at Q . We will thus be able to track $\overline{\mathcal{V}}$ along the singular orbit $\overline{\Gamma}$, and to show that the resulting manifold $\overline{\mathcal{M}}$ intersects $\overline{\mathcal{W}}^s$ transversely. This intersection will give the sought-after family of solutions to the boundary value problem (16),(17) for $\varepsilon \in (0, \varepsilon_0]$. The situation is illustrated in Figure 5.

4. EXISTENCE AND UNIQUENESS OF SOLUTIONS

In order to prove existence and uniqueness of solutions to (16),(17), we have to distinguish between the cases $n = 2$ and $n = 3$, due to the particularly degenerate structure of the problem for $n = 2$. In a first step, we consider the dynamics of (29) in charts K_1 and K_2 separately, which we then combine to obtain the main result of this paper:

Theorem 4.1. *For $\varepsilon \in (0, \varepsilon_0]$, with $\varepsilon_0 > 0$ sufficiently small and $n = 2, 3$, there exists a locally unique solution to (16),(17) close to the singular orbit Γ .*




FIGURE 4. Geometry in chart K_1 for (a) $n = 3$ and (b) $n = 2$.

4.1. The case $n = 3$.

4.1.1. *Dynamics in chart K_1 .* Let \mathcal{V}_1 denote the manifold $\bar{\mathcal{V}}$ in K_1 , i.e., let

$$(49) \quad \mathcal{V}_1 := \{(0, v_1, 1, \varepsilon_1) \mid |v_1 - 1| \leq \alpha, \varepsilon_1 \in [0, \varepsilon_0]\}$$

for some $0 < \alpha < 1$. To obtain the singular orbit $\bar{\Gamma}$ in K_1 , note that $\varepsilon = 0$ in (29) implies $\varepsilon_1 = 0$ in (36) due to $\varepsilon = r_1 \varepsilon_1$ and $r_1 > 0$ in (49). In general, given the initial conditions³

$$(50) \quad (u_1, v_1, r_1, \varepsilon_1)^T(0) = (0, v_{1_0}, 1, 0)^T,$$

equations (36) can easily be solved explicitly:

$$(51) \quad (u_1, v_1, r_1, \varepsilon_1)^T(\xi_1) = (v_{1_0} (1 - e^{-\xi_1}), v_{1_0} e^{-\xi_1}, e^{-\xi_1}, 0)^T.$$

Let $\gamma_1(\xi_1)$ now denote the orbit corresponding to $P_1 = (0, 1, 1, 0)$, i.e., to $v_{1_0} = 1$ in (51),

$$(52) \quad \gamma_1(\xi_1) := \{(1 - e^{-\xi_1}, e^{-\xi_1}, e^{-\xi_1}, 0) \mid \xi_1 \in [0, \infty)\},$$

and let $P_1^{in} := \gamma_1 \cap \Sigma_1^{in}$; note that $\gamma_1(\xi_1) \rightarrow Q_1 = (1, 0, 0, 0)$ as $\xi_1 \rightarrow \infty$. In analogy to Section 3, we thus define Γ_1 by

$$(53) \quad \Gamma_1 := \gamma_1 \cup \{Q_1\} \cup \{(1, 0, 0, \varepsilon_1) \mid \varepsilon_1 \in (0, \infty)\},$$

see Figure 6. As the *variational equations* of (36) along γ_1 are given by

$$(54) \quad \begin{aligned} \delta u'_1 &= \delta v_1, \\ \delta v'_1 &= -\delta v_1 - e^{-\xi_1} (1 - e^{-\xi_1}) \delta \varepsilon_1, \\ \delta r'_1 &= -\delta r_1, \\ \delta \varepsilon'_1 &= \delta \varepsilon_1, \end{aligned}$$

we obtain the following result:

Proposition 4.2. *Let $T_{P_1} \mathcal{V}_1$ denote the tangent space to \mathcal{V}_1 at P_1 , let $\mathbf{t}_{P_1} \in T_{P_1} \mathcal{V}_1$ be the tangent direction spanned by*

$$(55) \quad (\delta u_1, \delta v_1, \delta r_1, \delta \varepsilon_1)^T(0) = (0, 1, 0, 0)^T,$$

and let $\mathbf{t}_{\gamma_1} \in T_{\gamma_1} \mathcal{M}_1$ be the solution of (54) corresponding to (52). Then, $\mathbf{t}_{P_1^{in}} \in T_{P_1^{in}} \mathcal{M}_1$ is given by

$$(56) \quad (\delta u_1, \delta v_1, \delta r_1, \delta \varepsilon_1)^T(-\ln \rho) = (1 - \rho, \rho, 0, 0)^T,$$

where ρ is as in the definition of Σ_1^{in} .


Proof. For the proof, note that clearly $\delta r_1 \equiv 0 \equiv \delta \varepsilon_1$. The equations in (54) then reduce to

$$(57) \quad \begin{aligned} \delta u'_1 &= \delta v_1, \\ \delta v'_1 &= -\delta v_1, \end{aligned}$$

which can be solved to give

$$(58) \quad (\delta u_1, \delta v_1)^T(\xi_1) = (1 - e^{-\xi_1}, e^{-\xi_1})^T.$$

³It is no restriction to set $\xi_{1_0} = 0$ here; indeed, this can always be achieved by choosing the integration constant in $\xi_1(\xi) = \int r_1(\xi) d\xi$ appropriately.

FIGURE 6. Dynamics in chart K_1 for $n = 3$.

From (52), it follows that

$$(59) \quad \rho = e^{-\xi_1}$$


in Σ_1^{in} , which completes the proof. \square

Remark 6. For reasons which will become clear later on, the evolution of the tangent direction to \mathcal{V}_1 spanned by $(\delta u_1, \delta v_1, \delta r_1, \delta \varepsilon_1)^T(0) = (0, 0, 0, 1)^T$ is of no relevance to us and thus is not considered here. \square

The analysis of the transition of \mathcal{M}_1 from Σ_1^{in} to Σ_1^{out} past the line ℓ_1 of partially hyperbolic equilibria is more subtle. For hyperbolic equilibria, normal form transformations combined with cut-off techniques can be used to eliminate higher-order terms, see [Ste58]. For partially hyperbolic equilibria satisfying certain non-resonance conditions, a transformation to standard form can still be found, see [Tak71] or [Bon96]. By Lemma 3.3, however, the eigenvalues of (37) are obviously in *resonance*, both for $n = 3$ and for $n = 2$. Hence, the above techniques do not apply. We therefore have to proceed directly, i.e., by estimation, to obtain bounds on u_1 and v_1 in Σ_1^{out} . In fact, it is these resonances which are responsible for the occurrence of logarithmic switchback terms in the Lagerstrom model. This will become more evident in the upcoming paper [PS], where asymptotic expansions for the solution to (16),(17), as specified in Theorem 4.1, will be derived.

Remark 7. Note that for n irrational in (29), the resonances are destroyed, which explains the absence of logarithmic switchback in Lagerstrom's model then. \square

The following simple observation will prove quite useful:


 FIGURE 7. Evolution of \mathbf{t}_{P_1} under the flow of (54).

Lemma 4.3. *For any $u_{10}, v_{10} \geq 0$, the solutions $u_1(\xi_1)$ and $v_1(\xi_1)$ to (36) can be estimated as follows:*

$$(60) \quad \begin{aligned} u_{10} &\leq u_1(\xi_1) \leq u_{10} + v_{10} (1 - e^{-\xi_1}), \\ 0 &\leq v_1(\xi_1) \leq v_{10} e^{-\xi_1}. \end{aligned}$$

Proof. For⁴ $v_0 \geq 0$, $v(\xi) \geq 0$ follows from the invariance of $\{v = 0\}$ in (36). As

$$(61) \quad v' = -v - \varepsilon uv \leq -v,$$

integration yields $v(\xi) \leq v_0 e^{-\xi}$. Similarly, the estimate for $u(\xi)$ is obtained by integrating $0 \leq u' \leq v_0 e^{-\xi}$. \square

Proposition 4.2 asserts that \mathcal{M}_1 is very much tilted in the direction of u_1 by the flow of (36): despite \mathbf{t}_{P_1} being vertical, $\mathbf{t}_{P_1^{in}}$ is almost horizontal already, as δv_1 is almost annihilated during transport, whereas δu_1 is hugely expanded, see Figure 7. The next result shows that the transition from Σ_1^{in} to Σ_1^{out} only serves to make the tilt more pronounced, with δv_1 being even further contracted at the expense of δu_1 :

Proposition 4.4. *Let $\Pi : \Sigma_1^{in} \rightarrow \Sigma_1^{out}$ denote the transition map for (36), and let $P_1^{out} := \Pi(P_1^{in})$. Then, $D\Pi(\mathbf{t}_{P_1^{in}}) = \mathbf{t}_{P_1^{out}}$ is spanned by*

$$(62) \quad (\delta u_1^{out}, \delta v_1^{out}, \delta r_1^{out}, \delta \varepsilon_1^{out})^T(\infty) = (1, 0, 0, 0)^T.$$

Remark 8. Strictly speaking, there is of course no transition at all past ℓ_1 for $\varepsilon_1 = 0$; hence, Π and $D\Pi$ have to be defined by taking the limit as $\varepsilon_1 \rightarrow 0$. The following proof will show that this limit is in fact well-defined. \square

⁴Here and in most of the following proofs, we will omit the subscripts for the sake of readability.

Proof of Proposition 4.4. For $0 < \varepsilon^{in} < \frac{\varepsilon}{\rho}$, let $\tilde{P}^{in} := (\tilde{u}^{in}, \tilde{v}^{in}, \rho, \varepsilon^{in}) \in \mathcal{M} \cap \Sigma^{in}$, and let

$$(63) \quad \tilde{\Gamma}(\xi) := (\tilde{u}(\xi), \tilde{v}(\xi), \rho e^{-\xi}, \varepsilon^{in} e^{\xi})$$

be the solution to (36) starting in \tilde{P}^{in} (see Figure 8). The variational equations of (36) along $\tilde{\Gamma}$ are given by

$$(64) \quad \begin{aligned} \delta u' &= \delta v, \\ \delta v' &= -\varepsilon^{in} e^{\xi} \tilde{v} \delta u + (-1 - \varepsilon^{in} e^{\xi} \tilde{u}) \delta v - \tilde{u} \tilde{v} \delta \varepsilon, \\ \delta r' &= -\delta r, \\ \delta \varepsilon' &= \delta \varepsilon, \end{aligned}$$

with initial conditions in $T_{\tilde{P}^{in}} \mathcal{M}$, i.e.,

$$(65) \quad (\delta u, \delta v, \delta r, \delta \varepsilon)^T(0) = (\delta u^{in}, \delta v^{in}, 0, 0)^T.$$

As before, $\delta r \equiv 0 \equiv \delta \varepsilon$, and we obtain

$$(66) \quad \begin{aligned} \delta u' &= \delta v, \\ \delta v' &= -\delta v - \varepsilon^{in} e^{\xi} (\tilde{v} \delta u + \tilde{u} \delta v). \end{aligned}$$

To prove our assertion, we proceed by substituting (66) into

$$(67) \quad \left(\frac{\delta u}{\delta v} \right)' = \frac{\delta u' \delta v - \delta u \delta v'}{(\delta v)^2},$$

whence

$$(68) \quad z' = 1 + (1 + \varepsilon^{in} e^{\xi} \tilde{u}) z + \varepsilon^{in} e^{\xi} \tilde{v} z^2;$$

here, we have set $z := \frac{\delta u}{\delta v}$. From Lemma 4.3, we conclude that $\tilde{u}, \tilde{v} \geq 0$. With $z^{in} = \left(\frac{\delta u}{\delta v} \right)^{in} > 0$ for ε^{in} sufficiently small, this gives $z' \geq 1$ as long as z remains bounded, i.e., as long as $\delta v > 0$; therefore,

$$(69) \quad z(\xi) \geq \xi + z^{in}.$$

A similar argument for $w = z^{-1}$ shows that z indeed cannot become unbounded for finite ξ . As (63) yields

$$(70) \quad \delta = \varepsilon^{in} e^{\xi}$$

in Σ^{out} , the assertion now follows from

$$(71) \quad \left(\frac{\delta u}{\delta v} \right)^{out} \geq \ln \frac{\delta}{\varepsilon^{in}},$$

with $\varepsilon^{in} \rightarrow 0$. □

4.1.2. Dynamics in chart K_2 . Let $Q_2 = (1, 0, 0, r_2)$ in chart K_2 . The following observation is crucial for everything that follows:

Lemma 4.5. *Let $k \in \mathbb{N}$ be arbitrary.*

- (1) *There exists an attracting three-dimensional center manifold \mathcal{W}_2^c for (43) which is given by $\{v_2 = 0\}$.*
- (2) *For $|u_2 - 1|$, v_2 , η_2 , and r_2 sufficiently small, there is a stable invariant \mathcal{C}^k -smooth foliation \mathcal{F}_2^s with base \mathcal{W}_2^c and one-dimensional \mathcal{C}^k -smooth fibers.*

To obtain an approximation to \mathcal{W}_2^s through its tangent bundle $T\mathcal{W}_2^s$ along γ_2 , we consider the variational equations of (43). The latter are given by

$$(76) \quad \begin{aligned} \delta u'_2 &= \delta v_2, \\ \delta v'_2 &= -\left(\frac{2}{\xi_2} + 1\right) \delta v_2, \\ \delta \eta'_2 &= -\frac{2}{\xi_2} \delta \eta_2, \\ \delta r'_2 &= 0, \end{aligned}$$

where $' = \frac{d}{d\xi_2}$, as before. Note that the first and second equation from (76) combined yield

$$(77) \quad \delta u''_2 = -\left(\frac{2}{\xi_2} + 1\right) \delta u'_2;$$

this equation, which is precisely the (linear) *Oseen equation* from classical theory, has the solution

$$(78) \quad \delta u_2(\xi_2) = \alpha E_2(\xi_2) + \beta,$$

where

$$(79) \quad E_k(z) := \int_z^\infty e^{-t} t^{-k} dt, \quad z \in \mathbb{C}, \Re(z) > 0, k \in \mathbb{N},$$

and α, β are some constants which are as yet undetermined.⁵ From Lemma 3.4, we know that the tangent direction $\mathbf{t}_{Q_2} \in T_{Q_2}\mathcal{W}_2^{ss}$ to \mathcal{W}_2^{ss} is spanned by the vector

$$(80) \quad (\delta u_2, \delta v_2, \delta \eta_2, \delta r_2)^T(\infty) = (-1, 1, 0, 0)^T.$$

The following proposition describes the evolution of $\mathbf{t}_{\Gamma_2} \in T_{\Gamma_2}\mathcal{W}_2^s$, i.e., of \mathbf{t}_{Q_2} extended along Γ_2 as $\xi_2 \rightarrow 0$ (see Figure 9):

Proposition 4.7. *Let Q_2, \mathcal{W}_2^s , and \mathbf{t}_{Γ_2} be defined as above, and let $Q_2^{in} := \gamma_2 \cap \Sigma_2^{in}$. Then, $\mathbf{t}_{Q_2^{in}} \in T_{Q_2^{in}}\mathcal{W}_2^s$ is spanned by*

$$(81) \quad (\delta u_2, \delta v_2, \delta \eta_2, \delta r_2)^T(\delta) = (\delta u_2^{in}, 1, 0, 0)^T,$$

where

$$(82) \quad \delta u_2^{in} = \mathcal{O}(\delta).$$

Proof. As we know the solution to

$$(83) \quad \begin{aligned} \delta u' &= \delta v, \\ \delta v' &= -\left(\frac{2}{\xi} + 1\right) \delta v \end{aligned}$$


to be

$$(84) \quad (\delta u, \delta v)^T(\xi) = \left(\alpha \int_\xi^\infty e^{-t} t^{-2} dt + \beta, -\alpha e^{-\xi} \xi^{-2} \right)^T,$$

we can now determine α and β from the condition that

$$(85) \quad \lim_{\xi \rightarrow \infty} \frac{\delta u}{\delta v}(\xi) = \lim_{\xi \rightarrow \infty} \left[-e^\xi \xi^2 \int_\xi^\infty e^{-t} t^{-2} dt - \frac{\beta}{\alpha} e^\xi \xi^2 \right] = -1.$$

⁵Here, $\Re(z)$ denotes the real part of z .


 FIGURE 9. Geometry and dynamics in chart K_2 .

An easy application of *de l'Hôpital's rule* to the first term in (85) shows that we have to require $\beta = 0$; to fix α , we demand that $\delta v = 1$ in section Σ^{in} . Reverting to our original subscripts and recalling that $\xi_2 = \eta_2^{-1}$, in Σ_2^{in} we therefore have $\xi_2 = \delta$, so that $P_2^{in} = \gamma_2(\delta)$ and

$$(86) \quad \alpha = -e^\delta \delta^2.$$

For δu_2 , the substitution $\tau = \frac{t}{\delta}$ thus yields

$$(87) \quad \delta u_2 = -e^\delta \delta^2 \int_\delta^\infty e^{-t} t^{-2} dt = -e^\delta \delta \int_1^\infty e^{-\delta\tau} \tau^{-2} d\tau.$$

This latter integral, which we denote by

$$(88) \quad \tilde{E}_k(z) := \int_1^\infty e^{-z\tau} \tau^{-k} d\tau, \quad z \in \mathbb{C}, \Re(z) > 0, k \in \mathbb{N},$$

and its properties are well known, see e.g. [AS64]:

Lemma 4.8. For $|\arg z| < \pi$, $\tilde{E}_k(z)$ has the series expansions

$$(89) \quad \tilde{E}_1(z) = -\gamma - \ln z - \sum_{i=1}^{\infty} \frac{(-z)^i}{i \cdot i!}$$

for $k = 1$ and

$$(90) \quad \tilde{E}_k(z) = \frac{(-z)^{k-1}}{(k-1)!} [-\ln z + \psi(k)] - \sum_{\substack{i=0 \\ i \neq k-1}}^{\infty} \frac{(-z)^i}{(i-k+1) \cdot i!}$$

for $k \geq 2$; here, $\gamma = 0.5772\dots$ is Euler's constant and

$$(91) \quad \psi(1) = -\gamma, \quad \psi(k) = -\gamma + \sum_{i=1}^{k-1} \frac{1}{i}, \quad k \geq 2.$$

For $n = 3$, we thus obtain

$$(92) \quad \tilde{E}_2(z) = z [\ln z + \gamma - 1] - \sum_{\substack{i=0 \\ i \neq 1}}^{\infty} \frac{(-z)^i}{(i-1) \cdot i!},$$

and the proof is completed by substituting (92) into (87) and collecting powers of δ . \square

4.2. The case $n = 2$.

4.2.1. Dynamics in chart K_1 . As indicated already, our analysis in this section will have to differ substantially from the above, which is due to the complicated nature of the singular orbit Γ_1 for $n = 2$, as compared to $n = 3$. With

$$(93) \quad \gamma_1(\xi_1) := \{(0, 0, e^{-\xi_1}, 0) \mid \xi_1 \in [0, \infty)\}$$

denoting again the orbit through $P_1 = (0, 0, 1, 0)$ which is asymptotic to the origin, Γ_1 is given by

$$(94) \quad \Gamma_1 := \gamma_1 \cup \{O\} \cup \{(u_1, 0, 0, 0) \mid u_1 \in (0, 1)\} \cup \{Q_1\} \cup \{(1, 0, 0, \varepsilon_1) \mid \varepsilon_1 \in (0, \infty)\}$$

(see Figure 10). However, rather than investigating (36) for $\varepsilon_1 = 0$, as before, we now consider (36), with the perturbative terms $-\varepsilon_1 u_1 v_1$ omitted:

$$(95) \quad \begin{aligned} u_1' &= v_1, \\ v_1' &= 0, \\ r_1' &= -r_1, \\ \varepsilon_1' &= \varepsilon_1; \end{aligned}$$

here, the initial conditions are given by

$$(96) \quad (u_1, v_1, r_1, \varepsilon_1)^T(0) = (0, v_{1_0}, 1, \varepsilon_{1_0})^T.$$

The reason for considering (95) instead of (36) is that these equations can easily be solved, yielding

$$(97) \quad (u_1, v_1, r_1, \varepsilon_1)^T(\xi_1) = (v_{1_0} \xi_1, v_{1_0}, e^{-\xi_1}, \varepsilon_{1_0} e^{\xi_1})^T;$$


in a second step, we will prove that (97) is in fact a good approximation to the corresponding solution to (36), which justifies our approach. Note that due to (34), we have $\varepsilon_{1_0} = \varepsilon$ in (97).⁶ The manifold \mathcal{V}_1 of boundary conditions is given by

$$(98) \quad \mathcal{V}_1 := \{(0, v_1, 1, \varepsilon_1) \mid 0 \leq v_1 \leq \alpha, \varepsilon_1 \in [0, \varepsilon_0]\},$$

with $0 < \alpha < 1$. First, we show that for some suitable set $U_1^{out} \subset \Sigma_1^{out}$ containing $P_1^{out} := \Gamma_1 \cap \Sigma_1^{out}$ and any point $\tilde{P}_1^{out} \in U_1^{out}$, we can choose a $\tilde{P}_1 \in \mathcal{V}_1$ such that there is a solution of (95) passing through \tilde{P}_1^{out} .⁷

⁶In the following, we will use the two synonymously whenever there is no danger of confusion.

⁷Note that the size of U_1^{out} is restricted merely by the values of α and ε_0 in the definition of \mathcal{V}_1 .

FIGURE 10. Geometry in chart K_1 for $n = 2$.

Lemma 4.9. *There exists a set $U_1^{out} \subset \Sigma_1^{out}$ (specified in the proof below) such that for any $\tilde{P}_1^{out} \in U_1^{out}$, there is a unique $\tilde{P}_1 \in U_1 \subset \mathcal{V}_1$, with*

$$(99) \quad \tilde{P}_1^{out} \in \tilde{\Gamma}_1$$

for the solution $\tilde{\Gamma}_1(\xi_1)$ of (95) starting in \tilde{P}_1 ; here, U_1 is an appropriately defined subset of \mathcal{V}_1 containing P_1 .

Proof. Let $\tilde{P}^{out} := (\tilde{u}^{out}, \tilde{v}^{out}, \varepsilon\delta^{-1}, \delta)$, and let $U^{out} \in \Sigma^{out}$ be defined such that $|\tilde{u}^{out} - 1| \leq \beta$ for some $\beta > 0$ to be determined. From (97), we have

$$(100) \quad \delta = \varepsilon e^\xi$$

in Σ^{out} , whence

$$(101) \quad v_0 = \tilde{u}^{out} \left(\ln \frac{\delta}{\varepsilon} \right)^{-1} \in \left[(1 - \beta) \left(\ln \frac{\delta}{\varepsilon} \right)^{-1}, (1 + \beta) \left(\ln \frac{\delta}{\varepsilon} \right)^{-1} \right];$$

here, β is chosen such that $(1 + \beta) \left(\ln \frac{\delta}{\varepsilon_0} \right)^{-1} \leq \alpha$. The same is true of \tilde{v}^{out} , which, together with $\varepsilon \in [0, \varepsilon_0]$, determines both U^{out} and U , see Figure 10. \square

Let us fix $\tilde{u}_1^{out} = 1$ in the definition of \tilde{P}_1^{out} now, and take $\tilde{\Gamma}_1$ to be the corresponding solution to (95). Let $\tilde{P}_1^{in} := \tilde{\Gamma}_1 \cap \Sigma_1^{in}$, as before. With (95) being linear, the

Proposition 4.12. *Let $\tilde{\Pi} : \Sigma_1^{in} \rightarrow \Sigma_1^{out}$ denote the transition map for (95), and let \tilde{P}_1^{out} be defined as above. Then, $D\tilde{\Pi}(\mathbf{t}_{\tilde{P}_1^{in}}) = \mathbf{t}_{\tilde{P}_1^{out}}$ is spanned by*

$$(105) \quad (\delta u_1, \delta v_1, \delta r_1, \delta \varepsilon_1)^T \left(\ln \frac{\delta}{\varepsilon} \right) = \left(\ln \frac{\delta}{\varepsilon}, 1, 0, 0 \right)^T.$$

Proof. Equations (102) can be solved explicitly: from Proposition 4.10, one easily obtains $\delta r \equiv 0 \equiv \delta \varepsilon$ and

$$(106) \quad (\delta u, \delta v)^T(\xi) = (\xi, 1)^T.$$

The assertion then follows by taking $\xi = \ln \frac{\delta}{\varepsilon}$. \square

It only remains to show that, for ε small, the solution of the full problem (36) starting in \tilde{P}_1 does indeed stay close to $\tilde{\Gamma}_1$:

Lemma 4.13. *Let $u_{1_0} = 0$, let v_{1_0} be defined as in Lemma 4.9, and let u_1^{out} and v_1^{out} denote the values of u_1 and v_1 in Σ_1^{out} for the corresponding solution to (36). Then,*

$$(107a) \quad 1 - \varepsilon \ln \frac{\delta}{\varepsilon} \leq u_1^{out} \leq 1,$$

$$(107b) \quad \left(\ln \frac{\delta}{\varepsilon} \right)^{-1} - \varepsilon \leq v_1^{out} \leq \left(\ln \frac{\delta}{\varepsilon} \right)^{-1}.$$

Proof. For the proof, we first note that

$$(108a) \quad u(\xi) = u(\xi_0) + \int_{\xi_0}^{\xi} v(\xi') d\xi',$$

$$(108b) \quad v(\xi) = v(\xi_0) - \int_{\xi_0}^{\xi} \varepsilon(\xi') u(\xi') v(\xi') d\xi'$$

for any $0 \leq \xi_0 \leq \xi \leq \ln \frac{\delta}{\varepsilon}$. The upper bounds are obtained directly from Lemma 4.11, with $\xi_0 = 0$ and $\xi = \ln \frac{\delta}{\varepsilon}$; for the lower bounds, we estimate (108b) as

$$(109) \quad v(\xi) \geq v(\xi_0) - \varepsilon \int_{\xi_0}^{\xi} e^{\xi'} v(\xi') d\xi' \\ = (1 - e^{\xi_0}) v(\xi_0) + e^{\xi_0} v(\xi_0) - \varepsilon \int_{\xi_0}^{\xi} e^{\xi'} v(\xi') d\xi';$$

here, we have used that $\max_{\xi' \in [\xi_0, \xi]} u(\xi') \leq 1$ and $\varepsilon(\xi) = \varepsilon e^{\xi}$. To complete the proof, we require the following generalization of *Gronwall's inequality*, see [Bee75] or [Gol69]:

Lemma 4.14. *Let the real-valued functions $y(t)$ and $k(t)$ be continuous on $I := [\alpha, \beta] \subset \mathbb{R}$, and let the functions $b(t)$ and $k(t)$ be non-negative on I . If $x(t)$ is any function such that*

$$(110) \quad x(t) \geq y(t_0) - b(t) \int_{t_0}^t k(\tau) y(\tau) d\tau, \quad \alpha \leq t_0 \leq t \leq \beta,$$

then

$$(111) \quad x(t) \geq y(t_0) \exp \left[-b(t) \int_{t_0}^t k(\tau) d\tau \right], \quad \alpha \leq t_0 \leq t \leq \beta.$$

This result is optimal in the sense that equality in (110) implies equality in (111).

Setting $y(t) = e^t v(t)$, $k(t) \equiv 1$, and $b(t) \equiv \varepsilon$, we obtain

$$(112) \quad v(\xi) \geq (1 - e^{\xi_0}) v(\xi_0) + e^{\xi_0} v(\xi_0) e^{-\varepsilon(\xi - \xi_0)}$$

(with ξ instead of t as well as $\alpha = 0$ and $\beta = \ln \frac{\delta}{\varepsilon}$), whence

$$(113) \quad v^{out} \geq v_0 e^{-\varepsilon \ln \frac{\delta}{\varepsilon}} \geq v_0 \left(1 - \varepsilon \ln \frac{\delta}{\varepsilon}\right).$$

The estimate for u^{out} now follows from (108a) and (113). \square

Remark 10. One easily sees that the above estimates are equivalent to

$$(114) \quad u_1^{out} = 1 + \mathcal{O}(\varepsilon \ln \varepsilon), \quad v_1^{out} = \mathcal{O}((\ln \varepsilon)^{-1}).$$

A similar result might generally be expected if Lemma 4.13 were to be rephrased in terms of $\Pi : \Sigma_1^{in} \rightarrow \Sigma_1^{out}$, the transition map for (36). \square

4.2.2. Dynamics in chart K_2 . In contrast to the situation in K_1 , the dynamics in K_2 is no more involved for $n = 2$ than it was for $n = 3$. We will therefore not go into too many details here. Given Lemma 4.5, which is equally valid for $n = 2$, the variational equations along γ_2 are found to be

$$(115) \quad \begin{aligned} \delta u'_2 &= \delta v_2, \\ \delta v'_2 &= - \left(\frac{1}{\xi_2} + 1 \right) \delta v_2, \\ \delta \eta'_2 &= - \frac{2}{\xi_2} \delta \eta_2, \\ \delta r'_2 &= 0, \end{aligned}$$

where γ_2 is defined as in (74). The solution to

$$(116) \quad \delta u''_2 = - \left(\frac{1}{\xi_2} + 1 \right) \delta u'_2$$

is now given by

$$(117) \quad \delta u_2(\xi_2) = \alpha E_1(\xi_2) + \beta,$$

with α, β constant. Just as for $n = 3$, we have the following result:

Proposition 4.15. *Let Q_2 , \mathcal{W}_2^s , \mathfrak{t}_{Γ_2} , and Q_2^{in} be defined as above. Then, $\mathfrak{t}_{Q_2^{in}} \in T_{Q_2^{in}} \mathcal{W}_2^s$ is spanned by*

$$(118) \quad (\delta u_2, \delta v_2, \delta \eta_2, \delta r_2)^T(\delta) = (\delta u_2^{in}, 1, 0, 0)^T,$$

where

$$(119) \quad \delta u_2^{in} = \mathcal{O}(\delta \ln \delta).$$

Proof. The proof is exactly the same as for $n = 3$. \square




FIGURE 12. Illustration of transversality in Σ_2^{in} for (a) $n = 3$ and (b) $n = 2$.

5. PROOF OF THEOREM 4.1

Having finished the preparatory work, we are now ready to prove our main result:

Proof of Theorem 4.1. As noted before, it suffices to prove existence and uniqueness in the blown-up system. For (16),(17) proper, the assertion then follows by applying the appropriate blow-down transformations.

For $n = 3$, a direct computation using Lemma 3.5 and Proposition 4.4 shows that $\mathbf{t}_{P_1^{out}}$ corresponds to the vector $\mathbf{t}_{Q_2^{in}} \in T_{Q_2^{in}} \mathcal{M}_2$ spanned by

$$(120) \quad (\delta u_2, \delta v_2, \delta \eta_2, \delta r_2)^T(\delta) = (1, 0, 0, 0)^T;$$

here, $Q_2^{in} = \kappa_{12}(P_1^{out})$ and $\mathcal{M}_2 = \kappa_{12}(\mathcal{M}_1)$. With Proposition 4.7, this implies transversality in Σ_2^{in} for $r_2 = 0$, see Figure 12. From regular perturbation theory, Lemma 4.3, and the proof of Proposition 4.4, it now follows that \mathcal{M}_2 and \mathcal{W}_2^s still intersect transversely for $r_2 > 0$ sufficiently small.

For $n = 2$, we define $\tilde{P}_2^{in} := \kappa_{12}(\tilde{P}_1^{out})$ and $T\kappa_{12}|_{\Sigma_1^{out}}(\mathbf{t}_{\tilde{P}_1^{out}}) =: \mathbf{t}_{\tilde{P}_2^{in}} \in T_{\tilde{P}_2^{in}} \tilde{\mathcal{M}}_2$. By Proposition 4.12, $\mathbf{t}_{\tilde{P}_2^{in}} \cap \mathbf{t}_{Q_2^{in}}$ is then clearly transversal; with Lemma 4.13, the intersection remains transversal for (95) replaced by (36) (see again the proof of Proposition 4.4: the estimate in (69) is valid for $n = 2$, as well, as the relevant equation is

$$(121) \quad z' = 1 + \varepsilon^{in} e^{\xi} \tilde{u} z + \varepsilon^{in} e^{\xi} \tilde{v} z^2$$

now). □

Remark 11. The above proof shows why it suffices to consider only one tangent direction each both in $T\mathcal{M}_2$ and in $T\mathcal{W}_2^s$: as the equations in K_2 are completely independent of $r_2 = \varepsilon$, the question of transversality is reduced to two (instead of three) dimensions in Σ_2^{in} . □

The result of Theorem 4.1 can be interpreted as follows: for any value of $\varepsilon \in (0, \varepsilon_0]$, there is exactly one pair of values $(u_{2\varepsilon}, v_{2\varepsilon})$ that is singled out by the intersection of \mathcal{M}_2 and \mathcal{W}_2^s in Σ_2^{in} . (For $\varepsilon = 0$, of course, one again retrieves the singular orbits




FIGURE 13. $v_{1\epsilon}$ for (a) $n = 3$ and (b) $n = 2$.

discussed above. These pairs $(u_{2\epsilon}, v_{2\epsilon})$ form a curve parametrized by $\epsilon \in [0, \epsilon_0]$ which, after transformation to K_1 , determines a curve of boundary values in \mathcal{V}_1 , say $(0, v_{1\epsilon}, 1, \epsilon)$.⁸ It is precisely the function $v_{1\epsilon}$ which, if explicitly known, would give us the solution to (29). For an illustration of the above argument, see Figure 13. In the upcoming paper [PS], we will derive expansions for $v_{1\epsilon}$, both for $n = 3$ and for $n = 2$; as is to be expected, these expansions agree with those obtained in the literature by asymptotic matching.

Acknowledgements. The authors wish to thank Astrid Huber, Martin Krupa, and Martin Wechselberger for numerous helpful discussions. This work has been supported by the Austrian Science Foundation under grant Y 42-MAT.

REFERENCES

- [AS64] M. Abramowitz and I.A. Stegun, editors. *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*. Number 55 in National Bureau of Standards Applied Mathematical Series. United States Department of Commerce, Washington, 1964.
- [Bee75] P.R. Beesack. Gronwall inequalities. In *Carleton Mathematical Lecture Notes*, volume 11. Carleton University, Ottawa, 1975.
- [Bon96] P. Bonckaert. Partially hyperbolic fixed points with constraints. *Trans. Amer. Math. Soc.*, 348(3):997–1011, 1996.
- [Bus71] W.B. Bush. On the Lagerstrom mathematical model for viscous flow at low Reynolds number. *SIAM J. Appl. Math.*, 20(2):279–287, 1971.
- [CFL78] D.S. Cohen, A. Fokas, and P.A. Lagerstrom. Proof of some asymptotic results for a model equation for low Reynolds number flow. *SIAM J. Appl. Math.*, 35(1):187–207, 1978.
- [CLW94] S.N. Chow, C. Li, and D. Wang. *Normal Forms and Bifurcation of Planar Vector Fields*. Cambridge University Press, Cambridge, 1994.
- [Col68] J.D. Cole. *Perturbation Methods in Applied Mathematics*. Blaisdell Publishing Company, Waltham, 1968.

⁸Note that due to $r_1 = 1$ in \mathcal{V}_1 , we can again write ϵ instead of ϵ_1 .

- [DR91] Z. Denkowska and R. Roussarie. A method of desingularization for analytic two-dimensional vector field families. *Bol. Soc. Bras. Mat.*, 22(1):93–126, 1991.
- [DR96] F. Dumortier and R. Roussarie. Canard cycles and center manifolds. *Mem. Amer. Math. Soc.*, 121(577), 1996.
- [Dum93] F. Dumortier. Techniques in the Theory of Local Bifurcations: Blow-Up, Normal Forms, Nilpotent Bifurcations, Singular Perturbations. In D. Schlomiuk, editor, *Bifurcations and Periodic Orbits of Vector Fields*, number 408 in NATO ASI Series C, Mathematical and Physical Sciences, Dordrecht, 1993. Kluwer Academic Publishers.
- [Fen79] N. Fenichel. Geometric Singular Perturbation Theory for Ordinary Differential Equations. *J. Diff. Eqs.*, 31:53–98, 1979.
- [GH83] J. Guckenheimer and P. Holmes. *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, volume 42 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1983.
- [Gol69] H.E. Gollwitzer. A note on a functional inequality. *Proc. Amer. Math. Soc.*, 23:642–647, 1969.
- [Hsi73] G.C. Hsiao. Singular perturbations for a nonlinear differential equation with a small parameter. *SIAM J. Math. Anal.*, 4:283–301, 1973.
- [HTB90] C. Hunter, M. Tajdari, and S.D. Boyer. On Lagerstrom’s model of slow incompressible viscous flow. *SIAM J. Appl. Math.*, 50(1):48–63, 1990.
- [Jon95] C.K.R.T. Jones. Geometric Singular Perturbation Theory. In *Dynamical Systems*, volume 1609 of *Springer Lecture Notes in Mathematics*, New York, 1995. Springer-Verlag.
- [Kap57] S. Kaplun. Low Reynolds number flow past a circular cylinder. *J. Math. Mech.*, 6:595–603, 1957.
- [KL57] S. Kaplun and P.A. Lagerstrom. Asymptotic expansions of Navier-Stokes solutions for small Reynolds numbers. *J. Math. Mech.*, 6:585–593, 1957.
- [KS01] M. Krupa and P. Szmolyan. Extending geometric singular perturbation theory to non-hyperbolic points—fold and canard points in two dimensions. *SIAM J. Math. Anal.*, 33(2):286–314, 2001.
- [KW96] J.B. Keller and M.J. Ward. Asymptotics beyond all orders for a low Reynolds number flow. *J. Eng. Math.*, 30(1-2):253–265, 1996.
- [KWK95] M.C.A. Kropinski, M.J. Ward, and J.B. Keller. A hybrid asymptotic-numerical method for low Reynolds number flow past a cylindrical body. *SIAM J. Appl. Math.*, 55(6):1484–1510, 1995.
- [Lag66] P.A. Lagerstrom. A course on perturbation methods. Lecture Notes by M. Mortell, National University of Ireland, Cork, 1966.
- [Lag88] P.A. Lagerstrom. *Matched asymptotic expansions: ideas and techniques*, volume 76 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1988.
- [LC72] P.A. Lagerstrom and R.G. Casten. Basic concepts underlying singular perturbation techniques. *SIAM Rev.*, 14(1):63–120, 1972.
- [LR84] P.A. Lagerstrom and D.A. Reinelt. Note on logarithmic switchback terms in regular and singular perturbation expansions. *SIAM J. Appl. Math.*, 44(3):451–462, 1984.
- [Ose10] C.W. Oseen. Über die Stokes’sche Formel, und über eine verwandte Aufgabe in der Hydrodynamik. *Ark. Math. Astronom. Fys.*, 6(29), 1910.
- [PP57] I. Proudman and J.R.A. Pearson. Expansions at small Reynolds numbers for the flow past a sphere and a circular cylinder. *J. Fluid Mech.*, 2:237–262, 1957.
- [PS] N. Popović and P. Szmolyan. Rigorous asymptotic expansions for Lagerstrom’s model equation—a geometric approach. In preparation.
- [RS75] S. Rosenblat and J. Shepherd. On the asymptotic solution of the Lagerstrom model equation. *SIAM J. Appl. Math.*, 29(1):110–120, 1975.
- [Ski81] L.A. Skinner. Note on the Lagerstrom singular perturbation models. *SIAM J. Appl. Math.*, 41(2):362–364, 1981.
- [Ste58] S. Sternberg. On the structure of local homeomorphisms of Euclidean n -space, II. *Am. J. Math.*, 80:623–631, 1958.
- [Sto51] G.G. Stokes. On the effect of the internal friction of fluids on the motion of pendulums. *Trans. Camb. Phil. Soc.*, 9:8–106, 1851. Part II.
- [Tak71] F. Takens. Partially hyperbolic fixed points. *Topology*, 10:133–147, 1971.
- [vGKS] S. van Gils, M. Krupa, and P. Szmolyan. Asymptotic expansions using blow-up. To appear in *ZAMP*.

[Whi89] A.N. Whitehead. Second approximations to viscous fluid motion. *Quart. J. Math.*, 23:143–152, 1889.

E-mail address: `nikola.popovic@tuwien.ac.at`, `peter.szmolyan@tuwien.ac.at`

DEPARTMENT OF APPLIED MATHEMATICS AND NUMERICAL ANALYSIS, VIENNA UNIVERSITY OF TECHNOLOGY, WIEDNER HAUPTSTRASSE 8-10, A-1040 VIENNA, AUSTRIA, EU.