



Smooth Homology Spheres and their Fundamental Groups

Author(s): Michel A. Kervaire

Source: *Transactions of the American Mathematical Society*, Vol. 144 (Oct., 1969), pp. 67-72

Published by: American Mathematical Society

Stable URL: <http://www.jstor.org/stable/1995269>

Accessed: 04/01/2009 05:04

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ams>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Transactions of the American Mathematical Society*.

<http://www.jstor.org>

SMOOTH HOMOLOGY SPHERES AND THEIR FUNDAMENTAL GROUPS

BY
MICHEL A. KERVAIRE

Let M^n be a smooth homology n -sphere, i.e. a smooth n -dimensional manifold such that $H_*(M^n) \cong H_*(S^n)$. The fundamental group π of M satisfies the following three conditions:

- (1) π has a finite presentation,
- (2) $H_1(\pi) = 0$,
- (3) $H_2(\pi) = 0$,

where $H_i(\pi)$ denotes the i th homology group of π with coefficients in the trivial $\mathbb{Z}\pi$ -module \mathbb{Z} . Properties (1) and (2) are trivial and (3) follows from the theorem of Hopf [2] which asserts that $H_2(\pi) = H_2(M) / \rho\pi_2(M)$, where ρ denotes the Hurewicz homomorphism.

For $n > 4$ we will prove the following converse

THEOREM 1. *Let π be a group satisfying the conditions (1), (2) and (3) above, and let n be an integer greater than 4. Then, there exists a smooth manifold M^n such that $H_*(M^n) \cong H_*(S^n)$ and $\pi_1(M) \cong \pi$.*

The proof is very similar to the proof used for the characterization of higher knot groups in [5]. Compare also the characterization by K. Varadarajan of those groups π for which Moore spaces $M(\pi, 1)$ exist [9].

Not much seems to be known for $n \leq 4$. If M^3 is a 3-dimensional smooth manifold with $H_*(M) \cong H_*(S^3)$, then $\pi = \pi_1(M)$ possesses a presentation with an equal number of generators and relators. (Take a Morse function f on M with a single minimum and a single maximum. Then f possesses an equal number of critical points of index 1 and 2.)

Also, under restriction to *finite* groups there is the following

THEOREM 2. *Let M^3 be a 3-dimensional manifold such that $H_*(M) \cong H_*(S^3)$. Suppose that $\pi_1(M)$ is finite. Then, either $\pi_1(M) = \{1\}$ or else, $\pi_1(M)$ is isomorphic to the binary icosahedral group with presentation*

$$(x, y; x^2 = y^3 = (xy)^5).$$

This is implicitly well known: The hypotheses imply that $\pi = \pi_1(M)$ is a group of fix-point free transformations of a homotopy 3-sphere. Any such group belongs to a list established by Suzuki [8] and even to the shorter list of Milnor [7]. The

binary icosahedral group is the only (nontrivial) group in Milnor's list satisfying $H_1\pi=0$.

REMARK. Curiously enough, it seems that the trivial group and the binary icosahedral group are the only available examples of finite groups π satisfying $H_1\pi=0$ and having a presentation with an equal number of generators and relators.

For $n=4$, it follows from the proof below that every finitely presented group π with an equal number of generators and relators and satisfying $H_1\pi=0$ is the fundamental group of some homology 4-sphere. It seems unlikely that these conditions should characterize the fundamental groups of homology 4-spheres, but I do not know of a counterexample.

1. **Proof of Theorem 1.** We start with a finite presentation of π :

$$(x_1, \dots, x_\alpha; R_1, \dots, R_\beta)$$

and the manifold

$$M_0 = (S^1 \times S^{n-1}) \# \dots \# (S^1 \times S^{n-1}),$$

connected sum of α copies of $S^1 \times S^{n-1}$. Choosing as usual a contractible open set U in M_0 as "base point", the condition $n > 2$ implies that $\pi_1(M, U)$ is free on α generators. After identification of free generators of $\pi_1(M, U)$ with x_1, \dots, x_α , the elements R_1, \dots, R_β of the free group on x_1, \dots, x_α can be represented by disjoint differentiable imbeddings $\phi_i: S^1 \times D^{n-1} \rightarrow M_0, i=1, \dots, \beta$. Moreover, these can be chosen so that the spherical modification $\chi(\phi_1, \dots, \phi_\beta)$ can be framed (see [6]). The resulting manifold $M_1 = \chi(M_0; \phi_1, \dots, \phi_\beta)$ is stably parallelizable as was M_0 and $\pi_1(M_1) \cong \pi$ since $n > 3$.

From the homology exact sequences of the pairs $(M_0, \cup_i \phi_i(S^1 \times D^{n-1}))$ and $(M_1, \cup_i \phi'_i(D^2 \times S^{n-2}))$, where ϕ'_i denotes the natural imbedding $D^2 \times S^{n-2} \rightarrow M_1$ with $\phi'_i|S^1 \times S^{n-2} = \phi_i|S^1 \times S^{n-2}$, one concludes that $H_i(M_1) = 0$ for $i \neq 0, 2, n-2, n$, and that $H_2(M_1)$ is free abelian of rank $\gamma = \beta - \alpha$. (Observe that $\beta \geq \alpha$ since π abelianized is trivial.) Hence, there exist bases ξ_1, \dots, ξ_γ of $H_2(M_1)$ and $\eta_1, \dots, \eta_\gamma$ of $H_{n-2}(M_1)$ respectively such that $\xi_i \cdot \eta_j = \delta_{ij}$, where $\xi_i \cdot \eta_j$ is the homology intersection number, and δ_{ij} is the Kronecker delta.

By the theorem of Hopf mentioned above, $H_2\pi = H_2(M_1)/\rho\pi_2(M_1)$, and so by condition (3) on π , the Hurewicz homomorphism $\rho: \pi_2(M_1) \rightarrow H_2(M_1)$ is surjective.

Since $n > 4$, the classes ξ_1, \dots, ξ_γ can be represented by disjoint differentiable imbeddings $f_i: S^2 \rightarrow M_1, i=1, \dots, \gamma$, and M_1 being stably parallelizable, these extend to disjoint imbeddings $\psi_i: S^2 \times D^{n-2} \rightarrow M_1$. It follows from the arguments in [6, §5] that the manifold $M = \chi(M_1; \psi_1, \dots, \psi_\gamma)$ obtained by spherical modification is a homology sphere with $\pi_1(M) \cong \pi$.

2. Many of the homology spheres occurring in the literature are constructed as the boundary of a contractible manifold. We investigate this question whether, in general, homology spheres bound contractible manifolds.

The following well-known construction provides an example of a 3-dimensional homology sphere which does not bound a contractible manifold.


Let $W = D^4 + (\phi_1) + \dots + (\phi_8)$ be the manifold obtained from the 4-disc by attaching eight handles of type 2 using unknotted imbeddings (with disjoint images) $\phi_i: S^1 \times D^2 \rightarrow S^3$, such that the matrix of linking numbers

$$L(\phi_i(S^1 \times x_0), \phi_j(S^1 \times x_1))$$

with $x_0 \neq x_1, x_0, x_1 \in D^2$, is:

$$L = \begin{bmatrix} 2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 2 \end{bmatrix}$$

This matrix is unimodular and has signature 8. The picture shows the images



$S_i = \phi_i(S^1 \times 0)$ in $R^3 = S^3 - \text{point}$. From the handle decomposition of W one sees that $\pi_1 W = \{1\}$, and $H_2 W$ is free abelian on 8 generators. An easy calculation shows that $\pi_1(bW)$ is the group with presentation:

$$(x, y; x^5 = y^2 = (x^{-1}y)^3),$$

where x and y are the classes of the loops shown on the picture. Hence, bW is a homology sphere. If bW were the boundary of a contractible manifold V , one could form the closed 4-manifold $M = W \cup V$, and the above matrix L would be the matrix of intersection numbers of $H_2 M$. From L one reads off the Stiefel-Whitney class $w_2 M = 0$ and the signature $\sigma(M) = 8$, contradicting Rohlin's theorem

which says that if $w_2M=0$ for a closed orientable 4-manifold M , then $\sigma(M)\equiv 0 \pmod{16}$. (Cf. [4].)

In contrast, we have

THEOREM 3. *Every 4-dimensional homology sphere bounds a contractible manifold. If M^n is a smooth oriented homology sphere with $n \geq 5$, there exists a unique smooth homotopy sphere Σ_M^n such that $M^n \# \Sigma^n$ bounds a contractible smooth manifold.*

This has been obtained independently by the Hsiang brothers [3].

Let first M^4 be a smooth 4-dimensional homology sphere which we think of as imbedded in S^{k+4} with k large. It is easy to see that M has trivial normal bundle. Indeed, the only obstruction to trivializing the normal bundle is an element $\mathcal{O} \in H^4(M; \pi_3(SO_k)) = \mathbb{Z}$. It is known that $2\mathcal{O} = \pm p_1[M]$, where p_1 is the Pontryagin class. (Cf. e.g. [4].) The latter vanishes by the Thom-Hirzebruch formula $\frac{1}{2}p_1[M] = \sigma(M) = 0$. Now, choosing a trivialization of the normal bundle of M^4 in S^{k+4} , the Thom construction provides an element in $\pi_{k+4}(S^k)$. Since $\pi_{k+4}(S^k) = 0$, we see that M^4 is the boundary of a parallelizable manifold W^5 . This manifold can be modified to get a new 5-manifold V^5 with $bV^5 = bW^5 = M^4$ such that $\pi_1V = \{1\}$ and $\pi_2V = 0$. (Cf. [6, §5].) Clearly, V is contractible.

Assume then that $n \geq 5$.

We associate with the given smooth homology n -sphere M^n some smooth homotopy sphere Σ^n . Observe that M^n is stably parallelizable. Since

$$H^i(M; \pi_{i-1}(SO)) = 0 \quad \text{for } i < n,$$

there is only one possible obstruction $\mathcal{O} \in H^n(M; \pi_{n-1}(SO)) = \pi_{n-1}(SO)$ to producing a trivialization of the stable tangent bundle of M . It is known that \mathcal{O} belongs to the kernel of the homomorphism $J: \pi_{n-1}(SO) \rightarrow \Pi_{n-1}$, where $\Pi_{n-1} = \pi_{n+k-1}(S^k)$, k large. Thus $\mathcal{O} = 0$ by the same arguments as in [6, §3].

We obtain Σ^n from M^n by framed surgery in dimensions 1 and 2. Let x_1, \dots, x_α be a finite set of generators of $\pi_1(M, U)$, where U is a contractible open "base set" in M . Let $\phi_1, \dots, \phi_\alpha$ be smooth imbeddings of $S^1 \times D^{n-1}$ into M^n with disjoint images representing x_1, \dots, x_α respectively. We use $\phi_1, \dots, \phi_\alpha$ to attach α handles of type 2 to $I \times M$ along $(1) \times M$, where $I = [0, 1]$. Let

$$V_0 = I \times M + (\phi_1) + \dots + (\phi_\alpha)$$

be the resulting $(n+1)$ -manifold. We may assume that the imbeddings $\phi_1, \dots, \phi_\alpha$ have been chosen so that V_0 is parallelizable. The manifold $N = bV_0 - (0) \times M$ will be called the right-hand boundary of V_0 . It is easily checked that $\pi_1N = \pi_1V_0 = \{1\}$, and $H_iN = 0$ for $3 \leq i \leq n-3$ (if $n \geq 6$). The groups H_2N and $H_{n-2}N$ are free abelian of rank α and the inclusion $N \subset V_0$ induces an isomorphism $H_2N \cong H_2V_0$.

Let ξ_1, \dots, ξ_α and $\eta_1, \dots, \eta_\alpha$ be bases of H_2N and $H_{n-2}N$ respectively such that $\xi_i \cdot \eta_j = \delta_{ij}$. Then, representing the classes ξ_1, \dots, ξ_α by disjoint differentiable imbeddings $\psi_1, \dots, \psi_\alpha$ of $S^2 \times D^{n-2}$ into N we construct

$$V_1 = I \times M + (\phi_1) + \dots + (\phi_\alpha) + (\psi_1) + \dots + (\psi_\alpha)$$

by attaching α handles of type 3 to V_0 along N using $\psi_1, \dots, \psi_\alpha$. Again this can be done so that V_1 is parallelizable. Now, the right-hand boundary $bV_1 - (0) \times M$ is a smooth homotopy n -sphere Σ^n . We think of it as oriented so that the (oriented) boundary of V_1 is $-\Sigma^n - (0) \times M$, where the orientation of V_1 is given by the one of $I \times M$.

Claim: $M \# \Sigma^n$ bounds a contractible manifold. Indeed, let $t: (I \times D^n, bI \times D^n) \rightarrow (V_1, bV_1)$ be a smooth imbedding with $t(\text{int } I \times D^n) \subset \text{int } V_1$, then after rounding off corners we get a smooth contractible manifold $V = V_1 - t(I \times \text{int } D^n)$ whose boundary is diffeomorphic to $M \# \Sigma$.

It remains to prove the uniqueness of Σ . Let Σ_1 be a homotopy n -sphere such that $M \# \Sigma_1$ is the boundary of a contractible manifold W . Using a connected sum of V_1 (as constructed above) and $I \times \Sigma_1$ along $t(I \times S^{n-1})$ and $I \times \sigma^{n-1}$, where σ^{n-1} is the boundary of a smooth n -disc δ^n in Σ_1 , one gets a manifold W' whose homotopy type is S^n , and $bW' = (-\Sigma) \# \Sigma_1 + (-M) \# \Sigma_1$. (W' is obtained from the disjoint union

$$\{V_1 - t(I \times (0))\} + \{I \times \Sigma_1 - I \times (0)\}$$

under the identification of $t(x, ry) \in t(I \times D^n)$ with $(x, (1-r)y) \in I \times \delta^n$ for $x \in I$, $y \in S^{n-1}$ and $0 < r < 1$.)

Now, paste the manifold W along the left boundary $((0) \times M) \# \Sigma_1$ of W' by the identity diffeomorphism

$$bW = M \# \Sigma_1 \rightarrow ((0) \times M) \# \Sigma_1.$$

The resulting union $W \cup W'$ is a contractible manifold as follows easily using the van Kampen and Mayer-Vietoris theorems. The boundary of $W \cup W'$ is $(-\Sigma) \# \Sigma_1$. Therefore, Σ and Σ_1 are h -cobordant, and thus diffeomorphic since we assume $\dim \Sigma = \dim \Sigma_1 \geq 5$.

COROLLARY. Every combinatorial homology sphere K^n of dimension $n \neq 3$ is the boundary of a contractible combinatorial manifold.

By Hirsch's obstruction theory [1, Theorem 3.1], every combinatorial manifold K^n admits a smoothness structure in the neighborhood of its 7-skeleton. We need a smoothness structure in a neighborhood N of the 2-skeleton of K^n . Then N is parallelizable and we can apply the above surgery arguments requiring the imbeddings to have their images in the smooth subset. We obtain a manifold W^{n+1} such that $bW = \Sigma^n + M^n$, where Σ^n is a combinatorial homotopy n -sphere and the inclusion $\Sigma \subset W$ is a homotopy equivalence. Since $n \geq 5$, Σ^n is PL-homeomorphic to $b\Delta^{n+1}$, and pasting Δ^{n+1} to W^{n+1} by such a homeomorphism provides a contractible manifold V^{n+1} with boundary M^n .

Thus, as a by-product, we have proved the probably well-known fact that every combinatorial homology sphere admits a smoothness structure.

BIBLIOGRAPHY

1. M. Hirsch, *Obstruction theories for smoothing manifolds and maps*, Bull. Amer. Math. Soc. **69** (1963), 352–356.
2. H. Hopf, *Fundamentalgruppe und zweite Bettische Gruppe*, Comment. Math. Helv. **14** (1941), 257–309.
3. Wu-chung Hsiang and Wu-yi Hsiang, *Differentiable actions of compact connected classical groups. I*, Yale University Notes, 1966.
4. M. Kervaire, *On the Pontryagin classes of certain $SO(n)$ -bundles over manifolds*, Amer. J. Math. **80** (1958), 632–638.
5. ———, *Les noeuds de dimensions superieures*, Bull. Soc. Math. France **93** (1965), 225–271.
6. M. Kervaire and J. Milnor, *Groups of homotopy spheres*, Ann. of Math. **77** (1963), 504–537.
7. J. Milnor, *Groups which act on S^n without fixed points*, Amer. J. Math. **79** (1957), 623–630.
8. M. Suzuki, *On finite groups with cyclic Sylow subgroups for all odd primes*, Amer. J. Math. **77** (1955), 657–691.
9. K. Varadarajan, *Groups for which Moore spaces $M(\pi, 1)$ exist*, Ann. of Math. **84** (1966), 368–371.

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY,
NEW YORK, NEW YORK